Homoclinic orbit to a center manifold

Patrick Bernard

janvier 2000

Introduction

A saddle-center fixed point of a Hamiltonian system is a fixed point with precisely one pair of purely imaginary eigenvalues, and other eigenvalues all having non-zero real part. Such a fixed point is contained in a two dimensional invariant manifold, called the center manifold, associated with the pair of imaginary eigenvalues and filled with periodic orbits. Each of these periodic orbits is the transversal intersection between its energy shell and the center manifold, and is hyperbolic with respect to its energy shell. We are interested in the existence of orbits homoclinic to these periodic trajectories.

Let us consider an initial system with a saddle-center fixed point and an orbit homoclinic to it. The orbit structure near this homoclinic orbit of the initial system and of perturbed systems can be studied using appropriate local sections and the Poincaré return map along the homoclinic. This has been initiated by Conley in [12], and used in [21], [23] and [26] to prove the existence of homoclinic orbits in perturbed systems. Under suitable hypotheses, and if the phase space is four dimensional, these papers show the following behavior. Hamiltonian systems sufficiently close to the initial system have a saddle-center fixed point with a center manifold. We can suppose without loss of generality that the saddle-center fixed point always has zero energy in the systems under interest. For any sufficiently small fixed positive energy, the periodic motion on the center manifold at that energy has a homoclinic orbit in a system sufficiently close to the initial system. However, in a fixed system close to the initial system, the periodic orbits closest to the fixed point are not proved to have any homoclinic orbit. The homoclinic orbits of smallest energy are first destroyed by the perturbation. This is not surprising since the saddle-center fixed point itself does not have any homoclinic orbit in general. These works provide a much more detailed description of the orbit structure than we do in this paper, but their range is limited to the study of perturbations of initial systems with a homoclinic orbit to the saddle-center, which is an exceptional case, and to four dimensional phase spaces.

Variational methods provide global existence results on homoclinic orbits to a hyperbolic fixed point, see [5], [13] and many other papers, that can be viewed as non-perturbative analogs of the theory of Melnikov that studies the persistence of homoclinics under perturbation. In the same spirit, we attempt to provide a non-perturbative analog of the behavior described above around saddle-center fixed points. This paper is closely connected to [3], where we study the existence of homoclinic orbits to some hyperbolic periodic orbits of a Hamiltonian system in $\mathbb{C}^n$. Since the smallest periodic motion in the center manifold having a homoclinic orbit seems to go away from the fixed point when the system goes away from the initial system in the perturbative setting, it is natural to consider a global center manifold



Figure 1: Elastic pendulum

in order to find a global (i.e. non-perturbative) result. In [3], we study the vicinity of a prescribed energy shell sufficiently far from the origin, and obtained homoclinic orbits in a dense family of energy shells around the prescribed one. We abandon all pretension to find many homoclinic orbits, but focus on finding the orbit closest to the saddle-center. It is yet very unlikely that the orbit we find is indeed the closest to the saddle-center, but it is probably the closest among those which satisfy a certain estimate. We shall clarify this point later.

We study a model system where the center manifold is a plane with harmonic oscillations on it. We suppose that these periodic motions are hyperbolic with respect to their energy shells, so that the center manifold is a normally hyperbolic manifold. The setting is thus quite similar to the setting of [3], but we assume here that the total phase space is the product of this plane with the cotangent bundle of a compact manifold M, instead of $\mathbb{R}^{2n}$ in [3]. This product structure is a key to our result, since we shall obtain homoclinic orbits by comparison with product (uncoupled) flows. The existence of homoclinic orbits for product flows is reduced to well-known existence results on homoclinic motions to hyperbolic fixed points on compact Riemannian manifolds, see [5]. We shall moreover assume that the Hamiltonian is fiberwise convex on the total phase space $T^*(M \times \mathbb{R})$, so that a Lagrangian action functional can be used. It should be possible to avoid this restriction since the Hamiltonian action functional can be well studied in this context, see [19] or [10]. The Lagrangian functional remains simpler, and the results of this paper will be expressed in terms of Lagrangian systems.

Let us stress that, although the setting is Lagrangian, our result is very different from classical ones on homoclinic orbits in Lagrangian systems since the periodic orbits of the center manifold do not satisfy the minimality hypothesis needed in these results. In fact, the center manifold as a whole satisfies this hypothesis, and we shall look for orbits homoclinic to this manifold. An orbit homoclinic to the center manifold is homoclinic to one of the periodic orbits, by energy conservation. The difficulty is that the center manifold is not compact, and that we have to find a way to localize orbits.

Under suitable hypotheses, we prove the existence of an orbit homoclinic to one of the oscillations of the center manifold and give an estimate of its action and of its energy. These estimates are the main novelties compared with [3], they allow interesting new applications. The energy is close to zero (the energy of the fixed point) when the system is close to a product system and the homoclinic we find should be seen as the continuation, when a coupling is introduced, of the orbit homoclinic to the origin that existed in the product system. In

this sense we can say that the homoclinic we find is the closest to the origin, although new orbits, longer and closer, may appear. Both the center manifold and the homoclinic orbit are preserved by a small perturbation of the system *i.e.* a perturbed system still has an invariant manifold diffeomorphic to a plane and foliated by periodic orbits one of which has a homoclinic.

Among the applications let us give the example of the stiff elastic spatial pendulum. This is a pendulum where the bar has been replaced by a stiff spring which has variable length but remains always straight, see Figure 1. The center manifold here is the set of oscillations of the spring in unstable equilibrium. We obtain an orbit homoclinic to one of these oscillations when the spring is stiff enough. This homoclinic is moreover preserved by a small perturbation of the system. It is a very general process to introduce an additional degree of freedom highly confined to zero in a mechanical system (a previously frozen binding is now granted some freedom to oscillate). Under certain hypotheses, we see that a hyperbolic fixed point with a homoclinic orbit of the frozen system is turned to a saddle-center fixed point with a center manifold and an orbit homoclinic to the center manifold in the extended system.

A major interest of homoclinic orbits is their link with chaotic behavior. The orbit structure near a transversal homoclinic orbit to a hyperbolic fixed point of a periodic time-dependent system has by now been well described. The natural analog of this structure exists in an autonomous system around a transversal homoclinic orbit to an hyperbolic fixed point. It should be noted however that the behavior associated with homoclinic orbits to hyperbolic fixed points of autonomous systems is not as well understood, see [15] and [8] for some results on this subject. One of the interests of our work is that the homoclinic we find, if transversal, lead to the well described case, *i.e.* to a Bernoulli shift with positive entropy. Consider for example a classical plane pendulum, our results provide a new way to break integrabilty and introduce chaotic behavior. Instead of considering that there is some small influence from the exterior (a time dependent perturbation), one can consider that the bar has some elasticity. In this case, the unstable equilibrium is surrounded by unstable oscillations. We prove that one of these oscillations have a homoclinic orbit, this homoclinic can be made transversal by a perturbation, and the system then has positive topological entropy.

1 Results, comments and applications

Let M be a compact manifold, $TM \xrightarrow{\pi} M$ its tangent bundle. We provide M with a metric g, and note

$$\|z\| = \sqrt{g_{\pi(z)}(z,z)}$$

the norm of a tangent vector $z \in TM$. There is an associated metric on TM, and we note $d(z,z')$ the distance between two points of TM associated with this metric. Let us consider the smooth Lagrangian on $T(M \times \mathbb{R}) = TM \times \mathbb{R}^2$ given by

$$L(z,q,v) = a\left(v^2 - \omega^2 q^2\right) + G(z,q,v) \quad (z,q,v) \in TM \times \mathbb{R} \times \mathbb{R}, \tag{1}$$

where a and ω are positive real numbers and $G : TM \times \mathbb{R}^2 \longrightarrow \mathbb{R}$ is a smooth function satisfying the assumptions:

HG1 : There exists a $z_0 = (\theta_0, 0) \in TM$ such that $G(z_0,q,v) = 0$ and $dG(z_0,q,v) = 0$ for all $(q,v) \in \mathbb{R}^2$.

HG2 : There exists a $b > 0$ such that $G(z,q,v) \geqslant b\, d(z,z_0)^2$.
HG3 :

$$G(z,q,v) \geqslant \frac{1}{2}\left(q\frac{\partial G}{\partial q}(z,q,v) + v\frac{\partial G}{\partial v}(z,q,v)\right) + bd(z,z_0)^2.$$

Moreover, we assume that there exist two smooth fiberwise convex functions U and W on TM such that

$$U(z) \leqslant G(z,q,v) \leqslant W(z) \tag{2}$$

for all $(z,q,v) \in TM \times \mathbb{R}^2$, and that they both satisfy
HU1 : $U(z_0) = 0$ and $dU(z_0) = 0$,
HU2 : $U(z) \geqslant b\, d(z,z_0)^2$.
Finally, we assume that the Lagrangian L is fiberwise convex,
HL : The restriction of L to each fiber $T_\theta M \times T_q\mathbb{R}$ is convex.
No more control at infinity is necessary for our results to hold true, but we will use in the proofs systems satisfying the additional hypothesis
HG4 : There exists a function G_∞ on TM, a number $\alpha > 0$, a compact set $K \subset TM$ and a compact set $B \subset K \times \mathbb{R}^2$ such that $G(z,q,v) = G_\infty(z)$ outside B, and $G_\infty(z) = \alpha\|z\|^2$ outside K.
As a consequence of [HL], the trajectories of L on $M \times \mathbb{R}$ are the projections of the integral curves of a vector-field Y_L on TM, that is conjugated to the Hamiltonian vector field X_H on T^*M, where H is the fiberwise dual of L

$$H(\zeta,q,p) = \sup_{z\in\pi^{-1}(\pi^*(\zeta)),v\in\mathbb{R}} \langle\zeta,z\rangle + pv - L(z,q,v) \quad (\zeta,q,p) \in T^*M \times \mathbb{R}^2.$$

See Section 2 for more details. The flow of Y_L has an invariant manifold, the center manifold, of equation $z = z_0$. The center manifold is filled with periodic orbits, which are the liftings of

$$O_r(t) = (\theta_0, r\cos(\omega t)),$$

and can be described also by

$$O_r = \{(z_0,q,v) \in TM \times \mathbb{R}^2 \,/\, v^2 + \omega^2 q^2 = \omega^2 r^2\}.$$

We are looking for orbits homoclinic to O_r, i.e. trajectories $x = (\theta, q) : \mathbb{R} \longrightarrow M \times \mathbb{R}$ such that $\theta \not\equiv \theta_0$ and

$$\lim_{t\to\pm\infty}\left(\theta(t), \dot\theta(t), \dot q(t)^2 + \omega^2 q(t)^2\right) = \left(\theta_0, 0, \omega^2 r^2\right).$$

We will see in Section 5 that to any function U on TM satisfying [HU1,2] we can associate a number I(U) such that

$$U \leqslant W \Longrightarrow I(U) \leqslant I(W)$$

and

$$\left|1 - \frac{I(W)}{I(U)}\right| \leqslant \sup_z \frac{|W(z) - U(z)|}{b\, d^2(z,z_0)} \tag{3}$$

for all U and W satisfying [HU1-2]. Recall that b is the constant of [HU2]. The value $I(U)$ can be thought of as the action of an orbit homoclinic to z_0 for the Lagrangian system U on TM, although we can only prove that there is a homoclinic of action below $I(U)$. We are now in a position to state the main result of this paper:

Theorem 1 *Let us consider the Lagrangian system (1), and assume that G satisfies [HG1-3] and*

$$U(z) \leqslant G(z,q,v) \leqslant W(z)$$

with U and W satisfying [HU1,2]. There is a radius

$$r \leqslant \sqrt{\frac{I(W) - I(U)}{2\pi a \omega}} \tag{4}$$

such that the periodic or

4. Let E be the energy, see Section 2. It can be easily computed (see Section 6) that

$$E(O_r) = a\omega^2 r^2,$$

and the energy of the homoclinic obtained from Theorem 1 satisfies

$$0 \leqslant E(\partial X_\infty) \leqslant E_0 = \frac{\omega}{2\pi}\big(I(W) - I(U)\big).$$

5. The integral $\int_{\mathbb{R}} L(\partial X)$ is not defined for a homoclinic orbit because it has an oscillating tail. This is linked to the fact that the action

$$\int_0^t L(\partial O_r(s))\, ds$$

is not identically zero. We can nevertheless integrate by parts the expression

$$\int \dot{q}^2 + \omega^2 q^2 = [q\dot{q}] - \int q(\ddot{q} - \omega^2 q),$$

and using the Euler-Lagrange equation

$$2a(\ddot{q} + \omega^2 q) = \frac{\partial G}{\partial q} - \frac{d}{dt}\left(\frac{\partial G}{\partial v}\right),$$

and a second integration by parts we obtain

$$\mathcal{L}(X_\infty) = [q_\infty \dot{q}_\infty] + \int_{\mathbb{R}} G(\partial X_\infty) - \frac{1}{2} q_\infty \frac{\partial G}{\partial q}(\partial X_\infty) - \frac{1}{2}\dot{q}_\infty \frac{\partial G}{\partial v}(\partial X_\infty),$$

thus the integral can be thought as the action of the homoclinic orbit.

6. Although the setting is Lagrangian, the theory of Bolotin [6] can not be applied to our problem. Here the whole center manifold enjoys a minimizing property as used in [6], but the orbits O_r themselves do not. This is connected with the fact that the center manifold is hyperbolic with respect to the full phase space, while the periodic orbit O_r is not. For that reason, we shall rather search orbits homoclinic to the center manifold as a whole, and this is why we do not know precisely which of the periodic orbits O_r have a homoclinic.

7. It should be possible to extend Theorem 1 to more general Hamiltonian systems by using the analysis of [19] or pseudo-holomorphic curves as in [9] and [10], [11].

1.1 Normalization of the center manifold and persistence of the hypotheses

The hypotheses of Theorem 1 may appear to be very rigid. They imply for example that there is an invariant plane with elliptic linear motion on it. We see in this section a general method for normalizing center manifolds i.e. bringing them to a linear elliptic plane. This requires a change of coordinates and a reparametrisation. These operations preserve homoclinic orbits. This method can be applied to prove that the homoclinic obtained by Theorem 1 is not destroyed by a C^3-small perturbation of L.

Theorem 2 *Let* L

We define the normalized Hamiltonian $\tilde{H}_\epsilon = f_\epsilon \circ H_\epsilon \circ \Phi_\epsilon$, the associated Lagrangian $\tilde{L}_\epsilon$ can be written as (1) with a function $\tilde{G}_\epsilon$ satisfying [HG1], we have globally normalized the center manifold. Let us now compute

$$\begin{aligned} d^2\tilde{H}_\epsilon(x)\cdot(u,u) &= f_\epsilon'\big(H_\epsilon\circ\Phi_\epsilon(x)\big)\, d^2H_\epsilon(\Phi_\epsilon(x))\big(d\Phi_\epsilon(x)\cdot u\,,\, d\Phi_\epsilon(x)\cdot u\big) \\ &+ f_\epsilon'\big(H_\epsilon\circ\Phi_\epsilon(x)\big)\, dH_\epsilon(\Phi_\epsilon(x))\circ d^2\Phi_\epsilon(x)\cdot(u,u) \\ &+ f_\epsilon''\big(H_\epsilon\circ\Phi_\epsilon(x)\big)\big(dH_\epsilon(\Phi_\epsilon(x))\circ d\Phi_\epsilon(x)\cdot u\big)^2. \end{aligned}$$

We obtain that

$$\left\|\tilde{H}_\epsilon - H\right\|_{C^2} \xrightarrow[\epsilon\to 0]{} 0,$$

which implies that

$$\left\|\tilde{L}_\epsilon - L\right\|_{C^2} \xrightarrow[\epsilon\to 0]{} 0, \tag{6}$$

and we also easily see that

$$\tilde{L}_\epsilon - L = 0 \quad \text{outside } K. \tag{7}$$

We use this normalized form to prove Theorem 2.

Proof of Theorem 2: The first step is to replace the Lagrangian L_ϵ by a new Lagrangian, still noted L_ϵ, which satisfies

$$\|L_\epsilon - L\|_{C^3} \leqslant C_K\epsilon$$

and such that $L_\epsilon = L$ outside K. This can be done with a constant C_K depending only on K. When ϵ is small enough, the associated vector field has a global center manifold. We now consider the normalized Lagrangian

$$\tilde{L}_\epsilon = a\left(v^2 - \omega^2 q^2\right) + \tilde{G}_\epsilon(z,q,v) \quad (z,q,v)\in TM\times\mathbb{R}\times\mathbb{R},$$

as defined above. Let us stress that L_ϵ has an orbit homoclinic to a periodic trajectory $\mathcal{C}(\epsilon)\cap\{E=e\}$ for some $e\in[E_\epsilon(p(\epsilon)),E_0]$ if $\tilde{L}_\epsilon$ has an orbit homoclinic to a periodic trajectory O_r for some r satisfying $a\omega r^2 \leqslant E_0$. We apply Theorem 1 to find such a homoclinic orbit. There remains to check that the hypotheses of Theorem 1 are satisfied by $\tilde{L}_\epsilon$. The Lagrangian $\tilde{L}_\epsilon$ has been constructed to obtain [HG1]. The hypothesis [HL] is a direct consequence of (6) and (7) when ϵ is small enough. It is not harder to see that [HG2] holds with the constant $b/2$ instead of b for sufficiently small ϵ. We also obtain from (6) and (7) the existence of a function $c(\epsilon)>0$ with $\lim_{\epsilon\to 0} c(\epsilon) = 0$ such that

$$\left|L_\epsilon - L\right| \leqslant c(\epsilon)d^2(z,z_0)$$

and

$$\left|\left(q\frac{\partial L}{\partial q}(z,q,v) + v\frac{\partial L}{\partial v}(z,q,v)\right) - \left(q\frac{\partial \tilde{L}_\epsilon}{\partial q}(z,q,v) + v\frac{\partial \tilde{L}_\epsilon}{\partial v}(z,q,v)\right)\right| \leqslant c(\epsilon)d^2(z,z_0).$$

The hypothesis [HG3] is thus satisfied with the constant $b/2$ when ϵ is small enough. We moreover have the inequality

$$U_\epsilon = U - c(\epsilon)d^2(z,z_0) \leqslant \tilde{G}_\epsilon \leqslant W + c(\epsilon)d^2(z,z_0) \leqslant W_\epsilon.$$

The estimate (3) combined with the fact that [HG2] is satisfied with the constant $b/2$ implies

$$\big|I(W_\epsilon) - I(U_\epsilon)\big| \xrightarrow[\epsilon\to 0]{} \big|I(W) - I(U)\big|.$$

It is possible to apply Theorem 1 to $\tilde{L}_\epsilon$ when ϵ is small enough, and get a homoclinic orbit to O_r with

$$a\omega r^2 \leqslant \frac{\omega}{2\pi}\big(I(W_\epsilon) - I(U_\epsilon)\big) \leqslant E_0.$$

□

1.2 Perturbation from product systems

Let us first consider the case where G does not depend on (q, v). We can set $U = G = W$ in the notations of Theorem 1, and

$$Q(q, v) = a(v^2 - \omega^2 q^2), \quad (q, v) \in \mathbb{R}^2.$$

The system L is the uncoupled product between the linear oscillating Lagrangian system Q on $\mathbb{R}$ and the Lagrangian system U on M. It is well known that if [HU1,2] hold the Lagrangian system U has an orbit $h(t)$ homoclinic to θ_0, see Section 5. This can be recovered from Theorem 1. The hypothesis [HG3] always holds in this case, and Theorem 1 gives the existence of an orbit homoclinic to $(\theta_0, 0)$ for L, which of course implies the existence of the homoclinic of U. All the orbits O_r have a homoclinic for L in this case, given by

$$h_r(t) = (h(t), r\cos(\omega t)).$$

Let us now come back to the general case of a function $G(z, q, v)$. The theorem 1 can be seen as a perturbation result when a coupling is introduced in a product as above. Elementary dimension considerations show that the saddle-center fixed point do not have any homoclinic orbit in a generic coupled system. The theorem 1 yet gives the existence of an orbit homoclinic to some periodic orbit O_r if [HG3] holds. In view of the estimate (3) the quantity $I(W) - I(U)$ is a measure of the coupling, and we obtain that the radius r tends to zero when the coupling tends to 0. The orbit obtained by Theorem 1 can be considered as the continuation of the orbit homoclinic to the fixed point that existed in the uncoupled system. Moreover, the hypothesis [HG3] is satisfied when the coupling is small since it can be written

$$\frac{1}{2}\left(q\frac{\partial C}{\partial q} + v\frac{\partial C}{\partial v}\right) - C \leqslant U - cd^2(z, z_0)$$

if we separate the main part U and the coupling perturbation C of R: $G(z, q, v) = U(z) + C(z, q, v)$, and it is satisfied for example when

$$|C| + \left|q\frac{\partial C}{\partial q}\right| + \left|v\frac{\partial C}{\partial v}\right| \leqslant \epsilon d^2(z, z_0)$$

with a sufficiently small ϵ. Shortly, The homoclinic orbit to the fixed point that existed in the uncoupled system is turned to an orbit homoclinic to O_r when the coupling is introduced, with r as small as the coupling is small, and this homoclinic exists as long as [HG3] holds. Combining this with the methods of the preceding subsection, we obtain

Application 1 *Let us consider a smooth one parameter family L_ϵ of Lagrangian systems such that*

$$L_0(z,q,v) = a\left(v^2 - \omega^2 q^2\right) + U(z), \quad (z,q,v) \in TM \times \mathbb{R} \times \mathbb{R}$$

satisfies [HL] and [HU1,2]. There is an $\epsilon_0 > 0$ and a function $e(\epsilon) \geqslant 0$ satisfying $\lim_{\epsilon \to 0} e(\epsilon) = 0$ such that for $\epsilon \leqslant \epsilon_0$ the system L_ϵ has a saddle-center fixed point $p(\epsilon)$ and a center manifold $\mathcal{C}(\epsilon)$ intersecting transversally the energy level $E_\epsilon^{-1}\big(e(\epsilon)\big)$ along a hyper

Application 2 *The point* $(z_0, 0, 0)$ *is a saddle-center fixed point of* L^ω. *It has a center manifold* $z = z_0$, *which is filled with the periodic or*

6. The classical pendulum is described by the Lagrangian

$$G_0(\theta,\dot\theta) = \|\dot\theta\|^2 + (1-\cos\theta), \quad \theta \in S^1.$$

It is well known that integrability can be destroyed and chaotic behavior turned on by a time-dependent small perturbation. Let us consider a system

$$L^\omega(\theta,\dot\theta,q,\dot q) = \dot q^2 - \omega^2 q^2 + G(\theta,\dot\theta,q), \quad (\theta,q) \in S^1\times\mathbb{R}$$

with $G(\theta,\dot\theta,0) = G_0(\theta,\dot\theta)$, satisfying the hypotheses of the application. The homoclinic orbit obtained by the application can be made transversal by a small perturbation of G. This is a new way, also physically relevant, to introduce chaotic behavior in the classical pendulum.

Proof : We are interested in trajectories located around $q=0$, and it is first necessary to change the Lagrangian function outside a neighborhood of $q=0$. We need a smooth function $\varphi : [0,\infty] \longrightarrow [0,1]$ such that $\varphi|_{[0,1]} = 1$ and $\varphi|_{[2,\infty)} = 0$ and $0 \geqslant \varphi' \geqslant -2$. Let us fix $\delta > 0$ and define

$$G_\delta(\theta,q) = \varphi(q/\delta)G(\theta,q) + \big(1-\varphi(q/\delta)\big)G_0(\theta).$$

It is clear that G_δ satisfies HG1 when δ is small enough. To check the other hypotheses of Theorem 1 let us first notice that there is a constant $D>0$ such that

$$|G-G_0| \leqslant D\delta d^2(z,z_0) \quad \text{and} \quad \left|\frac{\partial G}{\partial q}\right| \leqslant D d^2(z,z_0)$$

when $|q| \leqslant 2\delta$. It follows from the first estimate above that $G_\delta(\theta,q) \geqslant b\, d^2(z,z_0)/2$ when δ is small enough. In view of the calculation

$$\begin{aligned} G_\delta - \frac{1}{2}q\frac{\partial G_\delta}{\partial q} &= \varphi(q/\delta)G + (1-\varphi(q/\delta))G_0 + \frac{1}{2}q\left(\varphi(q/\delta)\frac{\partial G}{\partial q} + (G-G_0)\varphi'(q/\delta)/\delta\right) \\ &\geqslant G_0 - \big(\varphi(q/\delta) - \varphi'(q/\delta)\big)|G-G_0| - \delta\varphi(q/\delta)\left|\frac{\partial G}{\partial q}\right| \\ &\geqslant G_0 - 4\delta D\, d^2(z,z_0), \end{aligned}$$

the hypotheses HG2 and HG3 are both satisfied with the constant $b/3$ when δ is small enough. We also obtain that

$$U_\delta = G_0(z) - D\delta d^2(z,z_0) \leqslant G_\delta \leqslant G_0(z) + D\delta d^2(z,z_0) = W_\delta,$$

and (3) yields

$$I(W_\delta) - I(U_\delta) \leqslant 3D\delta/b.$$

We are now in a position to apply Theorem 1 to $L^\omega_\delta = a\left(v^2 - \omega^2 q^2\right) + G_\delta(z,q,v)$, and obtain an orbit $h^\omega_\delta = (z^\omega_\delta,\, q^\omega_\delta,\, \dot q^\omega_\delta)$ homoclinic to O_r with $r \leqslant B\sqrt{\delta/\omega}$, where B is a constant that depends neither on ω nor on δ. The energy function E^ω_δ associated with L_δ is $E^\omega_\delta = a\left(v^2+\omega^2q^2\right) + E_0(z)$, where E_0, the energy associated to G_0, is bounded from below. Writing energy conservation along h^ω_δ yields

$$\omega^2\,|q^\omega_\delta|^2 \leqslant B^2\delta\omega + C.$$

The homoclinic h^ω_δ is thus an orbit of L^ω if

$$\omega^2\delta^2 \geqslant B^2\delta\omega + C.$$

Let us now choose $\delta = \beta/\omega$, where β is a sufficiently large fixed number, the inequality above is satisfied and the homoclinic $h^\omega = h^\omega_{\beta/\omega} = (z^\omega, q^\omega, \dot{q}^\omega)$ is a trajectory of L^ω, of bounded energy $E^\omega(h^\omega) \leqslant E^\infty = aB^2\beta$, and satisfying

$$\int_{\mathbb{R}} G_{\beta/\omega}(h_\omega) - \frac{1}{2}q^\omega \frac{\partial G_{\beta/\omega}}{\partial q}(h^\omega) \leqslant I(W_{\beta/\omega}).$$

In view of HG2, this estimate yields

$$\int d^2(z^\omega, z_0) \leqslant C.$$

The function $G_q(z) = G(z,q)$ is a fiberwise convex Lagrangian on TM for all q. Let us call Y_q the associated vectorfield. The curves z^ω satisfy the Euler-Lagrange equation

$$\dot{z}^\omega(t) = Y_{q^\omega(t)}(z^\omega(t))$$

hence z^ω is bounded in $C^1(\mathbb{R}, TM)$. Moreover, since $\|q^\omega\| \longrightarrow 0$, any limit curve z^∞ of z^ω satisfies the Euler-Lagrange equation

$$\dot{z}^\infty(t) = Y_0(z^\infty(t)),$$

where Y_0 is the Euler-Lagrange vector field of G_0. It is not hard to see that there is a constant $C > 0$ independent of ω such that all orbit Z of G_0 homoclinic to z_0, satisfies $\int d^2(Z, z_0) \geqslant C$, and all orbit $X = (Z, Q, \dot{Q})$ of L^ω homoclinic to the center manifold $z = z_0$ and lying in $E^\omega \leqslant E^\infty$ satisfies $\|d(Z, z_0)\|_\infty \geqslant C$. One now applies the concentration compactness principle, see [27], 4.3, to the function $d^2(z^\omega(t), z_0)$ in order to prove the last point of the theorem, see [13] for the use of concentration compactness with homoclinic orbits. In our situation, vanishing is impossible since $\|d^2(z^\omega, z_0)\|_\infty \geqslant C^2$, while only a finite number of bumps can appear since each bump satisfies $\int d^2(Z^i, z_0) \geqslant C$. To finish, the persistence is a direct consequence of Theorem 2. □

2 Lagrangian and Hamiltonian systems

We recall in this section some standard facts abound Lagrangian and Hamiltonian systems. This is an opportunity to introduce some notations and to state a simple estimate of the energy function that will be used throughout the paper.

Let N be a manifold, $TN \xrightarrow{\pi} N$ the tangent bundle and $T^*N \xrightarrow{\pi^*} N$ the cotangent bundle. The lifting ∂x of a curve $x : \mathbb{R} \longrightarrow N$ is the curve

$$\begin{aligned} \partial x : \mathbb{R} &\longrightarrow TN \\ t &\longmapsto dx(t, 1). \end{aligned}$$

We consider a smooth Lagrangian function $L : TN \longrightarrow \mathbb{R}$, that is uniformly convex on each fiber. The fiber derivative L_v of L is well defined, and the application

$$\begin{aligned} \phi : TN &\longrightarrow T^*N \\ v &\longmapsto L_v(v) \end{aligned}$$

is a diffeomorphism. We can associate to L an energy function

$$\begin{aligned} E : TN &\longrightarrow \mathbb{R} \\ v &\longmapsto \langle \phi(v), v \rangle - L(v) \end{aligned}$$

and a Hamiltonian function

$$\begin{aligned} H : T^*N &\longrightarrow \mathbb{R} \\ p &\longmapsto E \circ \phi^{-1}(p) = \langle v, \phi^{-1}(v) \rangle - L(\phi^{-1}(v)). \end{aligned}$$

There is a canonical symplectic structure Ω on T^*M, and we associate to H its Hamiltonian vector field X defined by the equation $i_X\Omega = -dH$. Let us define the action $\mathcal{L}(x)$ of a smooth curve $x : [t_0, t_1] \longrightarrow N$

$$\mathcal{L}(x) = \int_{t_0}^{t_1} L(\partial x(t))\, dt,$$

we say that x is a trajectory of L if it is a critical point of the action with respect to fixed endpoints variations. A curve $x : \mathbb{R} \longrightarrow N$ is a trajectory of L if and only if its restrictions to finite time intervals are trajectories of L. We will pay special attention to the periodic trajectories of L. Let $T > 0$ be a fixed period, a T-periodic curve $x : \mathbb{R} \longrightarrow N$ is a trajectory of L if and only if the loop $x_{|[0,T]}$ is a critical point of

$$\mathcal{L}_T(x) = \int_0^T L(\partial x)$$

on $C^1_T = \{x \in C^1([0,T], M) \,/\, x(0) = x(T)\}$. There is a one to one correspondence between trajectories x of L and integral curves z of X, given by

$$x \longrightarrow z = \phi(\partial x)\ , \quad z \longrightarrow x = \pi^*(z).$$

As a consequence, there is a vector-field Y on TM such that x is a trajectory of L if and only if ∂x is an integral curve of Y, and we have

$$Y(z) = (d\phi_z)^{-1}(X(\phi(z)).$$

In any canonical chart (q, v) of TM, the trajectories of L satisfy the Euler-Lagrange equations

$$\frac{d}{dt}\frac{\partial L}{\partial v}(q(t), \dot{q}(t)) = \frac{\partial L}{\partial q}(q(t), \dot{q}(t)).$$

The Hamiltonian function H is invariant along integral curves of X, and the energy E is invariant along integral curves of Y hence $E(\partial x)$ is constant if x is a trajectory of L. This construction can be reversed. Let $H : T^*N \longrightarrow \mathbb{R}$ be a Hamiltonian function. If the mapping

$$\begin{aligned} \psi : TN &\longrightarrow TN \\ z &\longmapsto H_v(z) \end{aligned}$$

is a diffeomorphism, which happens when H is fiberwise convex and proper, we define

$$L(z) = (z, \psi^{-1}(z)) - H(\psi^{-1}(z)),$$

the associated mapping ϕ is the diffeomorphism $\phi = \psi^{-1}$, and the correspondence described above between orbits of L and integral curves of H holds. Let us now come back to our main subject of interest and estimate the energy E associated to (1). We assume [HG1-4] and [HL]. Note that the configuration space of L is $N = M \times \mathbb{R}$.

Lemma 1 *Let* L

defined on T-periodic loops. The critical points of $\mathcal{L}_l$ are precisely the T-periodic solutions of L_l. We will see in Section 8, that it is possible to find a critical value $c_T(l)$ of $\mathcal{L}_l$ for all $T = 2\pi\tau/\omega$, $\tau \in \mathbb{N}$, and all $l \in (1, 1 + 1/\tau)$. This critical value satisfies

$$I_T(U) \leqslant c_T(l) \leqslant I_T(W),$$

the numbers $I_T(U)$ are defined in Section 5 together with the numbers $I(U)$. Since the function $l \longrightarrow c_T(l)$ is non-increasing, there is an $l(T) \in (1, 1+1/\tau)$ such that $c_T'(l(T))$ exists and $|c_T'(l(T))| \leqslant \tau(I_T(W) - I_T(U))$. We use this to find a critical point $X_T = (\theta_T, q_T)$ of $\mathcal{L}_{l(T)}$ at level $c_T(l(T))$ such that

$$2l(T)a\omega^2\|q_T\|_2^2 = \left|\frac{\partial \mathcal{L}_l(X_T)}{\partial l}\right|_{l(T)} \leqslant 1 + |c_T'(l(T))| \leqslant \tau(I_T(W) - I_T(U)) + 1.$$

The periodic orbit we obtain is not trivial i.e. $\theta_T \not\equiv \theta_0$, because it has nonzero action. All this is detailed in Section 8, where we prove

Proposition 2 *For all $\tau \in \mathbb{N}$ and $T = 2\pi\tau/\omega$, there exist a parameter $l(T)$ in the interval $(0, 1/\tau)$ and a trajectory $X_T = (\theta_T, q_T)$ of $L_{l(T)}$ such that*

$$\begin{aligned} \frac{1}{T}\|q_T\|_2^2 &\leqslant \frac{1}{4\pi a\omega}\left(I_T(W) - I_T(U) + \frac{1}{\tau}\right), \\ \mathcal{L}_{l(T)}(X_T) &= c_T(l(T)), \\ \theta_T &\not\equiv \theta_0. \end{aligned}$$

We now use the periodic orbits given by Proposition 2 to build the homoclinic orbit. Since $I(U) = \liminf_{T\to\infty} I_T(U)$, see Section 5, we can extract a subsequence T_n of T such that

$$\frac{1}{T_n}\|q_{T_n}\|_2^2 \longrightarrow r^2/2$$

with a radius

$$r \leqslant \sqrt{\frac{I(W) - I(U)}{2\pi a\omega}}.$$

We obtain a sequence X_n of T_n-periodic orbits of $L_{l(T_n)}$ which satisfies

- $l(T_n) \longrightarrow 0$
- $T_n \longrightarrow \infty$,
- $\mathcal{L}_{l(T_n)}(X_n) \leqslant I(W)$,
- $\|q_n\|^2/T_n \longrightarrow r^2/2$,
- $\theta_n \not\equiv \theta_0$,

and we prove in Section 7 that a homoclinic orbit can be found as an accumulation point of this sequence. Moreover, this homoclinic orbit satisfies all the conclusions of Theorem 1. $\square$

4 Control at infinity

All the systems considered in this paper are autonomous, and preserve an energy function. It is thus possible to change the Lagrangian function at infinity without changing the flow on prescribed compact energy shells. This observation is the basis of the proof of Proposition 1. The details are not simple since fiberwise convexity has to be preserved during the process. We now prove Proposition 1.

Let us consider a Lagrangian function $L = a(v^2 - \omega^2 q^2) + G$ satisfying all the hypotheses of Theorem 1. We build a new Lagrangian function that is equal to the old one on $E \leqslant E_0$ and that satisfies all the hypothesis of Theorem 1 and [HG4]. Recall that

$$E_0 = \frac{\omega}{2\pi}\big(I(W) - I(U)\big).$$

First step : Let K_0 be a compact subset of TM, we first build a function G_1 on $TM \times \mathbb{R}^2$ such that

$$G_1(z,q,v) = G(z,q,v) \quad \text{when } z \in K_0,$$
$$G_1(z,q,v) = \alpha\|z\|^2 \quad \text{when } z \notin K$$

for some $\alpha > 0$ and some compact set $K \subset TM$, and $G_1 + av^2$ is fiberwise convex. Let us set $d = \sup_{z\in K_0} \|z\|$. There exists a $d_1 > d$ such that for all $\alpha > 0$ there exists a convex function $f_\alpha : [0,\infty) \longrightarrow [0,\infty)$ satisfying $f_\alpha(x) = 0$ when $x \leqslant d$ and $f_\alpha(x) = \alpha x^2$ when $x \geqslant d_1$. We now consider a function $\varphi : \mathbb{R}^+ \longrightarrow [0,1]$ such that $\varphi(x) = 1$ when $x \leqslant d_1$ and $\varphi(x) = 0$ when $x \geqslant d_2$ for some $d_2 > d_1$ and we set

$$G_1(z,q,v) = \varphi(\|z\|)G(z,q,v) + f_\alpha(\|z\|^2).$$

It is easy to see that the function $G_1 + av^2$ is fiberwise convex except maybe on the compact set $d_1 \leqslant \|z\| \leqslant d_2$. To prove that $G_1 + av^2$ is also fiberwise convex on $d_1 \leqslant \|z\| \leqslant d_2$ we will use a local canonical chart $z = (\theta, \nu)$ of TM and prove that the function $G_1 + av^2$ is convex in (ν, v) on $d_1 \leqslant \|z\| \leqslant d_2$. We first note that

$$d_v^2(G_1 + av^2) = \varphi(\|z\|)d_v^2 G + 2a$$

is positive since $d_v^2 G + 2a$ is positive. On the other hand, it is not hard to see that given $\beta > 0$ one can take α large enough so that $d_\nu^2 G_1 \geqslant \beta Id$ on $d_1 \leqslant \|z\| \leqslant d_2$. Since the cross derivatives $d_\nu d_v(G_1 + av^2)$ do not depend on α, one can check that the Hessian

$$\begin{bmatrix} d_\nu^2(G_1 + av^2) & d_v d_\nu(G_1 + av^2) \\ d_\nu d_v(G_1 + av^2) & d_v^2(G_1 + av^2) \end{bmatrix}$$

is positive definite when α is large enough. The function $G_1 + av^2$ is thus fiberwise convex, as well as the function $L_1 = a(v^2 - \omega^2 q^2) + G_1$. The function G_1 satisfies [HG1-3] with the same constant b. Let us define the functions

$$U_1(z) = \varphi(\|z\|)U(z) + f_\alpha(\|z\|^2), \tag{9}$$

and W_1 in the same way, so that

$$U_1 \leqslant G_1 \leqslant W_1.$$

If α has been chosen large enough, the functions U_1 and W_1 are fiberwise convex and satisfy

$$U_1(z) = U(z) \text{ and } W_1(z) = W(z) \quad \text{when } z \in K_0$$

and

$$U_1(z) = W_1(z) = \alpha\|z\|^2 \quad \text{when } z \notin K.$$

If K_0 is sufficiently large, the definition of I is $I(U) = I(U_1)$ and $I(W) = I(W_1)$ (see Definition 2 of Section 5). So the maximal energy E_0 has not been changed.

SECOND STEP : We now want to control the behavior for large v. Let B_0 be a compact subset of $TM \times \mathbb{R}^2$, we will define a function $\tilde{G}$ such that $\tilde{G} = G_1$ on B_0 and $\tilde{G} = U_1$ outside a compact subset $B \supset B_0$. To do this we first observe that $U_1 - G_1$ is bounded, which easily implies that $d_v G_1$ is bounded since $d_v^2 G_1 \geqslant -2a$. We now take a compactly supported function $\psi : \mathbb{R}^2 \longrightarrow [0,1]$ such that $\psi(q,v) = 1$ when there exists a $z \in TM$ with $(z,q,v) \in B_0$, and such that $\|d\psi\|_{C^1} \leqslant \epsilon$ and $qd_q\psi + vd_v\psi \leqslant 0$. The function

$$\tilde{G}(z,q,v) = \psi(q,v)G_1(z,q,v) + (1-\psi(q,v))U_1(z),$$

satisfies [HG4] with $G_\infty = U_1$. With the notation $z = (\theta, \nu)$, we derive

$$\begin{aligned} d_\nu^2\tilde{G} &= \psi d_\nu^2 G_1 + (1-\psi)d_\nu^2 U_1 \\ d_v d_\nu \tilde{G} &= \psi d_v d_\nu G_1 + d_v\psi(d_\nu G_1 - d_\nu U_1) \\ d_v^2\tilde{G} &= \psi d_v^2 G_1 + d_v^2\psi\,(G_1 - U_1) + 2d_v\psi\, d_v G_1, \end{aligned}$$

and get

$$\Big\| d^2_{(v,\nu)}(\tilde{G} + av^2) - \psi d^2_{(v,\nu)}(G_1 + av^2) - (1-\psi)d^2_{(v,\nu)}(U_1 + av^2)\Big\|_\infty \xrightarrow[\epsilon\to 0]{} 0,$$

which implies that $\tilde{G}$ is fiberwise convex when ϵ is small enough. The function $\tilde{G}$ moreover satisfies [HG1-2] with the same constant b and [HG3] follows from :

$$\begin{aligned} \tilde{G} - \frac{1}{2}\big(qd_q\tilde{G} + vd_v\tilde{G}\big) =& \psi\Big(G_1 - \frac{1}{2}\big(qd_qG_1 + vd_vG_1\big)\Big) \\ &+ (1-\psi)U_1 + \frac{1}{2}(U_1 - G_1)\big(qd_q\psi + vd_v\psi\big) \\ &\geqslant \psi b\, d^2(z,z_0) + (1-\psi)b\, d^2(z,z_0) = b\, d^2(z,z_0). \end{aligned}$$

The function $\tilde{L} = a(v^2 - \omega^2 q^2) + \tilde{G}$ satisfies all the hypotheses of Theorem 1 with the same constants a, b and ω, and with U and W replaced by U_1 and W_1. Let us assume that K_0 and B_0 have been taken large enough so that

$$\{E \leqslant E_0\} \subset B_0 \cap \big(K_0 \times \mathbb{R}^2\big).$$

If the conclusions of Theorem 1 hold for $\tilde{L}$, they give the existence of a homoclinic of energy below E_0, this orbit is also an orbit of L since the function have not been changed in this region, the conclusion of Theorem 1 thus hold for L. □

5 Systems with a hyperbolic fixed point.

We now define the number $I(U)$ for a fiberwise convex Lagrangian $U : TM \longrightarrow \mathbb{R}$ satisfying [HU1-2]. These hypotheses imply that z_0 is a hyperbolic fixed point of the system, see Section 6, and that it has a homoclinic orbit. Homoclinics for this kind of Lagrangian were first studied variationally by Bolotin [5], and then by several authors (see e.g. [2] ,[25]). These works have also been extended to more general Hamiltonian systems in [16] and [10], [11]. We just give here a presentation of these results that will be useful in the proof of Theorem 1. We shall first study Lagrangians U with the additional hypothesis
HU3 : There exists an $\alpha > 0$ such that $U(z) = \alpha\|z\|^2$ outside a compact set of TM.
Although we are interested mainly in the homoclinic orbit, we shall use a variational setting for T-periodic orbits of U: Let Λ_T be the manifold of H^1-loops

$$\gamma : S_T = \mathbb{R}/T\mathbb{Z} \to M,$$

the action functional

$$\begin{aligned} \mathcal{U}_T : \Lambda_T &\longrightarrow \mathbb{R} \\ \gamma &\longmapsto \int_0^T U(\partial\gamma(t))\, dt \end{aligned}$$

is smooth and satisfies the Palais-Smale condition. We note H the rational cohomology. A pointed set (S, s) is a set S with a distinguished element $s \in S$. We will use the notation $H(S, s)$ for the relative cohomology $H(S, \{s\})$. For any closed subset S of Λ_T containing the constant loop $\theta \equiv \theta_0$ we consider the morphism

$$i_S^* : H(\Lambda_T, \theta_0) \longrightarrow H(S, \theta_0)$$

associated with the inclusion.

Definition 1 *We call Σ_T the family of all compact su*

PROOF OF LEMMA 3: This is very classical, and we shall only outline the proof. If M is not simply connected, there is a non contractible curve $\gamma \in \Lambda_1^0$. We take $K = \{\gamma, \theta_0\}$, and see that $i_K^*(H^0(\Lambda_1, \theta_0)) \neq 0$. Things are much harder when M is simply connected. Let us set $C = C^0(S^1, M)$ and $C^0 = \{\gamma \in C^0(S^1, M) \ / \ \gamma(0) = \theta_0)$, the inclusion $i_{\Lambda_0} : (\Lambda^0, \theta_0) \longrightarrow (\Lambda, \theta_0)$ is homotopy equivalent to the inclusion $ic : (C^0, \theta_0) \longrightarrow (C, \theta_0)$. A theorem of Sullivan gives the existence of infinitely many nonzero rational Betti numbers of the space $C^0(S^1, M)$ if $\pi_1(M) = 0$, see [28], page 46. Then, we consider the Serre fibration

$$\begin{aligned} C &\longrightarrow M \\ \gamma &\longmapsto \gamma(0) \end{aligned}$$

of fiber C^0 to prove that ic^* is nonzero, hence $i_{\Lambda_0}^*$ is nonzero. We now use broken geodesics approximation, see [7], to find a compact K representing this cohomology. □

PROOF OF LEMMA 2: For any $T \geqslant 1$, we can extend loops in Λ_1^0 to $[0, T]$ by fixing them in θ_0 outside $[0, 1]$, this defines the injection

$$\begin{aligned} j_T : (\Lambda_1^0, \theta_0) &\longrightarrow (\Lambda_T^0, \theta_0) \\ \theta(t) &\longmapsto j_T(\theta(t)) = \theta(\min(1, t)) \end{aligned}$$

which is homotopic to the diffeomorphism

$$\begin{aligned} (\Lambda_1^0, \theta_0) &\longrightarrow (\Lambda_T^0, \theta_0) \\ \theta(t) &\longmapsto S_T(\theta(t)) = \theta(t/T). \end{aligned}$$

It follows that $j_T(K) \in \Sigma_T$, thus

$$I_T(U) \leqslant \sup_{j_T(K)} \mathcal{U}_T = \sup_K \mathcal{U}_1$$

because the trajectory $t \longmapsto \theta_0$ has zero action. This ends the proof of the lemma since $\sup_K \mathcal{U}_1$ is a finite number. □

There is a T-periodic trajectory γ_T such that $\mathcal{U}_T(\gamma_T) = I_T(U)$. We shall not prove it since it is very classical, and involves arguments simpler than those of Section 8. Here non-trivial means that $\gamma_T \not\equiv \theta_0$. We can define the number

$$I(U) = \liminf_{T \longrightarrow \infty} I_T(U).$$

There must be a nontrivial homoclinic orbit to z_0 such that

$$\int_{-\infty}^{\infty} U(\dot{\gamma}(t))\, dt \leqslant I(U),$$

we obtain it as an accumulation point of the sequence γ_T of periodic orbits, compare Section 7. The following proposition is useful for applications

Proposition 3 *The function $U \longmapsto I(U)$ is increasing and continuous:*

$$U \leqslant W \Longrightarrow I(U) \leqslant I(W)$$

$$\left| 1 - \frac{I(W)}{I(U)} \right| \leqslant \sup_z \frac{|W(z) - U(z)|}{b\, d^2(z, z_0)}$$

for all U and W satisfying [HU1-3]. Recall that b is the constant of [HU2].

PROOF : The monotonicity is clear, we shall prove regularity. Let us set

$$\lambda = \sup_z \frac{|W(z) - U(z)|}{b\, d^2(\pi(z), \theta_0)},$$

we obtain using [HU2] that

$$(1-\lambda)U \leqslant W \leqslant (1+\lambda)U.$$

This yields

$$(1-\lambda)I_T(U) \leqslant I_T(W) \leqslant (1+\lambda)I_T(U).$$

We thus have for any T

$$\left|1 - \frac{I_T(W)}{I_T(U)}\right| \leqslant \lambda,$$

and we obtain the proposition by taking the limit. □

Let us come back to Lagrangians systems U on TM satisfying only [HU1,2] but not [HU3]. The energy function E_U is proper and the sets $E_U^e = \{E_U \leqslant e\}$ are compact. Let $\mathcal{E}_U$ be the set of all Lagrangians U_1 satisfying [HU1-3] and such that $U_1 = U$ on E_U^e for some $e > 0$. Elements of $\mathcal{E}_U$ can be constructed by the methods of Section 4.

Definition 2 *For all Lagrangian function U satisfying [HU1,2], we set*

$$I(U) = I(U_1)$$

for any $U_1 \in \mathcal{E}_U$. The proposition 3 holds for U and W satisfying [HU1,2] with this extended definition of I.

PROOF : One has to prove that the number $I(U_1)$ does not depend on the choice of the Lagrangian $U_1 \in \mathcal{E}_U$. Let us take two Lagrangians U_1 and U_0 in $\mathcal{E}_U$, define $U_t = tU_1 + (1-t)U_0$, $t \in [0,1]$, and let E_t be the energy function associated with U_t. There is an energy $e > 0$ such that $U_t(z) = U(z)$ for all $z \in E_U^e$ and all $t \in [0,1]$. The Lagrangians U_t satisfy [HU1,2] with the same constant b, and $U_t = \alpha_t \|z\|^2$ at infinity hence $E_t = \alpha_t \|z\|^2$ at infinity. Since α_t, $t \in [0,1]$ is bounded there is a constant $C > 0$ independent of t such that $E_t \leqslant C d^2(z, z_0)$, see Lemma 1. For all $T > 0$ there is a T-periodic trajectory γ_T^t of U_t such that $\mathcal{U}_T^t(\gamma_T^t) = I_T(U_t)$. One can build by the methods of Section 4 a Lagrangian $U_2 \in \mathcal{E}_U$ such that $U_2 \geqslant \max(U_0, U_1)$ and thus $U_2 \geqslant U_t$ for all $t \in [0,1]$. It follows that $\mathcal{U}_T^t(\gamma_T^t) = I_T(U_t) \leqslant I_T(U_2)$ is bounded, and

$$E_t(\gamma_T^t) \leqslant \frac{C}{T} \int d^2(\partial\gamma_T^t, z_0) \leqslant \frac{C}{Tb} \int U_t(\partial\gamma_T^t) \leqslant \frac{C'}{T}.$$

As a consequence, there exists a $T_0 > 0$ such that all the periodic orbits $\partial\gamma_T^t$ with $T \geqslant T_0$ are contained in $\{E_t \leqslant e\}$, which is nothing but E_U^e. The curves γ_T^t with $T \geqslant T_0$ are thus all trajectories of U and of U_0, and the value $I_T(U_t)$ is critical for the Lagrange action $\mathcal{U}_T^0$ associated with U_0. The set of critical values of $\mathcal{U}_T^0$ has measure zero. This is a non-trivial application of Sard's Theorem, see for example [11], Lemma 3.1, for a result of this kind. On the other hand, we see from Proposition 3 that the function $t \longmapsto I_T(U_t)$ is continuous, hence constant since it takes values in a set of measure zero. We have proved that $I_T(U_1) = I_T(U_0)$ when T is large enough, hence $I(U_1) = I(U_0)$, and the definition is meaningful. We now prove that Proposition 3 holds with this extended definition. Let us consider two Lagrangian functions U and W satisfying [HU1-2]. We use the construction of

Section 4 to build distinguished elements of $\mathcal{E}_U$ and $\mathcal{E}_W$. We take K_0 containing E^0_U and E^0_W in its interior, and define $U_1 \in \mathcal{E}_U$ and $W_1 \in \mathcal{E}_W$ by the same expression (9). It is clear that $U_1 \leqslant W_1$ if $U \leqslant W$, and that $|I(U_1) - I(W_1)| \leqslant |I(U) - I(W)|$. Proposition 3 for U_1 and W_1 thus implies Proposition 3 for U and W. □

Let us now come back to the full system.

6 Local structure.

In this section, we focus our attention o the vicinity of the center manifold $z = z_0$. Let us define the balls $D_\delta = B(\theta_0, \delta) \in M$ and $B_\delta = B(z_0, \delta) \in TM$. We will work in a local chart of M around θ_0, that is we identify D_δ with a neighborhood of 0 in $\mathbb{R}^n$. The local form of the Lagrangian function is

$$L(\theta, \nu, q, v) = a\left(v^2 - \omega^2 q^2\right) + G(\theta, \nu, q, v), \quad (\theta, \nu, q, v) \in D_\delta \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R},$$

and we can compute the associated energy function, see Section 2

$$E(\theta, \nu, q, v) = a\left(v^2 + \omega^2 q^2\right) + \left(\nu, \frac{\partial G}{\partial \nu}\right) + \left(v, \frac{\partial G}{\partial v}\right) - G. \tag{10}$$

The hypothesis [HG2] implies

[HG2 loc] :

$$\frac{\partial^2 G}{\partial(\theta, \nu)^2}(0, 0, q, v) \geqslant b.$$

We will only use this local minimizing property in this section. The following lemma will not be used in the sequel, but Lemma 5 below is the key to nontriviality.

Lemma 4 (Description of the local orbit structure) *If the hypotheses [HR1-2] are satisfied, the flow has a saddle-center fixed point* $(0, 0) \in B_\delta \times \mathbb{R}^2$ *with a 2-dimensional elliptic space and a 2n-dimensional hyper*

we apply ϕ^{-1} to obtain the expression of the associated orbits of Y. We now prove hyperbolicity. The hypersurface

$$\Sigma = \{(z, q, v) \in TM \times \mathbb{R}^2 \,/\, q > 0 \text{ and } v = 0\} = TM \times \mathbb{R}^+_*$$

is transversal to the flow around $\{0\} \times \mathbb{R}^+_*$, and we define the associated Poincaré return map Φ. Let us fix a $r > 0$, we want to study the eigenvalues of modulus 1 of the linearized map $d\Phi(0, r)$. Note that $\Phi_{|\{0\}\times\mathbb{R}^+_*} = Id$, thus $d\Phi(0, r)_{|\{0\}\times\mathbb{R}} = Id$. It follows that for all $\epsilon > 0$ there is a fully resonant approximation Ψ of $d\Phi(0, r)$ such that

$$\|\Psi(q, z) - d\Phi(0, r)(q, z)\| \leqslant \epsilon \|z\|.$$

By fully resonant, we mean that all the eigenvalues of modulus 1 of Ψ are roots of the unity. We can moreover take ϵ small enough so that Ψ and $d\Phi(0, r)$ have the same number of eigenvalues of modulus 1. Since $\Psi_{|\{0\}\times\mathbb{R}^+_*} = Id$ there exists a neighborhood of $(0, r) \in \Sigma$ where

$$|\Phi(z, q) - d\Phi(0, r)(z, q - r) - (0, r)| \leqslant \epsilon \|z\|^2.$$

As a consequence, there exists a function G_1 satisfying [HG1] and [HG2] with a smaller constant b_1 and such that Poincaré map Φ_1 of the flow associated to

$$L_1(\theta, \nu, q, v) = a\left(v^2 - \omega^2 q^2\right) + G_1(\theta, \nu, q, v)$$

satisfies $\Phi_1(z, q) = (0, r) + \Psi(z, q - r)$ in a neighborhood of $(0, r)$. Let us consider an eigenspace of Ψ associated with a pair of eigenvalues of modulus one, which are therefore root of the unity. This eigenspace is filled with periodic points, moreover given $\delta > 0$ there exists a neighborhood of 0 in the eigenspace such that all the points in this neighborhood have their Ψ-orbit contained in the zone where $\Phi_1(z, q) = (0, r) + \Psi(z, q - r)$, and such that the periodic orbits of L_1 associated with these Φ_1-orbits are contained in $B_\delta \times \mathbb{R}^2$. We now apply the lemma 5 below to L_1 and obtain that the periodic orbits we just constructed must be the trivial ones, corresponding to the fixed space $\{0\} \times \mathbb{R}$ of Ψ. As a consequence, the linearized Poincaré map $d\Phi(0, r)$ can have no eigenvalue of modulus 1 except the one associated with this fixed space. □

Lemma 5 *Let L*

in $B_\delta \times \mathbb{R}^2$. Let us now consider a closed trajectory $(\theta(t), q(t))$ such that $(\theta, \dot\theta) \in B_\delta$, the equation

$$\int \left\langle \theta, \frac{\partial G}{\partial \theta} \right\rangle = \int \left\langle \theta, \frac{d}{dt}\frac{\partial G}{\partial \nu} \right\rangle = -\int \left\langle \dot\theta, \frac{\partial G}{\partial \nu} \right\rangle$$

yields

$$\frac{b}{2}\int \|\theta\|^2 \leqslant \int \left\langle \theta, \frac{d}{dt}\frac{\partial G}{\partial \nu} \right\rangle \leqslant -\frac{b}{2}\int \|\dot\theta\|^2.$$

It follows that $\|\theta\| \equiv \|\dot\theta\| \equiv 0$. □

7 Convergence of sequences of periodic orbits.

In this section, we prove the convergence of good sequences of periodic orbits to homoclinic orbits. We first state the strong minimizing property of the subspace $z = z_0$.

Lemma 6 *Any T-periodic trajectory $X = (\theta, q)$ of L satisfies*

$$\mathcal{L}_T(X) \geqslant b\int_0^T d(\partial\theta, z_0)^2.$$

Proof : If $X = (\theta, q)$ is a trajectory, q must satisfy the first Euler-Lagrange equation

$$2a\left(\ddot q + \omega^2 q\right) = \frac{\partial G}{\partial q} - \frac{d}{dt}\left(\frac{\partial G}{\partial v}\right).$$

If X is closed, we can integrate by parts to write its action

$$\begin{aligned}\mathcal{L}(X) &= \int -aq\left(\ddot q + \omega^2 q\right) + G(\partial\theta, q, \dot q)\\ &= \int \left(G - \frac{1}{2}q\frac{\partial G}{\partial q}\right) + \int \frac{1}{2}q\frac{d}{dt}\left(\frac{\partial G}{\partial v}\right),\end{aligned}$$

and integrating by part again the last term,

$$\mathcal{L}(X) = \int G - \frac{1}{2}q\frac{\partial G}{\partial q} - \frac{1}{2}\dot q\frac{\partial R}{\partial v} \geqslant b\int d(\partial\theta, z_0)^2.$$

□

This lemma roughly implies that if there exists a sequence of periodic orbits of L of unbounded period and bounded action, there must be an orbit homoclinic to the center manifold. Unfortunately, there is no confinement in the q direction, and we must have some estimate of the q part of the periodic orbits in order to be able to prove convergence. As explained in the sketch of proof, we must allow the parameter ω to vary. Consider now a sequence ω_n of pulsations, with a limit ω, and the associated Lagrangian and action L_n and $\mathcal{L}_n$. We have the following convergence property:

Proposition 4 *If there exist a constant M, a radius r and a sequence $X_n = (\theta_n, q_n)$ of T_n-periodic or*

- $T_n \longrightarrow \infty$,
- $\mathcal{L}_n(X_n) \leqslant M$,
- $\|q_n\|^2/T_n \longrightarrow r^2/2$,
- $\theta_n \not\equiv \theta_0$,

then there exists an or

is finite. Since the curve $\partial\theta_\infty$ has bounded derivative this yields

$$\lim_{t\to\pm\infty} \partial\theta(t) = z_0.$$

Using once more the lemma 1 we get that

$$a(\dot{q}_\infty^2 + \omega^2 q_\infty^2) \underset{t\to\pm\infty}{\longrightarrow} E_\infty = a\omega^2 r^2.$$

This is the definition we have taken for a homoclinic orbit. The last inequality follows from

$$\int_{-T_n/2}^{T_n/2} G(\partial X_n) - \frac{1}{2} q_n \frac{\partial G}{\partial q}(\partial X_n) - \frac{1}{2}\dot{q}_n \frac{\partial G}{\partial v}(\partial X_n) = \mathcal{L}(X_n) \leqslant M$$

since the integrand is non-negative. □

8 Existence of periodic orbits.

Let us fix a period $T = 2\pi\tau/\omega$, $\tau \in \mathbb{N}$. For any $l \in \mathbb{R}$, the functional

$$\begin{aligned} \mathcal{Q}_l : C_T^\infty(\mathbb{R}) &\longrightarrow \mathbb{R} \\ x(t) &\longmapsto a\int_0^T \dot{x}(t)^2 - l^2\omega^2 x(t)^2 \end{aligned}$$

can be computed using Fourier expansion:

$$\mathcal{Q}_l\left(\sum_k q_k e^{ik\omega t/\tau}\right) = aT\sum_k \left(\frac{k^2\omega^2}{\tau^2} - l^2\omega^2\right)|q_k|^2.$$

It follows that $\mathcal{Q}_l$ can be extended to

$$E_T = H^1(S_T = \mathbb{R}/T\mathbb{Z}, \mathbb{R})$$

as a continuous quadratic form. It has a two dimensional kernel when $l \in \mathbb{Z}/\tau$, and is non-degenerate for other values of l. Let us set

$$\begin{aligned} E^+ &= \{q \text{ such that } q_k = 0 \text{ when } |k| \leqslant \tau\} \\ E^- &= \{q \text{ such that } q_k = 0 \text{ when } |k| > \tau\}, \end{aligned}$$

there is an orthogonal splitting

$$E_T = E^+ \oplus E^-,$$

such that $\pm\mathcal{Q}_l|_{E^\pm}$ is positive definite for all $l \in (1, 1+1/\tau)$. Notice that E^- is finite dimensional, which is the usual feature of Lagrangian formulations. Let us define the functionals

$$\begin{aligned} \mathcal{G} : A_T = \Lambda_T \times E_T &\longrightarrow \mathbb{R} \\ x(t) = (\theta(t), q(t)) &\longmapsto \int_0^T G(\partial\theta(t), q(t), \dot{q}(t))\, dt. \end{aligned}$$

and

$$\begin{aligned}\mathcal{L}_l : A_T = &\Lambda_T \times E_T \quad \longrightarrow \mathbb{R} \\ x(t) = &(\theta(t), q(t)) \longmapsto \mathcal{L}_l(x) = \mathcal{Q}_l(q) + \mathcal{G}(x).\end{aligned}$$

We also define the projection $P_\Lambda : \Lambda_T \times E_T \longrightarrow \Lambda_T$.

Lemma 7 *For any l in the interval $(1, 1/\tau)$, the functional $\mathcal{L}_l$ is C^1 and satisfies the Palais-Smale condition. The critical points of $\mathcal{L}_l$ are the T-periodic smooth trajectories of the Lagrangian*

$$L_l(z, q, v) = a\left(v^2 - l^2\omega^2 q^2\right) + G(z, q, v), \quad (z, q, v) \in TM \times \mathbb{R} \times \mathbb{R}.$$

PROOF : We will often omit the subscript l in the following proof. Recall that Λ_T is a smooth manifold, and that the mappings

$$\begin{aligned}\exp_c : H^1(\mathcal{O}_c) &\longrightarrow H^1(S_T, M) \\ \xi(t) &\longmapsto \exp(\xi(t))\end{aligned}$$

are charts of this manifold, where $c \in C^\infty(S_T, M)$, $\mathcal{O}_c$ is a sufficiently small neighborhood of the zero section in the bundle c^*TM of tangents vectors of M along c, and $\exp : TM \longrightarrow M$ is the exponential map associated with some spray on M, see [20]. Let $\mu_T : \mathbb{R} \longrightarrow S_T$ be the natural projection, the induced vector bundle $\mu^* c^* TM$ is trivial since it is a vector bundle over $\mathbb{R}$, we have the commutative diagram

$$\begin{array}{ccccc} \mathbb{R} \times \mathbb{R}^n & \xrightarrow{\Phi} & \mu^* c^* TM & \xrightarrow{\tilde{\mu}} & c^* TM \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{R} & \xrightarrow{id} & \mathbb{R} & \xrightarrow{\mu} & S_T, \end{array}$$

where Φ is a vector bundle isomorphism and we define the covering

$$r_c = \tilde{\mu} \circ \Phi : \mathbb{R} \times \mathbb{R}^n \longrightarrow c^* TM.$$

A H^1 section $\xi : S_T \longrightarrow c^* TM$ has a unique lifting $\tilde{\xi} : \mathbb{R} \longrightarrow \mathbb{R}^n$ such that the diagram

$$\begin{array}{ccc} \mathbb{R} \times \mathbb{R}^n & \xrightarrow{r_c} & c^* TM \\ {\scriptstyle (id, \tilde{\xi})}\uparrow & & \uparrow{\scriptstyle \xi} \\ \mathbb{R} & \xrightarrow{\mu} & S_T \end{array}$$

commutes. Let us take a compact neighborhood U_c of the origin in $\mathbb{R}^n$ such that $\mathbb{R} \times U_c \subset r_c^{-1}(\mathcal{O}_c)$, and suppose without loss of generality that $\mathcal{O}_c = r_c(\mathbb{R} \times U_c)$. The mapping

$$\begin{aligned}\rho : H^1(c^* TM) &\longrightarrow H^1([0, T], \mathbb{R}^n) \\ \xi &\longmapsto \tilde{\xi}_{|[0,T]}\end{aligned}$$

is a linear isomorphism onto its image $T\tilde{H} \subset H^1([0,T],\mathbb{R}^n)$. We will also note ρ the mapping (ρ, id_{E_T}), and we call $\tilde{H}$ the set $\tilde{H} = T\tilde{H} \cap H^1([0,T],U_c)$. Let us define the smooth map

$$\begin{aligned}\tilde{L}_c : \mathbb{R}\times U_c \times \mathbb{R}^n \times \mathbb{R}\times\mathbb{R} &\longrightarrow \mathbb{R}\\ (t,\tilde{\xi},\nu,q,v) &\longmapsto L\big(\exp\circ r_c(t,\tilde{\xi})\,,\\ &\qquad d_1(\exp\circ r_c)(t,\tilde{\xi}).1 + d_2(\exp\circ r_c)(t,\tilde{\xi}).\nu\,,\ q\,,\ v\big),\end{aligned}$$

and the functional

$$\begin{aligned}\tilde{\mathcal{L}} : H^1([0,T],U_c)\times E_T &\longrightarrow \mathbb{R}\\ (\tilde{\xi}(t),q(t)) &\longmapsto \int_0^T \tilde{L}_c(t,\tilde{\xi}(t),\tilde{\xi}'(t),q(t),\dot{q}(t))\,dt,\end{aligned}$$

we have $\mathcal{L} = \tilde{\mathcal{L}}\circ\rho$. One can check from [HG4] and the expression of $\tilde{L}_c$ above that the estimates

$$\begin{aligned}|\tilde{L}_c(t,\tilde{\xi},\nu,q,v)| &\leqslant C(1+q^2+|\nu|^2+v^2)\\ \left|\left(\frac{\partial\tilde{L}_c}{\partial\tilde{\xi}},\frac{\partial\tilde{L}_c}{\partial q}\right)(t,\tilde{\xi},\nu,q,v)\right| &\leqslant C(1+q+|\nu|^2+v^2)\\ \left|\left(\frac{\partial\tilde{L}_c}{\partial\nu},\frac{\partial\tilde{L}_c}{\partial v}\right)(t,\tilde{\xi},\nu,q,v)\right| &\leqslant C(1+q^2+|\nu|+v)\end{aligned}$$

hold on $\mathbb{R}\times U_c\times\mathbb{R}^n\times\mathbb{R}\times\mathbb{R}$. These growth conditions imply by well-known results (see [22]) that $\tilde{\mathcal{L}}$, and thus $\mathcal{L}$, are continuously differentiable. We also have the local expression of the differential:

$$d\tilde{\mathcal{L}}(\tilde{\xi},q) = \int_0^T \frac{\partial\tilde{L}_c}{\partial\tilde{\xi}}d\tilde{\xi} + \frac{\partial\tilde{L}_c}{\partial\nu}(d\tilde{\xi})' + \frac{\partial\tilde{L}_c}{\partial q}dq + \frac{\partial\tilde{L}_c}{\partial v}(dq)'$$

and $d\mathcal{L}(\xi,q) = d\tilde{\mathcal{L}}(\tilde{\xi},q)\circ\rho$. Let us now prove that the Palais-Smale condition is satisfied. We take a Palais-Smale sequence (θ_n,q_n). The sequence

$$\mathcal{L}_l(\theta_n,q_n) = \mathcal{Q}_l(q_n) + \mathcal{G}(\theta_n,q_n)$$

is bounded. Since $\mathcal{Q}_l$ is a non-degenerate quadratic form, there exists an operator $A_l : E_T \longrightarrow E_T$ such that

$$d\mathcal{Q}_l(q).A_lq = |\mathcal{Q}_l(q)| \geqslant C\|q\|^2_{H^1}.$$

Let us now write using [HG4] and that $\big\|d\mathcal{L}(\theta_n,q_n)\big\| = \epsilon_n \longrightarrow 0$

$$\begin{aligned}\epsilon_n\|q_n\|_{H^1} \geqslant d\mathcal{L}_l(\theta_n,q_n)(0,A_lq_n) &= d\mathcal{Q}_l(q_n).A_lq_n + d\mathcal{G}(\xi_n,q_n).(0,A_lq_n)\\ &\geqslant C\|q_n\|^2_{H^1} + \int\frac{\partial G}{\partial q}\cdot A_lq_n + \frac{\partial G}{\partial v}\cdot A_l\dot{q}_n\\ &\geqslant C\|q_n\|^2_{H^1} - C'\|q_n\|_{W^{1,1}}\\ &\geqslant C\|q_n\|^2_{H^1} - C''\|q_n\|_{H^1}.\end{aligned}$$

It follows that the sequence $\|q_n\|_{H^1}$ is bounded. Plugging this into the action

$$C \geqslant \mathcal{L}_l(\theta_n,q_n) \geqslant \int G + \mathcal{Q}_l(q_n) \geqslant b\int d^2(\partial\theta_n,z_0) - C\|q_n\|^2_{H^1}$$

yields that $\int \|\partial\theta_n\|^2$ is also bounded. By a standard application of the theorem of Ascoli, see [20], Lemma 1.4.4, we can find a C^0-convergent subsequence of θ_n, and by extracting another subsequence we can obtain that q_n also has a uniform limit. From now on, we will suppose that

$$(\theta_n, q_n) \xrightarrow{C^0} (\theta, q).$$

It remains to prove that the limit holds in $\Lambda \times E_T$, that is in H^1-norms. Since the continuous limit θ can be approximated by a smooth curve c, all the curves θ_n lie in a single chart $\exp_c$ of Λ for n large enough. We call ξ_n the local representatives of θ_n, and we can use the local expressions given above. It is useful to define the mapping

$$\begin{aligned} \Omega_c : \mathbb{R} \times U_c \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} &\longrightarrow \mathbb{R} \times U_c \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \\ (t, \tilde{\xi}, \nu, q, v) &\longmapsto \left(t, \tilde{\xi}, \frac{\partial \tilde{L}_c}{\partial \nu}, q, \frac{\partial \tilde{L}_c}{\partial v} \right). \end{aligned}$$

It is straightforward from the explicit expression of $\tilde{L}_c$ and from [HL] that Ω_c is a diffeomorphism, and the estimate

$$\frac{1}{C}|X| - C \leqslant \Omega_c(t, X) \leqslant C(|X| + 1)$$

is a consequence of [HG4]. A theorem of Krasnoselskii implies that the mapping

$$\begin{aligned} \Upsilon_c : L^2([0,T], \mathbb{R}^{2n+2}) &\longrightarrow L^2([0,T], \mathbb{R}^{2n+2}) \\ X(t) &\longmapsto \Omega_c(t, X(t)) \end{aligned}$$

is a homeomorphism. It is not hard to see that the sequence

$$\left(\frac{\partial \tilde{L}_c}{\partial \tilde{\xi}}(\tilde{\xi}_n, \tilde{\xi}'_n, q_n, \dot{q}_n), \frac{\partial \tilde{L}_c}{\partial \tilde{q}}(\tilde{\xi}_n, \tilde{\xi}'_n, q_n, \dot{q}_n) \right)$$

is bounded in $L^1([0,T], \mathbb{R}^{n+1})$, thus its zero averaged primitive $P_n \in W^{1,1}([0,T], \mathbb{R}^{n+1})$ has a subsequence that is convergent in $L^2([0,T], \mathbb{R}^{n+1})$. We suppose that

$$P_n \xrightarrow{L^2} P.$$

Since $\|d\mathcal{L}(\xi_n, q_n)\| \longrightarrow 0$, we have $\|d\tilde{\mathcal{L}}(\tilde{\xi}_n, q_n)_{|T\tilde{H}}\| \longrightarrow 0$, and the inequality

$$\left| \int \langle \dot{P}_n, (d\tilde{\xi}, dq) \rangle + \frac{\partial \tilde{L}_c}{\partial \nu} d\tilde{\xi}' + \frac{\partial \tilde{L}_c}{\partial v} d\dot{q} \right| \leqslant \epsilon_n \|(d\tilde{\xi}, dq)\|_{H^1}$$

holds for all variations $(d\tilde{\xi}, dq) \in H_0^1([0,T], \mathbb{R}^n) \subset T\tilde{H}$. The sequence

$$m_n = \frac{1}{T} \int_0^T \left(\frac{\partial \tilde{L}_c}{\partial \nu}(\tilde{\xi}_n(t), \tilde{\xi}'_n(t), q_n(t), \dot{q}_n(t)), \frac{\partial \tilde{L}_c}{\partial v}(\tilde{\xi}_n(t), \tilde{\xi}'_n(t), q_n(t), \dot{q}_n(t)) \right) dt$$

is bounded, and we can suppose taking a subsequence that it has a limit m. Integrating by parts in the inequality above yields

$$\left\langle \left(\frac{\partial \tilde{L}_c}{\partial \nu}, \frac{\partial \tilde{L}_c}{\partial v} \right) - P_n - m_n \, , \, (d\tilde{\xi}', d\dot{q}) \right\rangle_{L^2} \leqslant \epsilon_n \|(d\tilde{\xi}', d\dot{q})\|_{L^2}$$

and we obtain

$$\left\| \left(\frac{\partial \tilde{L}_c}{\partial \nu}, \frac{\partial \tilde{L}_c}{\partial v} \right) - P_n - m_n \right\|_{L^2} \leqslant \epsilon_n.$$

We thus have

$$\left(\frac{\partial \tilde{L}_c}{\partial \tilde{\xi}}(\tilde{\xi}_n, \tilde{\xi}'_n, q_n, \dot{q}_n), \frac{\partial \tilde{L}_c}{\partial \tilde{q}}(\tilde{\xi}_n, \tilde{\xi}'_n, q_n, \dot{q}_n) \right) \xrightarrow{L^2} P - m,$$

and the sequence

$$\left(\tilde{\xi}_n, \tilde{\xi}'_n, q_n, \dot{q}_n \right) = \Upsilon_c^{-1} \left(\tilde{\xi}_n \, , \, \frac{\partial \tilde{L}_c}{\partial \tilde{\xi}}(\tilde{\xi}_n, \tilde{\xi}'_n, q_n, \dot{q}_n) \, , \, q_n \, , \, \frac{\partial \tilde{L}_c}{\partial \tilde{q}}(\tilde{\xi}_n, \tilde{\xi}'_n, q_n, \dot{q}_n) \right)$$

has a limit in L^2. The sequence $(\tilde{\xi}_n, q_n)$ thus has a limit in $H^1([0,T], \mathbb{R}^{n+1})$, and the sequence $(\xi_n, q_n) = \rho^{-1}(\tilde{\xi}_n, q_n)$ has a limit in $H^1(\mathcal{O}_c) \times E_T$. □

We now have to study the topology of the functional. Let us define a group Γ of admissible deformations of A_T:

Definition 3 *A homeomorphism* $h : A_T \longrightarrow A_T$

- $T_s(\theta, q) = 0$ for all q such that $\|q\| \geqslant c$ and all s,
- $T_s(\theta_0, q) = 0$ for all q and all s,

and we can apply Dold's fixed point transfer, see [14] and [19], page 433, that asserts the injectivity of the morphism $P_\Lambda^* : H^*(\sigma, \theta_0) \longrightarrow H^*(F_s, (\theta_0, 0))$. We now take $s = 1$ and have the commutative diagram

$$\begin{array}{ccc} H^*(I_1, (\theta_0, 0)) & \xrightarrow{h^*} & H^*(F_1, (\theta_0, 0)) \\ {\scriptstyle P_\Lambda^*}\uparrow & & \uparrow{\scriptstyle P_\Lambda^*} \\ H^*(h.\sigma, \theta_0) & & H^*(\sigma, \theta_0) \\ {\scriptstyle i_{h.\sigma}^*}\uparrow & & \uparrow{\scriptstyle i_\sigma^*} \\ H^*(\Lambda, \theta_0) & \xrightarrow{h_\Lambda^*} & H^*(\Lambda, \theta_0), \end{array}$$

where h_Λ^* is the isomorphism that makes the following diagram commute

$$\begin{array}{ccc} H^*(\Lambda \times E, (\theta_0, 0)) & \xrightarrow{h^*} & H^*(\Lambda \times E, (\theta_0, 0)) \\ {\scriptstyle P_\Lambda^*}\uparrow & & \uparrow{\scriptstyle P_\Lambda^*} \\ H^*(\Lambda, \theta_0) & \xrightarrow{h_\Lambda^*} & H^*(\Lambda, \theta_0). \end{array}$$

Coming back to the first diagram, we see that $i_{h.\sigma}^*$ can not be zero because $P_\Lambda^* \circ i_\sigma^* \circ h_\Lambda^*$ is nonzero. □

For all G satisfying [HR1-4] and all $l \in (1, 1 + 1/\tau)$ we define

$$c_T^G(l) = \inf_{\sigma \in \Sigma} \inf_{h \in \Gamma} \sup \mathcal{L}_l \,|_{h(\sigma \times E^-)}$$

We have the estimate:

Lemma 9 *If G satisfies (2) then the inequality*

$$I_T(U) \leqslant c_T^G(l) \leqslant I_T(W)$$

holds.

Proof : Since $G \longrightarrow c_T^G(l)$ is an increasing function this is an easy consequence of the following lemma.

Lemma 10 *For all U satisfying [HU1-3], we have*

$$c_T^U(l) = I_T(U).$$

Proof : Recall that

$$\begin{aligned} c_T^U(l) &= \inf_{\sigma \in \Sigma} \inf_{h \in \Gamma} \sup_{(z,x) \in h(\sigma \times E^-)} \mathcal{Q}_l(x) + \mathcal{U}(z) \\ I_T(U) &= \inf_{\sigma \in \Sigma} \sup_{z \in \sigma} \mathcal{U}(z) \end{aligned}$$

We can take $h = Id$ in the definition of c to obtain

$$c \leqslant \inf_{\sigma\in\Sigma} \sup_{(z,q^-)\in\Lambda\times E^-} \mathcal{Q}_l(q^-) + \mathcal{U}(z) = \inf_{\sigma\in\Sigma} \sup_{z\in\Lambda} \mathcal{U}(z) = I.$$

To obtain the other inequality, we apply Lemma 8 and get

$$\sup_{(z,x)\in h(\sigma\times E^-)} \mathcal{Q}_l(x) + \mathcal{U}(z) \geqslant \sup_{z\in h.\sigma} \mathcal{U}(z) \geqslant I.$$

□

We are now in a position to prove Proposition 2.

Proof of Proposition 2: First, notice that the third conclusion is a consequence of the two other ones since the only T periodic solution of L_l satisfying $\theta_T \equiv \theta_0$ is the constant curve $(\theta_0, 0)$, and has zero action, which is forbidden by the second conclusion since $c_T(l(T)) \geqslant I_T(U) > 0$. Let us now choose $l(T)$. The function $l \longrightarrow c_T(l)$ is non-increasing thus almost everywhere differentiable. Moreover, the inequality

$$\int_1^{1+1/\tau} c_T'(l)dl \geqslant I_T(U) - I_T(W)$$

holds and we can choose an $l(T)$ in the interval $(1, 1/\tau)$ such that

$$c' = |c_T'(l(T))| \leqslant \tau(I_T(W) - IT(U)).$$

Let us set $c = c_T(l(T))$ and recall that

$$\mathcal{L}_l(\theta, q) = \mathcal{L}(\theta, q) - al^2\omega^2\|q\|_2^2.$$

We shall prove that there exists a critical point $X_T = (\theta_T, q_T)$ of $\mathcal{L}_{l(T)}$ such that

$$2a\omega^2\|q_T\|_2^2 \leqslant 1 + c.'$$

Arguing by contradiction we assume that there is no critical point of $L_{l(T)}$ at level c satisfying $2a\omega^2\|q_T\|_2^2 \leqslant 1 + c'$. We can then find using a standard deformation argument an ϵ in the interval $(0, c/2)$ and a homeomorphism $h_0 \in \Gamma$ satisfying

$$\mathcal{L}_{l(T)}(h_0(X)) \leqslant \mathcal{L}_{l(T)}(X)$$

for all $X \in A_T$, and such that

$$\mathcal{L}_{l(T)}(h_0(X)) \leqslant c - \epsilon$$

for all $X = (\theta, q) \in A_T$ satisfying

$$\mathcal{L}_{l(T)}(X) \leqslant c + \epsilon \quad \text{and} \quad 2a\omega^2\|q\|_2^2 \leqslant c' + 1/2.$$

Let l_n be an increasing sequence converging to $l(T)$, and let $c_n = c_T(l_n)$ and $\mathcal{L}_n = \mathcal{L}_{l_n}$. We can choose $\sigma_n \in \Sigma$ and $h_n \in \Gamma$ such that

$$\sup \mathcal{L}_n \big|_{h_n(\sigma_n\times E^-)} \leqslant c_n + (l(T) - l_n)/10.$$

When n is large enough this implies

$$\begin{aligned}\mathcal{L}_{l(T)}\big|_{h_n(\sigma_n\times E^-)} \leqslant \mathcal{L}_n\big|_{h_n(\sigma_n\times E^-)} &\leqslant c_n + (l(T)-l_n)/10\\ &\leqslant c + (c'+1/10)(l(T)-l_n) + (l(T)-l_n)/10\\ &\leqslant c + (c'+1/5)(l(T)-l_n).\end{aligned}$$

Take a loop $X=(\theta,q)\in h_n(\sigma_n\times E^-)$, either

$$\mathcal{L}_{l(T)}(X) \leqslant c - (l(T)-l_n)/5$$

and

$$\mathcal{L}_{l(T)}(h_0(X)) \leqslant c - (l(T)-l_n)/5$$

or

$$\mathcal{L}_{l(T)} \geqslant c - (l(T)-l_n)/5.$$

In the second case,

$$\begin{aligned}(l(T)^2 - l_n^2)a\omega^2\|q\|^2 = \mathcal{L}_n(X) - \mathcal{L}_{l(T)}(X) &\leqslant c + (l(T)-l_n)(c'+1/5) - c + (l(T)-l_n)/5\\ &\leqslant (c'+1/2)(l(T)-l_n),\end{aligned}$$

thus

$$2a\omega^2\|q\|^2 \leqslant (l(T)+l_n)a\omega^2\|q\|^2 \leqslant c'+1/2,$$

and we get

$$\mathcal{L}_{l(T)}(h_0(X)) \leqslant c - \epsilon,$$

when n is large enough. We have seen that

$$\mathcal{L}_{l(T)}(h_0\circ h_n(\sigma_n\times E^-)) \leqslant c - (l(T)-l_n)/5,$$

which is a contradiction since $h_0\circ h_n\in\Gamma$. □

References

[1] Arnold V. I. : *Mathematical methods of classical mechanics,* Springer (1978).

[2] Benci V. - Giannoni F. : Homoclinic orbits on compact manifolds, J. Math. Anal. Appl. **157** (1991), 568-576.

[3] Bernard P. : Homoclinic orbitsin families of hypersurfaces with hyperbolic periodic orbits. Preprint CEREMADE.

[4] Bernard P. : Recherche variationnelle d'orbites homoclines dans les systèmes dynamiques hamiltoniens, thèse, 2000.

[5] Bolotin S.V. : Libration motions of natural dynamical systems, Vestnik Mosk. Univ. Ser I Mat. **6**(1978),72-77.

[6] Bolotin S.V. :The existence of homoclinic motions, Vestnik Mosk. Univ. Ser I Mat. Mekh. **6**(1983),98-102.

[7] Bott R. : Lectures on Morse theory, old and new, BAMS **7** (1982) , No. 2, 331-358.

[8] Buffoni B. - Séré E. : A global condition for quasi-random behavior in a class of conservative systems, Comm. Pure and App. Math. , **49** (1996), 285-305.

[9] Cielebak K.: Pseudoholomorphic curves and periodic orbits on cotangent bundles, J. Math. Pures Appl. (a) **73**(1994), 251-278.

[10] Cielebak K. - Séré E. : Pseudoholomorphic curves and multiplicity of homolcinic orbits, Duke Math. J.**77**(1995), 483-518.

[11] Cielebak K. - Séré E. : Pseudoholomorphic curves and the shadowing lemma, Duke Math. J.**98**(1999).

[12] Conley C. C. : On the ultimate behavior of orbits with respect to an unstable critical point I. oscillating, asymptotic and capture orbits, J. Diff. Eq. **5**(1969), 136-158.

[13] Coti Zelati V., Ekeland I., and Sere E. : A Variational Approach to Homoclinic Orbits in Hamiltonian Systems, Math. Ann. **288** (1990), 133-160.

[14] Dold A. : The fixed point transfer of fibre preserving maps, Math. Z. **148** (1976), 215-244.

[15] Devaney R. L. : Homoclinic orbits in hamiltonian systems, J. Diff. Eq. **21** (1976), 431-438.

[16] Felmer P. : Heteroclinic orbits for spatially periodic hamiltonian systems, Ann. Inst. H. Poincaré, Anal. Non Linéaire **8** (1991), 477-497.

[17] Fenichel N. : Persistence and smoothness of invariant manifolds for flows, Ind. Univ. Math. Jour. **26** (1971), 193-225.

[18] Hirsch M. W. - Pugh C. C. - Shub M. : *Invariant Manifolds*, Lecture Notes in Math. **583** (1977), Springer.

[19] Hofer H. - Viterbo C. : The Weinstein conjecture in cotangent bundles and related results, Ann. Scuola Normale Superiore di Pisa, Serie 5,**15** (1988), 411-445.

[20] Klingenberg W. :*lectures on closed geodesics,* Grundlehren der Math. Wiss. **230** (1978), Springer.

[21] Lerman L. M. : Hamiltonian systems with loops of a separatrix of a saddle-center, Selecta Math. Sov. **10** (1991), 297-306.

[22] Mawhin J. - Willem M. : *Critical point theory and Hamiltonian systems,* App. Math. Sciences **105** (1989), Springer.

[23] Mielke A. - Holmes P. and O'Reilly O. : Cascades of homoclinic orbits to, and chaos near, a Hamiltonian saddle-center, J. Dyn. Diff. Eq. **4**(1992), 95-126.

[24] Rabinowitz P. H. : *Minimax methods in critical point theory with epplications to differential equations*, CBMS, **65**, AMS, Providence (1986).

[25] Rabinovitz P. : Periodic and heteroclinic orbits for a periodic Hamiltonian system, Ann. Inst. H. Poincaré, Anal. Non Linéaire **6** (1989), 331-346.

[26] Ragazzo C. G. : Irregular dynamics and homoclinic orbits to hamiltonian saddle-centers, Comm. Pure Appl. Math., **49**(1996) 1-37.

[27] Struwe M. : *Variational Methods*, Springer, Second Edition, (1996).

[28] Sullivan D. : Differential forms and the topology of manifolds, Proc. Int. Conf. on Manifolds and Related Topics in Topology, Tokyo (1973), 37-49.

Patrick BERNARD, CEREMADE, Université Paris Dauphine,
Place du Maréchal de Lattre de Tassigny, 75775 Paris Cedex 16.
pbernard@clipper.ens.fr