The Eulerian limit for 2D statistical hydrodynamics.

Sergei B. Kuksin

Dedicated to Giovanni Jona-Lasinio on his 70-th birthday

Abstract

We consider the 2D Navier-Stokes system, perturbed by a random force, proportional to the square root of the viscosity, and study its solutions when the viscosity goes to zero. We prove that under this limit the Reynolds number grows to infinity, and the solutions converge in distribution to non-trivial stationary solutions of the (free) Euler equation. We study the convergence and the limiting solutions.

Contents

1 Introduction **1**

2 Preliminaries **5**

3 The results **8**
3.1 Tightness 8
3.2 Estimates for U_{ν_j} and U 9
3.3 The equation for U 10
3.4 A phase-space for the Euler equation 11
3.5 Homogeneous solution 12
3.6 One degenerate example 13

4 Proofs **14**
4.1 Proof of Theorem 3.1 14
4.2 Proof of Lemma 3.2 15
4.3 Proof of Lemma 3.5 17

1 Introduction

In this work we study the small-viscosity 2D Navier-Stokes (NS) system, perturbed by a small random force:

$$\dot{u} - \nu\Delta u + (u\cdot\nabla)u + \nabla p = \sqrt{\nu}\,\hat{\eta}(t,x), \quad 0 < \nu \le 1, \tag{1.1}$$

$$\operatorname{div} u = 0, \quad u = u(t,x), \quad p = p(t,x), \quad x \in T^2 = \mathbb{R}^2/(2\pi L\mathbb{Z}^2).$$

It is assumed that $\int u\,dx \equiv \int \hat\eta\,dx \equiv 0$, and that the force $\hat\eta(t,x)$ is a random field, white in time and smooth in x (see below (2.4)).

It is known that under certain nondegeneracy assumptions on the force, the equation (1.1), interpreted as a Markov system in the space H of square-integrable divergence-free vector fields $u(x)$, has a unique stationary measure μ_ν. Moreover, distribution $\mathcal{D}(u_\nu(t))$ of any solution of (1.1) converges to μ_ν exponentially fast (see below in section 2). Equation (1.1) has a stationary solution $U_\nu(t)$ such that $\mathcal{D}U_\nu(t) \equiv \mu_\nu$, and the exponential convergence implies that

$$\operatorname{dist}(\mathcal{D}u_\nu(t), \mathcal{D}U_\nu(t)) \le Ce^{-\sigma t}, \quad \sigma > 0, \tag{1.2}$$

for any solution u_ν. Well known calculations, based on the Ito formula (repeated in section 2) show that

$$\mathbb{E}\,\|U_\nu\|_1^2 = \frac{1}{2}B_0, \qquad \mathbb{E}\,\|U_\nu\|_2^2 = \frac{1}{2}L^{-2}B_1, \tag{1.3}$$

where B_0 and B_1 are explicit constants, calculated in terms of the force $\hat\eta$, and for $m = 0,1,2,\dots$ we denote by $\|\cdot\|_m$ the norm in the Sobolev space $H^m \subset H$. For $m = 0$ $\|\cdot\|_0$ is the L_2-norm, it will be denoted $|\cdot|$.

The estimates (1.3) imply that the averaged energy [1] $\mathbb{E}\,|U_\nu(t)|^2$ of the flow U_ν is of order L^2 (see below (2.15)). Hence, its Reynolds number is

$$R(U_\nu) = \frac{L(\mathbb{E}\,|U_\nu|^2)^{1/2}}{\nu} \sim \frac{L^2}{\nu}.$$

The fact that the energy of the flow is of order L^2 while its Reynolds number is of order $L^2\nu^{-1}$ makes our scaling of the NS system convenient to study the 2D turbulence. We note that the substitution $u(t) = \nu^{1/5}v(\tau)$, $\tau = \nu^{1/5}t$, transforms (1.1) to the equation

$$v_\tau - \delta\Delta v + (v\cdot\nabla)v + \nabla p' = \tilde\eta(\tau,x), \quad \delta = \nu^{4/5}, \tag{1.4}$$

where $\tilde\eta$ is a new random field, distributed as $\hat\eta$. Stationary solutions for (1.4) have energy of order $L^2\delta^{-1}$ and the Reynolds number of order $L^2\delta^{-3/2}$. So (1.1) is equivalent to the NS system (1.4), perturbed by a random force of order one, but the former scaling is more convenient than the latter.

Our goal in this work is to examine the inviscous limit of the process U_ν, i.e. its limit as $\nu \to 0$. The period L is a parameter of the problem which we keep fixed.

In Theorem 3.1 and Lemma 3.2 we show that any sequence $\tilde\nu_j \to 0$ contains a subsequence $\nu_j \to 0$ such that the process U_{ν_j} a.s. converges to a limiting process $U(\cdot)$. Its energy $|U(t)|^2$ and enstrophy $\|U\|_1^2 = |\operatorname{rot} U|^2$ are time-independent random constants:

$$|U(t)|^2 = E, \quad \|U(t)\|_1^2 = \Omega \qquad \forall\, t.$$

[1] more exactly, the doubled energy per unit area since $|U_\nu|^2 = (2\pi L)^{-2}\int |U_\nu|^2\,dx$.

The expectation of the enstrophy is known, while the expectation of the energy is bounded from below and from above:

$$\mathbb{E}\Omega = \frac{1}{2}B_0, \qquad \frac{L^2 B_0^2}{2B_1} \le \mathbb{E}E \le \frac{L^2 B_0}{2}. \tag{1.5}$$

Moreover, certain exponential moments of E and Ω are finite, and

$$\mathbb{E}\|U(t)\|_2^2 \le \frac{1}{2}L^{-2}B_1 \quad \forall t. \tag{1.6}$$

Crucial property of the process U, established in Theorem 3.4, is that it satisfies the Euler equation

$$\dot{U} + (U\cdot\nabla)U + \nabla p = 0, \quad \operatorname{div} U = 0, \quad \int U\,dx = 0. \tag{1.7}$$

Note that although U solves the free equation (1.7), relations (1.5) show that it remembers some characteristics of the force $\hat{\eta}$. In particular, U is not identically zero. In section 3.5 it is proved that if the force $\hat{\eta}(t,x)$ is stationary in x, then the process U has zero mean-value:

$$\mathbb{E}U(t,x) = 0 \qquad \forall\, t, x\,.$$

Due to (1.5) this implies that the process U is 'genuinely random'.

Due to (1.6), $U(0) \in H^2$ a.s. No existence and uniqueness theorem for the Euler equation with Cauchy data in H^2 is known. Still, as we show in section 3.4, estimates (1.5) and (1.6) imply that with probability one trajectories of the process U belong to a uniqueness class for the Euler equation, and that they are trajectories of a continuous dynamical system which the equation defines in a certain functional space H^E. Accordingly, the distribution $\mathcal{D}U(t)$ of the process U at any point t is an invariant measure for the Euler equation in the space H^E (see Theorem 3.6).

Jointly with (1.2) the established results mean that the double limit

$$\lim_{\nu_j\to 0}\lim_{T\to\infty} u_{\nu_j}(T+\cdot) = U(\cdot) \tag{1.8}$$

exists (the convergences are understood as $*$-weak convergences of distributions of random processes $\{u(t),\, t\ge 0\}$), and the limiting stationary process U is a random solution of the deterministic Euler equation (1.7). The limits in (1.8) do not commute. Indeed, when $\nu_j \to 0$, $u_{\nu_j}(t)$ converges to a deterministic solution $u_0(t)$ of (1.7). When $t\to\infty$ this solution has no reason to converge to a limit and to remember characteristics of the force $\tilde{\eta}$ (cf. (1.5)).

Our results show that the scaling (1.1) of the 2D NS system is the right one to study the 2D turbulence since under the limit $\nu \to 0$ stationary solutions of the equation converge (at least, along a subsequence) to a regular limit, while their Reynolds numbers diverge to infinity. In the 3D case the right scaling is different, if we believe that predictions of the Kolmogorov theory apply to space-periodic

turbulent flows. Indeed, let u_ν be a stationary (in time) solution of the 3D NS system (1.1) with $\sqrt{\nu}\,\hat\eta$ replaced by $\hat\eta$. [2] Then, applying the Ito formula, we get that $\nu\mathbb{E}\,\|u_\nu\|_1^2 = \frac{1}{2}B_0$ (cf. (1.3)). In particular, $\nu\mathbb{E}\,\|u_\nu\|_1^2$ (trivially) converges to a positive finite limit when $\nu \to 0$, as predicted by the Kolmogorov theory. So in the 3D case exactly this scaling (without the factor $\sqrt{\nu}$ in the r.h.s.) should be correct. We also note that our results apparently disagree with the Kraichnan theory of 2D turbulence, see [Kra67] and [KM80, Section 4.2]. Indeed, if u_ν is a stationary solution of (1.1), then $\mathbb{E}\,\|u_\nu(t)\|_2^2 = O(1)$ as $\nu \to 0$, see (1.3). In particular $\lim \nu\mathbb{E}\,\|u_\nu\|_2^2 = 0$. But it is postulated in the theory that this limit is positive and finite. [3]

Due to (1.8), solutions of the Euler equation approximate the velocity fields of 2D turbulent flows with hight Reynolds numbers. This property of the 2D turbulence was predicted by many physicists, cf. [Ons49, KM80]. Accordingly, one can try to use the Euler equation to calculate characteristics of solutions u_ν for (1.1) with $\nu \ll 1$. For example, let us take a continuous functional f on H, consider an observable variable $f(u_\nu(t))$ and try to find its average in ensemble. If we can use some ergodic arguments to calculate the averages $\langle f\rangle$ of f along trajectories of the Euler flow, then due to (1.8) we will have $\lim_{\nu_j\to 0}\lim_{t\to\infty}\mathbb{E}\,f(u_{\nu_j}(t)) = \langle f\rangle$, independently of the sequence ν_j. So $\langle f\rangle$ would approximate asymptotic in time value of $\mathbb{E}\,f(u_\nu(t))$ for solutions with small viscosity ν. As we show in Proposition 3.7, the quantities $I_g = \int g(\mathrm{rot}\,U(t,x))\,dx$ are integrals of motion for the limiting processes U. Hence, their trajectories cannot be ergodic in the codimension-two surfaces $H_{a,b} = \{|u| = a, \|u\|_1 = b\}$. So it is unlikely that the 'right' average $\langle f\rangle$ can be found, assuming equidistribution of trajectories on the surfaces $H_{a,b}$ (concerning such calculations see [KM80]). On the contrary, it is plausible that the energy $|U|^2$ and the quantities I_g give a complete list of integrals of motion for the process U. If so, then the theory of Robert (e.g., see in [RS94]) can be used to calculate the averages of f along the corresponding surfaces of infinite codimension. Next one can hope to integrate the resulting quantities in the parameter, which parameterizes the surfaces (at least, to integrate approximately). This would give us approximate value of the expectations $\mathbb{E}\,f(u_\nu(t))$ for large time t and small viscosity ν.

The approach to study limiting behaviour of the NS system, suggested in this work, applies to other damped/driven Hamiltonian PDEs with two or more integrals of motion. In particular, to the damped/driven nonlinear Schrödinger equation

$$\dot u - \nu\Delta u + i(\Delta u - |u|^2 u) = \sqrt{\nu}\,\eta(t,x), \quad \dim x \le 4.$$

Now the limiting processes U are non-trivial stationary solutions of the nonlinear Schrödinger equation, and the distributions $\mathcal{D}U(0)$ are its invariant measures. Details will be given in a joint paper with Armen Shirikyan which is now under preparation.

[2] We boldly believe that, as in the 2D case, such solutions exist.

[3] Note that for any other scaling of the 2D NS equation (e.g., for (1.4)) the limit $\lim_{\nu\to 0} u_\nu$ does not exist.

Notations. $\mathcal{P}(X)$ is the set of probability Borel measures on a metric space X, provided with the $*$-weak topology. $C_b(X)$ is the space of bounded continuous functions on X. By C, C_1, etc. we denote various ν-independent finite constants.

Acknowledgements. I thank Oliver Penrose and Peter Constantin for discussions, related to the statistics of the 2D turbulence and the 2D Euler equation, respectively. This research was supported by EPSRC, grant N63055/01.

2 Preliminaries

Let H be the Hilbert space of square-integrable divergence-free vector fields on T^2 with zero mean-value, given the scalar product $(u,v) = (2\pi L)^{-2}\int u\cdot v\,dx$ and the norm $|u| = (u,u)^{1/2}$. Let Π be L_2-orthogonal projector to H. Applying Π to (1.1) we get

$$\dot u(t) + \nu Au(t) + B(u(t),\, u(t)) = \nu^{1/2}\eta, \tag{2.1}$$

where $A = -\Pi\Delta$, $B(u,u) = \Pi(u\cdot\nabla)u$ and $\eta = \Pi\hat\eta$. In (2.1) we view u and η as curves in H. See e.g. [CF88, Gal01].

For $\nu = 0$ equation (1.1) becomes the Euler equation (1.7). Application of the projector Π transforms it to the equation 2.1 with $\nu = 0$:

$$\dot u(t) + B(u(t),\, u(t)) = 0.$$

Let $\{e_s, s\in\mathbb{Z}^2_0 := \mathbb{Z}^2\backslash\{0\}\}$ be the Hilbert basis of H, formed by trigonometric vector fields:

$$\{e_s,\, e_{-s}\} = \{c_s s^\perp \cos L^{-1}s\cdot x,\; c_s s^\perp \sin L^{-1}s\cdot x\}\quad \forall\, s\in\mathbb{Z}^2_0,$$

where $s^\perp = \begin{pmatrix}-s_2\\ s_1\end{pmatrix}$ for any vector $\begin{pmatrix}s_1\\ s_2\end{pmatrix}\in\mathbb{Z}^2_0$, and $c_s = \sqrt{2}\,|s|^{-1}L^{-1}$ is the L_2-normalising factor. Then

$$Ae_s = \alpha_s e_s,\quad \alpha_s = L^{-2}|s|^2\quad \forall\, s.$$

For any $r\ge 0$ we define the Sobolev space H^r as $H^r = \{u\in H \mid \|u\|_r = |A^{r/2}u| < \infty\}$. In particular, $H^0 = H$ and $\|\cdot\|_0 = |\cdot|$. For $r<0$ we set H^r to be equal to the completion of H in the norm $\|\cdot\|_r$, defined as above.

Since for $u = \sum u_s e_s$ we have $\|u\|_1^2 = \sum \alpha_s |u_s|^2$ and $\alpha_s \ge L^{-2}$ for each s, then

$$\|u\|_1 \ge L^{-1}|u|. \tag{2.2}$$

The norms $\|\cdot\|_j$ satisfy the interpolation inequalities. In particular,

$$\|u\|_1^2 \le |u|\cdot\|u\|_2. \tag{2.3}$$

The random force η is assumed to be

$$\eta(t) = \frac{d}{dt}\zeta(t),\quad \zeta = \sum_{s\in\mathbb{Z}^2_0} b_s\beta_s(t)e_s(x), \tag{2.4}$$

where $\{\beta_s\}$ are independent standard Wiener processes, defined for $t \in \mathbb{R}$, and $\{b_s \geq 0\}$ are constants. For $j = 0, 1, \dots$ we set

$$B_j = \sum_{s \in \mathbb{Z}_0} |s|^{2j} b_s^2 \leq \infty.$$

Everywhere below we assume that

$$B_0, B_1 < \infty. \tag{2.5}$$

Integrating (2.1) from τ to $T > \tau$ we get

$$u(T) + \int_\tau^T (\nu Au + B(u,u))\, dt = u(\tau) + \sqrt{\nu}\, \zeta(T) - \sqrt{\nu}\, \zeta(\tau). \tag{2.6}$$

An a.s. continuous random process $u(t) \in H$, $t \geq \tau$, such that its norm $\|u(t)\|_1$ is square-integrable on every finite time-interval is called a *solution* for (2.1), if the relation (2.6) holds a.s. for each $T > \tau$. Here both parts of (2.6) are understood as elements of H^{-1}. It is known (see e.g. [VF88]) that for any $u_0 \in H$ the equation (2.1) has a unique solution $u(t)$, $t \geq \tau$, equal u_0 for $t = \tau$.

A random process $\tilde{u}(t) \in H_0$, $t \geq \tau$, is called a *weak solution* for (2.1) if one can find a process $\tilde{\zeta}(t) \in H$, distributed as $\zeta(t)$, such that $\tilde{u}$ is a solution for (2.1) with ζ replaced by $\tilde{\zeta}$. Its distribution $\mathcal{D}(\tilde{u}(\cdot))$ (i.e. a measure in $C([\tau, \infty);\ H)$) also is called a week solution, cf. [VF88, Chapter X]. A solution $u(t)$ (weak or strong), defined for all t, is called a *stationary solution* if u is a stationary process in H. For a random variable $v \in H$ with a finite second moment, independent of the process $\eta(t)$, $t \geq 0$, we denote by $u(t; v)$, $t \geq 0$, a strong solution for (2.1), equal v at $t = 0$ (such a solution exists, see [VF88]). A measure $\mu \in \mathcal{P}(H)$ is called a *stationary measure* for (2.1) if

$$\mathcal{D}(v) = \mu \Rightarrow \mathcal{D}u(t; v) = \mu \quad \forall\, t \geq 0.$$

If $u(\cdot)$ is a stationary solution, then the distribution $\mathcal{D}u(t)$, $t \in \mathbb{R}$, is a (t-independent) stationary measure in H. Conversely, if μ is a stationary measure, then there exists a (weak) stationary solution u such that $\mathcal{D}u(t) \equiv \mu$ (e.g., see in [KS02]).

It is known that every equation (2.1) (satisfying (2.5)) has a stationary measure μ, see [VF88]. Its uniqueness is proven under the additional assumption that

$$b_s \neq 0 \quad \forall\, |s| \leq N_\nu, \tag{2.7}$$

where N_ν is a sufficiently large number. See [KS00, KS02] (also see [EMS01, BKL02] and the review [Kuk02]). Moreover, under the assumption (2.7),

$$|\mathbb{E} f(u(t;\ v)) - \int f(u)\, \mu du| \leq C_\nu (1 + \mathbb{E}\, |v|^2) e^{-\sigma_\nu t},\ \sigma_\nu > 0, \tag{2.8}$$

for any bounded Lipschitz functional f on H such that $|f| \leq 1$ and $\mathrm{Lip}(f) \leq 1$. Let us introduce in $\mathcal{P}(H)$ the distance

$$\mathrm{dist}(\mu_1,\ \mu_2) = \sup(\mu_1 - \mu_2,\ f),$$

where the supremum is taken over all f as above. It is known that $(\mathcal{P}(H),\,\mathrm{dist})$ is a complete metric space, and that convergence with respect to this distance is equivalent to the $*$-weak convergence of measures, see [Dud89]. We can rewrite (2.8) as

$$\mathrm{dist}(\mathcal{D}u(t;\ v),\,\mu) \le C_\nu(1+\mathbb{E}\,|v|^2)e^{-\sigma_\nu t}. \tag{2.9}$$

We note that if

$$b_s \neq 0 \quad \forall\, s, \tag{2.10}$$

then the assumption (2.7) holds for each $\nu>0$, so for any $\nu>0$ equation (2.1) has a unique stationary measure μ_ν.

Lemma 2.1. *If* (2.10) *holds, then the measure* $\mu_\nu\in\mathcal{P}(H)$ *continuously depends on* $\nu>0$.

Proof. Due to (2.9) it is sufficient to check that $\mathcal{D}u_\nu(t)\in\mathcal{P}(H)$ continuously depends on ν for each $t\ge 0$, where $u_\nu(\cdot)$ is a solution of (2.1), vanishing at $t=0$. To verify this property we make in (2.1), (2.4) the substitution $u=\nu U(\tau\nu)$ and get for U the equation

$$U_\tau + AU + B(U,\,U) = \nu^{-5/2}\frac{\partial}{\partial\tau}\zeta(\nu\tau) =: \frac{\partial}{\partial\tau}\zeta_\nu(\tau).$$

To prove the continuity we may assume that $\nu>\delta>0$. Then $0\le\tau\le t/\nu< T=t/\sigma$. It is easy to check that a solution $U\in C([0,\,T];\ H)=\mathcal{H}$ continuously depends on $\zeta_\nu\in C([0,\,T];\ H^1)=\mathcal{H}^1$ (e.g., see Lemma 4.2 in [KS03]). Since a.s. $\zeta\in C([0,\,\infty),\,H^1)$, then a.s. $\zeta_\nu\in\mathcal{H}^1$ continuously depends on $\nu\in[\delta,\,1]$, as well as $U\in\mathcal{H}$. As $u_\nu(t)=\nu U(t\nu^{-1})$, then a.s. $u_\nu(t)\in H$ continuously depends on ν. Hence, $\mathcal{D}u_{\nu'}(t)\to\mathcal{D}u_\nu(t)$ as $\nu'\to\nu$, and the lemma is proven. □

Let $u(t)$ be a stationary solution of (2.1) (weak or strong). Due to (2.5) the Ito formula applies to the functional $|u(t)|^2$. Since $(B(u,u),\,u)=0$, then taking expectation of the formula we get that

$$\mathbb{E}\,|u(t)|^2 - \mathbb{E}\,|u(0)|^2 = -2\nu\int_0^t \mathbb{E}\,(Au,\ u)\,d\tau + \nu\sum_s\int_0^t b_s^2|e_s|^2\,d\tau,$$

(see [VF88, EMS01, Shi02]). As u is a stationary process, then this relation implies that

$$\mathbb{E}\|u(\tau)\|_1^2 = \frac{1}{2}B_0 \quad \forall\,\tau. \tag{2.11}$$

Since $u=u(x)$ is a divergence-free vector field, then

$$\|u\|_1^2 = (2\pi L)^{-2}\int_{T^2}|\operatorname{rot} u|^2dx.$$

So the l.h.s. of (2.11) is the averaged enstrophy of u (per unit area).

Similar, applying the Ito formula to the enstrophy functional $f(u)=\|u\|_1^2$ and using that $(B(u,u),\,Au)=0$, we obtain

$$\mathbb{E}\|u(\tau)\|_2^2 = \frac{1}{2}L^{-2}B_1 \quad \forall\,\tau. \tag{2.12}$$

Two useful inequalities below follow by applying the Ito formula to the functionals $\exp(\gamma|u|^2)$ and $\exp(\gamma\|u\|_1^2)$, where

$$\gamma = (2L^2 b_{max}^2)^{-1}, \quad b_{max} = \max\{b_s,\, s \in \mathbb{Z}_0^2\}.$$

Namely,

$$\mathbb{E}\exp(\gamma|u(\tau)|^2) \le \mathfrak{B}_0 = B_0 \exp\left(\frac{B_0+1}{2b_{max}^2}\right),$$

$$\mathbb{E}\exp(\gamma\|u(\tau)\|_1^2) \le \mathfrak{B}_1 = L^{-2}B_1 \exp\left(\frac{L^{-2}B_1+1}{2b_{max}^2}\right). \tag{2.13}$$

See [VF88, p. 395] and [Shi02]. Due to (2.13),

$$\mathbb{E}\|u(\tau)\|_1^p \le C_p \gamma^{-p/2} \mathfrak{B}_1 \tag{2.14}$$

for any $p \ge 1$.

Equalities (2.11) and (2.12) imply lower and upper bounds for $\mathbb{E}|u(\tau)|^2$. Indeed, taking the expectation of (2.3) we find that

$$\mathbb{E}\|u(\tau)\|_1^2 \le (\mathbb{E}|u(\tau)|^2)^{1/2} (\mathbb{E}\|u(\tau)\|_2^2)^{1/2}.$$

Now (2.11), (2.12) imply a lower bound for $\mathbb{E}|u(\tau)|^2$. Combining it with an upper bound, which follows from (2.2) and (2.11), we get

$$\frac{1}{2}\frac{L^2 B_0^2}{B_1} \le \mathbb{E}|u(\tau)|^2 \le \frac{L^2 B_0}{2}. \tag{2.15}$$

3 The results

3.1 Tightness

The equality (2.12) and the Prokhorov theorem immediately imply that the set $\{\mu_\nu, \nu \in (0,1]\}$ of stationary measures for equation (2.1) is tight in $H^{2-\epsilon}$ for any $\epsilon > 0$. Jointly with Lemma 3.2 it means that if $b_s \ne 0$ for all s, then the measures $\mu_\nu \in \mathcal{P}(H^{2-\epsilon})$ continuously depend on $\nu \in (0,1]$.

What is much more important, distributions $m_\nu = \mathcal{D}(u_\nu(\cdot)) \in \mathcal{P}(C(\mathbb{R};\, H))$ of the weak stationary solutions u_ν for (2.1) also are tight. Below we prove the tightness for the most important case $\nu \to 0$.

Let us fix any $\varepsilon \in (0,\, 1)$ and define the following spaces of trajectories:

$$\begin{aligned} Z &= L_{2\,\mathrm{loc}}(\mathbb{R},\, H^{2-\varepsilon}) \cap C(\mathbb{R},\, H^{-1-\varepsilon}), \\ Z_n &= L_2([-n,\, n],\, H^{2-\varepsilon}) \cap C([-n,\, n],\, H^{-1-\varepsilon}), \quad n \in \mathbb{N}. \end{aligned}$$

The norms in the spaces Z_n define a countable system of semi-norms $[\cdot]_n$ in Z, and define there the distance

$$\mathrm{dist}(u,\, v) = \sum 2^{-n} \frac{[u-v]_n}{[u-v]_n + 1},$$

cf. [VF88, p. 340]. All the spaces Z_n and Z are Polish (i.e., complete and separable).

Theorem 3.1. *Any sequence* $\{m_{\tilde\nu_j}\}$, $\tilde\nu_j \to 0$, *contains a converging subsequence*

$$m_{\nu_j} \to m \in \mathcal{P}(Z). \tag{3.1}$$

The theorem is proven in section 4.1.

The measures m_{ν_j} are stationary since they are distributions of stationary processes, so the limiting measure m is stationary as well. By the Skorokhod theorem (see [IW89, section I.2], we can find random processes $U_{\nu_j}(t)$, $j \geq 1$, and $U(t)$, defined on the same probability space, such that

$$\mathcal{D}U_{\nu_j} = m_{\nu_j} \quad \forall\, j, \quad \mathcal{D}U = m,$$

and

$$U_{\nu_j} \to U \quad \text{in} \quad Z \quad \text{a.s.} \tag{3.2}$$

Below we study properties of the limiting random process U.

3.2 Estimates for U_{ν_j} and U

For any $N \geq 1$ we consider the projection

$$P_N : H \to H, \quad \sum_{s\in\mathbb{Z}^2_0} u_s e_s \mapsto \sum_{|s|\leq N} u_s e_s,$$

and define the functionals

$$f(u) = e^{\gamma\|u\|_1^2}, \quad f_N(u) = e^{\gamma\|P_N u\|_1^2 \wedge N}, \quad N \geq 1.$$

Clearly f is continuous on H^1, the functionals f_N are bounded continuous on H^{-2} and

$$0 < f_N(u) \nearrow f(u) \leq \infty \quad \forall\, u \in H^{-2}. \tag{3.3}$$

Everywhere below in this section

$$\nu \in \{\nu_1, \nu_2, \dots\},$$

where $\nu_1, \nu_2, \dots$ is the sequence in (3.1).

Due to (3.2), for each $\tau \in \mathbb{R}$ we have $\mathbb{E}f_N(U_{\nu_j}(\tau)) \to \mathbb{E}f_N(U(\tau))$ as $j \to \infty$. Hence, $\mathbb{E}f_N(U(\tau)) \leq \mathfrak{B}_1$ for each N, due to (2.13). Evoking (3.3) and the Levi theorem we get that

$$\mathbb{E}\exp(\gamma\|U(\tau)\|_1^2) \leq \mathfrak{B}_1 \quad \forall\, \tau. \tag{3.4}$$

Similar,

$$\mathbb{E}\exp(\gamma|U(\tau)|^2) \leq \mathfrak{B}_0 \quad \forall\, \tau. \tag{3.5}$$

A remarkable property of the process U is that a.s. its energy $|U(t)|^2$ and enstrophy $\|U(t)\|_1^2$ are time-independent random constants:

Lemma 3.2. *Almost surely we have*

$$|U(t)|^2 = E \quad \text{and} \quad \|U(t)\|_1^2 = \Omega \quad \text{for a. a.} \quad t \in \mathbb{R}. \tag{3.6}$$

The expectations of the random constants E and Ω satisfy

$$\mathbb{E}\,\Omega = \frac{1}{2}B_0, \quad \frac{L^2 B_0^2}{2B_1} \le \mathbb{E}\,E \le \frac{L^2 B_0}{2}. \tag{3.7}$$

Besides,

$$\mathbb{E}\,\|U(\tau)\|_2^2 \le \frac{1}{2}L^2 B_1 \quad \forall\,\tau. \tag{3.8}$$

The lemma is proven below in section 4.2.

3.3 The equation for U

Since U_ν, $\nu \in \{\nu_j\}$, is a weak solution for (2.1), then for any $T > \tau$ it satisfies (2.6), where ζ is replaced by a process $\tilde{\zeta}$ with the same distribution. We rewrite (2.6) as follows:

$$\begin{aligned} \Psi(U_\nu) \;&:=\; U_\nu(T) - U_\nu(\tau) + \int_\tau^T B(U_\nu,\, U_\nu)dt = \\ &=\; -\nu\int_\tau^T AU_\nu(s)\,ds + \sqrt{\nu}(\tilde{\zeta}(T) - \tilde{\zeta}(\tau)) =: G_\nu. \end{aligned} \tag{3.9}$$

Lemma 3.3. *The nonlinearity B defines continuous quadratic maps $H^2 \to H^1$ and $H^1 \to L_p = L_p(T^2, \mathbb{R}^2) \cap H$, for any $p < 2$.*

The proof of the lemma's assertions is obvious since the multiplication of functions defines continuous bi-linear maps $H^2(T^2) \times H^1(T^2) \to H^1(T^2)$ and $H^1(T^2) \times L_2(T^2) \to L_p(T^2)$, and since the projection Π is continuous in the L_p-spaces with $1 < p < \infty$.

Due to the lemma, Ψ defines a continuous map $Z \to H_{-1-\varepsilon}$. So $\Psi(U_{\nu_j}) \to \Psi(U)$ in $H_{-1-\varepsilon}$ as $\nu_j \to 0$, almost surely. From the other hand, since

$$\mathbb{E}\,|G_\nu| \le C\sqrt{\nu}$$

due to (2.12) and (2.4), then $G_{\nu_j} \to 0$ in $H_{-1-\varepsilon}$ in probability. Passing in (3.9) to the limit in probability as $\nu_j \to 0$ we get that $\Psi(U) = 0$ a.s. That is, almost surely

$$\dot{U}(t) + B(U(t),\, U(t)) = 0. \tag{3.10}$$

So, a.s. $U(\cdot) \in Z$ is a weak solution of the Euler equation (3.10), equivalent to (1.7). Due to (3.6), a.s. $\|U(t)\|_1 = \text{const}$. Using (3.8) and the Fubini theorem we get that

$$U(\cdot) \in L_{2\,\mathrm{loc}}(\mathbb{R};\; H^2) \quad \text{a.s.}\,. \tag{3.11}$$

Therefore due to Lemma 3.3 and (3.10),

$$\dot{U} \in L_{1\,\mathrm{loc}}(\mathbb{R};\; H^1) \cap L_\infty(\mathbb{R};\; L_p) \quad \text{a.s.}, \quad \forall\, p < 2. \tag{3.12}$$

We have proven the following result:

Theorem 3.4. *Any limiting measure $m \in \mathcal{P}(Z)$ as in* (3.1) *is a weak stationary solution for the Euler equation* (3.10) *in the following sense: There exists a stationary random process $U(t) \in H$, distributed as m, which a.s. satisfies* (3.10). *Moreover,*

1) energy $|U(t)|^2$ of this process a.s. is a time-independent random constant E such that

$$\frac{L^2 B_0^2}{2B_1} \le \mathbb{E}\, E \le \frac{L^2 B_0}{2},$$

2) its enstrophy $\|U(t)\|_1^2$ a.s. is a time-independent random constant Ω such that $\mathbb{E}\Omega = \frac{1}{2}B_0$;

3) the process satisfies the estimate (3.8), *the exponential estimates* (3.4) *and* (3.5), *and its time-derivative satisfies* (3.12).

3.4 A phase-space for the Euler equation

By (3.11) and (3.12), with probability one trajectories of the process U belong to the space

$$\mathcal{K} = \{u \in L_{2\,\mathrm{loc}}(\mathbb{R};\ H^2) \mid \dot{u} \in L_{1\,\mathrm{loc}}(\mathbb{R};\ H^1)\}.$$

Since $\mathcal{K} \subset C(\mathbb{R}; H^1)$, then due to the lemma below $\mathcal{K}$ is an uniqueness class for the Euler equation.

Lemma 3.5. *Let $u(t)$ and $v(t)$, $t \in [t_1, t_2]$, be two solutions of the Euler equation* (3.10) *such that*

$$\|u(t)\|_1 + \|v(t)\|_1 \le C_1 \quad \forall\, t, \tag{3.13}$$

$$\int_{t_1}^{t_2} \|u(t)\|_2^2\, dt \le C_2, \tag{3.14}$$

and u equals v at some point $t_3 \in [t_1, t_2]$. Then $u(t) \equiv v(t)$.

The lemma is proved in section 4.3.

Let us denote by $\mathcal{K}^E$ the set of trajectories in $\mathcal{K}$ which satisfy the Euler equation. Clearly $\mathcal{K}^E$ is a closed subset of $\mathcal{K}$, invariant under the time-translations $u(\cdot) \to u(T + \cdot)$, $T \in \mathbb{R}$. Due to Lemma 3.5 the map

$$\pi : \mathcal{K} \to H^1, \quad u \mapsto u(0),$$

restricted to $\mathcal{K}^E$, defines a continuous embedding $\mathcal{K}^E \to H^1$. We denote by H^E its image $\pi(\mathcal{K}^E)$, and provide H^E with the topology, induced from $\mathcal{K}^E$.

Due to [BKM84], for any $u_0 \in H^3$ the Euler equation has a unique solution $u \in \mathcal{K} \cap L_{\infty\,\mathrm{loc}}(\mathbb{R};\ H^3)$, equal u_0 at $t = 0$. This solution, as a point in $\mathcal{K}$, continuously depends on u_0. So

$$H^3 \subset H^E \subset H^1, \tag{3.15}$$

and the inclusions are continuous. We point out that we do not know if the (topological) space H^E is linear, or not.

For any $u_0 \in H^E$, $u = \pi^{-1}(u_0)$ is the unique solution of the Euler equation in then space $\mathcal{K}$, equal u_0 at $t = 0$. Since $u(t) = \pi(u(t + \cdot))$ and the time-translation maps are continuous in $\mathcal{K}$, then the Euler equation defines a group of continuous automorphisms $S_t : H^E \to H^E$, $t \in \mathbb{R}$. As $S_t \circ U(\tau) = U(t + \tau)$ and the process U is stationary, then we have:

Theorem 3.6. *The distribution $\mathcal{D} = \mathcal{D}\,U(0)$ of the process U as in Theorem 3.4 is an invariant Borel measure for the flow which the Euler equation defines in H^E. The space H^E satisfies* (3.15)*, and $\mathcal{D}(H^E \cap H^2) = 1$.*

Since vector fields from H^2 have the modulus of continuity [4] $g(r) = r \max(1, \log r^{-1})$, then any $u(t, x) \in \mathcal{K}$ has the modulus of continuity in x, equal to $\phi(t)g(r)$, where $\phi \in L_{2\,\mathrm{loc}}$. Therefore by the Osgood criterion [Har64], the differential equation

$$\dot{x}(t) = u(t, x), \quad x(t_1) = y,$$

has a unique solution. So the flow-maps $S_{t_1}^{t_2} : T^2 \to T^2$, $y \mapsto x(t_2)$, are well defined and continuous (see [Che98]). Accordingly, due to the classical arguments, for any $u \in \mathcal{K}^E$ we have $\omega(t, x) \equiv \omega_0(S_t^0(x))$, where $\omega(t, x) = \mathrm{rot}\, u(t, x)$ (and $\omega_0 = \mathrm{rot}\, u_0$).

Approximating $u(t, x)$ by smooth divergence-free vector fields $u_\epsilon(t, x)$ (say, obtained as convolutions with scalar mollifiers), we approximate any continuous flow-map $S = S_{t_1}^{t_2}$ by measure-preserving smooth maps $T^2 \to T^2$, converging to S at each point, see [Har64, Theorem 2.1, Section V]. Passing to the limit as $\epsilon \to 0$, we find that $\int_{T^2} f(S(x))\,dx = \int_{T^2} f(x)\,dx$ for each continuous function f. Hence, the flow-maps preserve the Lebesgue measure, and

$$\int_{T^2} g(\omega(t, x))\,dx = \int_{T^2} g(\omega_0 \circ S_t^0)(x)\,dx = \int_{T^2} g(\omega_0(x))\,dx\,,$$

for any t and any $g \in C_b(\mathbb{R})$. That is,

$$\int_{T^2} g(\mathrm{rot}\, U(t, x))\,dx = \mathrm{const} \quad \forall\, t \in \mathbb{R}, \tag{3.16}$$

for each $U \in \mathcal{K}^E$. In particular, we have the following result:

Proposition 3.7. *For a.a. trajectory of the process U and any bounded continuous function g, the integral in* (3.16) *is a time-independent random constant.*

3.5 Homogeneous solution

Let us assume that

$$b_s = b_{-s} \neq 0 \quad \forall\, s \in \mathbb{Z}_0^2. \tag{3.17}$$

[4] Indeed, by the classical embedding theorem, spaces H^r with non-integer $r > 1$ are embedded in Hölder spaces C^{r-1}. So by the interpolation, the space H^2 is embedded in the interpolating Hölder space C_*^1 (the Sigmund space). The latter is formed by functions, which have the modulus of continuity $g(r)$, see [Che98, p. 31].

Then the random field $\zeta(t,\,x)$ is homogeneous in x, i.e. translations of x do not change its distribution (see [Kuk02]), and (2.1) has a unique stationary measure μ_ν. As a consequence, the measure μ_ν and a corresponding stationary solution $U_\nu(t,\,x)$ both are homogeneous in x, see [Kuk02]. Since U_ν a.s. converges to the field $U(t) = U(t,x)$ as in Theorem 3.4, then U is homogeneous (i.e., stationary in x).

That is, under the assumption (3.17) the limiting measures m as in (3.1) correspond to random solutions of (1.7) which are stationary both in time and space. In particular, since $\int U\,dx \equiv 0$, then in this case we have

$$\mathbb{E}\,U(t,x) \equiv 0$$

(i.e., $\mathbb{E}U(t) = 0 \in H^{-1-\varepsilon}$ for all t).

3.6 One degenerate example

Let us consider a force η of the form (2.4), such that $b_s = 0$ if $|s| > 1$. Then $B_0 = B_1$, so by Theorem 3.4 we have

$$\mathbb{E}\,E = \frac{L^2 B_0}{2} = L^2 \mathbb{E}\,\Omega. \tag{3.18}$$

Let us denote $s_1 = (1,\,0)^t$ and $s_2 = (0,\,1)^t$. Due to (2.2), $|u|^2 \le L^2 \|u\|_1^2$, and the equality holds if and only if $u \in M^4$, where

$$M^4 = \operatorname{span}\{e_{\pm s_1}, e_{\pm s_2}\} = \Big\{ C_1 s_1 \sin\big(a_1 + \frac{x_2}{L}\big) + \\ C_2 s_2 \sin\big(a_2 + \frac{x_1}{L}\big) \mid C_1, C_2 \in \mathbb{R},\ a_1, a_2 \in \mathbb{R}/2\pi\mathbb{Z} \Big\}.$$

So (3.18) implies that a limiting solution $U(t)$ as in Theorem 3.4 belongs to M^4 for all t. For any $u \in M^4$ we have

$$(u\cdot\nabla)u = L^{-1} C_1 C_2 \Big(s_1 \cos\big(a_1 + \frac{x_2}{L}\big) \sin\big(a_2 + \frac{x_1}{L}\big) + \\ s_2 \sin\big(a_1 + \frac{x_2}{L}\big) \cos\big(a_2 + \frac{x_1}{L}\big) \Big) = -C_1 C_2 \nabla\big(\cos(a_1 + \frac{x_2}{L}) \cos(a_2 + \frac{x_1}{L})\big), \tag{3.19}$$

so $B(u,u) = 0$. That is, $\dot U \equiv 0$ and U is a constant process

$$U(t) = C_1 s_1 \sin\big(a_1 + \frac{x_2}{L}\big) + C_2 s_2 \sin\big(a_2 + \frac{x_1}{L}\big) \quad \forall\, t, \tag{3.20}$$

where C_1, C_2 and a_1, a_2 are random variables.

The processes U above are very degenerate because due to (3.19), the 4-dimensional space M^4 is invariant for equation (2.1) with $\eta = 0$. The space M^4 is a maximal one with this property since if M is a linear subspace of H, invariant for (2.1) with $\eta = 0$, and $M^4 \subset M$, $M^4 \ne M$, then M is dense in H, see [EM01]. Accordingly, if $b_s \ne 0$ for $|s| \le 2$, then the corresponding processes U have no chances to be as degenerate as (3.20).

4 Proofs

4.1 Proof of Theorem 3.1

Our arguments in this section closely follow [VF88], Chapter X.

First we note that to prove the theorem it is sufficient to check that for any $n \geq 1$ the sequence of measures $\{m^n_{\bar\nu_j},\ j = 1, 2, \dots\}$ is tight in Z^n, where

$$m^n_{\bar\nu_j} = \mathcal{D}(u_{\bar\nu_j}|_{[-n,\,n]}) \in \mathcal{P}\big(C([-n,\,n];\ H)\big).$$

Indeed, we need to verify that the sequence $\{m_{\bar\nu_j}\}$ contains a subsequence $\{m_{\nu_j}\}$ such that

$$(m_{\nu_j},\, f) \to (m,\, f), \quad m \in \mathcal{P}(Z), \tag{4.1}$$

for any $f \in C_b(Z)$. It is known that it suffices to check (4.1) for bounded Lipschitz functionals f_L (see [Dud89], Theorem 11.3.3). It is easy to approximate any such f_L by functionals $f^n_L = g^n_L \circ \pi_n$, where π_n is the natural projection $Z \to Z_n$, and g^n_L is a bounded Lipschitz functional on Z_n. Since $(m_\nu,\, g^n_L \circ \pi_n) = (m^n_\nu,\, g^n_L)$, then (4.1) holds if

$$m^n_{\nu_j} \to m^n \quad \forall\, n, \tag{4.2}$$

where $m^n = (\pi_n)_* m$. If we know that the sequence $\{m^n_{\bar\nu_j}\}$ is tight in Z_n for each n, then we can use the diagonal process to construct a subsequence $\{\nu_j\}$ such that (4.2) holds for every n. The measures $m^n \in \mathcal{P}(Z_n)$ form a compatible family and define a measure $m \in \mathcal{P}(Z)$ that satisfies (4.1)

So we need to check that the sequence of measures $\{\mu^n_{\bar\nu_j}\}$ is tight for any fixed n. Let u_ν be a stationary solution for (2.1) such that $\mathcal{D}(u_\nu) = m_\nu$. We abbreviate $u = u_\nu$ and take any $t_1,\, t_2 \in [-n,\, n]$, satisfying $t_1 \leq t_2$ and $|t_1 - t_2| \leq 1$. We have

$$u(t_2) - u(t_1) = -\nu \int_{t_1}^{t_2} Au(s)\, ds - \int_{t_1}^{t_2} B(u(s),\, u(s))\, ds + \sqrt{\nu}\,(\zeta(t_2) - \zeta(t_1)) \tag{4.3}$$

(strictly speaking, since u is a weak solution, then ζ should be replaced by a suitable process $\tilde\zeta$, distributed as ζ. This inaccuracy makes no difference for what follows).

The H-norm of the first integral in the r.h.s. of (4.3) is bounded by

$$\nu \left(\int_{t_1}^{t_2} \|u\|_2\, ds \right) \leq \nu |t_2 - t_1|^{\frac12} \Big(\int_{t_1}^{t_2} \|u\|_2^2\, ds \Big)^{\frac12} \leq \nu |t_2 - t_1|^{\frac12} \Big(\int_{-n}^{n} \|u\|_2^2\, ds \Big)^{\frac12}.$$

Due to (2.12),

$$\mathbb{E} \Big(\int_{-n}^{n} \|u\|_2^2 \Big)^{\frac12} \leq (nL^{-2} B_1)^{\frac12}.$$

Since $\|B(u,\, u)\|_{-1} \leq C \|u\|_1^2$, then H^{-1}-norm of the second integral is bounded by

$$|t_2 - t_1| \sup_{[t_1,\, t_2]} \|u(t)\|_1^2 \leq |t_2 - t_1| \sum_{m=-n}^{n-1} \sup_{[m,\, m+1]} \|u(t)\|_1^2.$$

An application of the Ito lemma to $\|u(t)\|_1^2$ (cf. (4.6) in the next subsection) bounds the expectation of oscillation of $\|u(t)\|_1^2$ on any segment $[m, m+1]$ by $C\sqrt{\nu}$. So due to (2.11),

$$\mathbb{E} \sum_{m=-n}^{n-1} \sup_{[m,m+1]} \|u(t)\|_1^2 \le C_1 n.$$

Finally, due to basic properties of the Brownian motion, the H-norm of the third term in the right-hand side of (4.3) is bounded by

$$\sqrt{\nu}\,|t_2 - t_1|^{\frac{1}{3}}\xi,$$

where $\xi \ge 0$ and $\mathbb{E}\xi \le C_2(n)$.

Let us consider the Hölder space $C^{1/3}\left([-n,\, n],\, H^{-1}\right) =: \mathcal{C}_n^{1/3}$. Due to (2.11) and the estimates for the norms of the three terms in the r.h.s. of (4.3),

$$\mathbb{E}\,|u_\nu\,|_{\mathcal{C}_n^{1/3}} \le C(n). \tag{4.4}$$

Let Y_n be the space $Y_n = \mathcal{C}_n^{1/3} \cap L_2\left([-n,\, n],\, H^2\right)$. Since

$$\mathbb{E}\,|u_\nu\,|^2_{L_2([-n,\,n],\,H^2)} = nL^{-2}B_1$$

by (2.12), then using (4.4) we have

$$\mathbb{E}\,|u_\nu\,|_{Y_n} \le C'(n) \quad \forall\, \nu.$$

As the space Y_n is compactly embedded in the separable space Z_n ([VF88], Lemma X.3.1), then the family of measures $\mu_\nu^n = \mathcal{D}u_\nu|_{[-n,\,n]}$ is tight in Z_n by the Prokhorov theorem. Theorem 3.1 is proven. □

4.2 Proof of Lemma 3.2

Let us apply the Ito formula to the functional $\xi_\nu(t) = \|U_\nu(t)\|_1^2$. For any $t_1 \le t_2$ we have:

$$\begin{aligned}\xi_\nu(t_2) - \xi_\nu(t_1) &= -2\nu \int_{t_1}^{t_2} \|U_\nu(t)\|_2^2\, dt \\ &+ \nu \int_{t_1}^{t_2} \sum_s b_s^2 (e_s,\, Ae_s)\, dt + 2\sqrt{\nu} \sum_s b_s \int_{t_1}^{t_2} (U_\nu(t),\, Ae_s)\, d\beta_s(t).\end{aligned}$$

Noting that the second term in the r.h.s. equals $\nu L^{-2} B_1 (t_2 - t_1)$, we see that

$$\text{osc}\ \xi_\nu|_{[0,\,1]} := \sup_{0 \le t_1 \le t_2 \le 1} |\xi_\nu(t_1) - \xi_\nu(t_2)| \le \nu L^{-2} B_1 +$$

$$2\nu \int_0^1 \|U_\nu(t)\|_2^2\, dt \;+\; 4\sqrt{\nu} \sup_{0 \le \tau \le 1} \left| \int_0^\tau (AU_\nu(t),\, \sum b_s e_s d\beta_s(t)) \right|. \tag{4.5}$$

Writing $U_\nu(t)$ as $\sum U_{\nu\, s}(t)e_s$ and applying the Burkholder–Devis–Gundi inequality, we see that the expectation of the third term in the r.h.s. of (4.5) is bounded by

$$4\sqrt{\nu}\,\mathbb{E}\left(\sum_{s\in Z_0^2}\int_0^1 \alpha_s^2 b_s^2 U_{\nu s}(t)^2 dt\right)^{1/2} \leq 4\sqrt{\nu}\, b_{max}\left(\mathbb{E}\int_0^1 \|U_\nu(t)\|_2^2 dt\right)^{1/2}$$
$$= 4\sqrt{\nu}\, b_{max} L^{-1}\sqrt{\frac{B_1}{2}},$$

where to get the last equality we used (2.12). Using (2.12) once again to estimate the expectation of the second term in the r.h.s. of (4.5) we get that

$$\mathbb{E}\,(\mathrm{osc}\;\; \xi_\nu|_{[0,\,1]}) \leq C\sqrt{\nu}. \tag{4.6}$$

Let us denote by $\mathcal{E}$ and g the following functionals on Z:

$$\mathcal{E}(u) = \int_0^1 \|u(t)\|_1^2\, dt, \quad g(u) = \int_0^1 \Big|\, \|u(t)\|_1^2 - \mathcal{E}(u)\Big|\, dt.$$

By $\mathcal{E}_N$ and g_N, $N \in \mathbb{N}$, we denote their regularized versions:

$$g_N(u) = g(u) \wedge N, \quad \mathcal{E}_N(u) = \mathcal{E}(u) \wedge N.$$

All these functionals are continuous non-negative, and

$$g_N \nearrow g, \quad \mathcal{E}_N \nearrow \mathcal{E} \quad \text{as} \quad N \to \infty.$$

Due to (2.11),

$$\mathbb{E}\,\mathcal{E}(U_\nu) = \int_0^1 \mathbb{E}\,\|U_\nu(t)\|_1^2\, dt = \frac{1}{2}B_0. \tag{4.7}$$

Besides, due to (2.14) we have

$$\mathbb{E}\,\mathcal{E}(U_\nu)^2 = \mathbb{E}\left(\int_0^1 \|U_\nu\|_1^2\, dt\right)^2 \leq \mathbb{E}\int_0^1 \|U_\nu\|_1^4\, dt \leq C'.$$

Therefore,

$$\mathbb{E}\,\mathcal{E}(U_\nu) I_{\mathcal{E}(U_\nu)>M} \leq \frac{1}{M}\mathbb{E}\,\mathcal{E}(U_\nu)^2 \leq M^{-1}C'.$$

So the system of random variables $\{\mathcal{E}(U_\nu)\}$ is uniformly integrable, and (3.2) implies that

$$\mathbb{E}\,\mathcal{E}(U) = \lim \mathbb{E}\,\mathcal{E}(U_\nu) = \frac{1}{2}B_0,$$

see [Dud89, Theorem 10.3.6]. Since the process $U(t)$ is stationary, then

$$\mathbb{E}\,\mathcal{E}(U) = \mathbb{E}\int_0^1 \|U(t)\|_1^2\, dt = \mathbb{E}\,\|U(t)\|_1^2.$$

That is,

$$\mathbb{E}\,\|U(t)\|_1^2 = \frac{1}{2}B_0 \quad \forall\, t \in [0,1]. \tag{4.8}$$

Now we return to estimate (4.6). As $|\xi_\nu(t) - \int_0^1 \xi_\nu(\tau)d\tau| \le \operatorname{osc}\ \xi_\nu|_{[0,\,1]}$ for $0 \le t \le 1$, then due to (4.6) we have

$$0 \le \mathbb{E}\, g_N(U_\nu) \le \mathbb{E}\, g(U_\nu) \le C\sqrt{\nu}.$$

Passing to the limit as $\nu_j \to 0$ using (3.2), we get that $\mathbb{E}\, g_N(U) = 0$ for each N. That is, $g(U) = 0$ a.s., and $\|U(t)\|_1^2 = \mathcal{E}(U)$ a.e. in $[0,\,1]$. Repeating these arguments for all half-integer time-intervals $\left[\frac{n}{2},\, \frac{n}{2}+1\right]$, $n \in \mathbb{Z}$, we get that a.s.

$$\|U(t)\|_1^2 = \Omega \quad \text{for a.a.} \quad t \in \mathbb{R}. \tag{4.9}$$

Here Ω is a random constant which can be defined, say, as

$$\Omega = \mathcal{E}(U) = \int_0^1 \|U(t)\|_1^2\, dt.$$

Using (4.8) we see that expectation of Ω satisfies the first relation in (3.7).

Applying the Ito formula to the functional $|U_\nu|^2$, we get a bound for oscillation of $|U_\nu(t)|^2$, analogous to (4.5). Next we repeat derivation of (4.9) to find that the energy of the process $U(t)$ a.s. is a time-independent random constant E.

Passing to the limit in (2.12) using (3.2) and the Levi theorem we get (3.8) (cf. the derivation of (3.4)). Repeating the arguments that proved (2.15) and using (4.8) and (3.8), we find that

$$\frac{L^2 B_0^2}{2B_1} \le \mathbb{E}\, E \le \frac{L^2 B_0}{2}.$$

Now the lemma is proven. □

4.3 Proof of Lemma 3.5

Our proof of the lemma is a variation of the classical arguments due to Yudovitch [Yud63] (also see in [Che98]). To simplify notations we assume that $t_1 = t_3 = 0$ and $t_2 = T > 0$. Below the constants C, C_1, etc. are independent of the parameter $p > 2$.

Let us denote $b(u_1, u_2, u_3) = (B(u_1, u_2), u_3)$ and $w = u - v$. Subtracting from the equation for u the equation for v, multiplying the result in H by w and using that $b(v, w, w) = 0$ we get:

$$\frac{1}{2}\frac{d}{dt}|w|^2 + b(w, u, w) = 0. \tag{4.10}$$

Abbreviating $b(w, u, w) = b$ and applying the Hölder inequality, we get

$$|b| \le C|\nabla u|_p |w|_{2q}^2, \quad \frac{1}{p} + \frac{1}{q} = 1, \quad p > 2 \tag{4.11}$$

(here and below $|\cdot|_r$ stands for the L_r-norm). It is known that the Sobolev embedding theorem for the space H^1 can be specified as follows:

$$|V|_p \le C\sqrt{p}\,\|V\|_1 \quad \forall\, V \in H^1, \tag{4.12}$$

where $C = C(L)$, see [LL97, Theorem 8.5]. [5] Therefore,

$$|\nabla u|_p \le C\sqrt{p}\,\|u\|_2.$$

To estimate $|w|_{2q}$ we write

$$\int |w|^{2q}\,dx = \int |w|^a |w|^{2q-a}\,dx \le |w|_2^a |w|_{p(2q-a)}^{2q-a}, \quad a = \frac{2}{q},$$

where we used the Hölder inequality with $P = 2/a = q$, $Q = p$. Since $\|w\|_1 \le \|u\|_1 + \|v\|_1$, then due to (3.13), (4.12) we have

$$|w|_{2q}^2 \le C|w|_2^{2q^{-2}} \big(C_1^2 p(2q-a)\big)^{\frac{2q-a}{2q}}.$$

Noting that $0 < \frac{2q-a}{2q} = \frac{q+1}{pq} \le \frac{2}{p}$, we obtain

$$|w|_{2q}^2 \le C'|w|_2^{2q^{-2}} p^{2/p} \le C|w|_2^{2q^{-2}}.$$

So

$$|b| \le C\sqrt{p}\,\|u(t)\|_2 |w|_2^{2q^{-2}}. \tag{4.13}$$

Let us denote $g = |w|_2^2$. Relations (4.10) and (4.13) imply the differential inequality

$$\dot g(t) \le C\sqrt{p}\; g(t)^{q^{-2}} \|u(t)\|_2\,.$$

As $g(0) = 0$ and $1 - q^{-2} = (q+1)/pq$, then applying the Gronwall lemma we get an upper bound for g:

$$\begin{aligned} g(t) &\le \left(C(1-q^{-2})\sqrt{p} \int_0^t \|u(s)\|_2\,ds \right)^{\frac{pq}{q+1}} \\ &\le \left(C_1 p^{-1/2}\sqrt{t}\,\big(\int_0^t \|u(s)\|_2^2\,ds \big)^{1/2} \right)^{\frac{pq}{q+1}}. \end{aligned}$$

Using (3.14) and passing to the limit as $p \to \infty$, we see that $g(t) = 0$ for all $0 \le t \le T$, as stated.

[5] In the statement of the theorem in [LL97] the value of the constant that bounds the norm of the Sobolev embedding is given incorrectly, but right value of the constant immediately follows from the proof (which is correct).

References

[BKL02] J. Bricmont, A. Kupiainen, and R. Lefevere, *Exponential mixing for the 2D stochastic Navier–Stokes dynamics*, Comm. Math. Phys. **230** (2002), no. 1, 87–132.

[BKM84] J. Beale, T. Kato, and A. Majda, *Remarks on the breakdown of smoothness for the 3-D Euler equations*, Comm. Math. Phys. **94** (1984), 61–66.

[CF88] P. Constantin and C. Foias, *Navier-Stokes Equations*, University of Chicago Press, Chicago, 1988.

[Che98] J.-Y. Chemin, *Perfect Incompressible Fluids*, Oxford University Press, Oxford, 1998.

[Dud89] R. M. Dudley, *Real Analysis and Probability*, Wadsworth & Brooks/ Cole, Pacific Grove, California, 1989.

[EM01] W. E and J. C. Mattingly, *Ergodicity for the Navier–Stokes equation with degenerate random forcing: finite dimensional approximation*, Comm. Pure Appl. Math **54** (2001), 1386–1402.

[EMS01] W. E, J. C. Mattingly, and Ya. G. Sinai, *Gibbsian dynamics and ergodicity for the stochastically forced Navier–Stokes equation*, Comm. Math. Phys. **224** (2001), 83–106.

[Gal01] G. Gallavotti, *Foundations of Fluid Dynamics*, Springer-Verlag, Berlin, 2001.

[Har64] Ph. Hartman, *Ordinary Differential Equations*, John Wiley & Sons, New York, 1964.

[IW89] N. Ikeda and S. Watanabe, *Stochastic Differential Equations and Diffusion Processes*, North-Holland, Amsterdam, 1989.

[KM80] R. H. Kraichnan and D. Montgomery, *Two-dimensional turbulence*, Rep. Prog. Phys. **43** (1980), 547–619.

[Kra67] R. H. Kraichnan, *Inertial ranges in two-dimensional turbulence*, Phys. Fluids **10** (1967), 1417–1423.

[KS00] S. B. Kuksin and A. Shirikyan, *Stochastic dissipative PDE's and Gibbs measures*, Comm. Math. Phys. **213** (2000), 291–330.

[KS02] S. B. Kuksin and A. Shirikyan, *Coupling approach to white-forced nonlinear PDE's*, J. Math. Pures Appl. **81** (2002), 567–602.

[KS03] S. B. Kuksin and A. Shirikyan, *Some limiting properties of randomly forced 2D Navier–Stokes equations*, Proceedings A of the Royal Society of Edinburgh (2003), to appear.

[Kuk02] S. B. Kuksin, *Ergodic theorems for 2D statistical hydrodynamics*, Rev. Math. Phys. **14** (2002), 585–600.

[LL97] E. H. Lieb and M. Loss, *Analysis*, Graduate Studies in Mathematics, vol. 14, AMS, Providence, 1997.

[Ons49] L. Onsager, *Statistical hydrodynamics*, Nuovo Cimento Suppl. **6** (1949), 279–287, also in "The collected works of Lars Onsager", World Scientific, 1996.

[RS94] R. Robert and J. Sommeria, *Statistical equilibrium states for two-dimensional flows*, J. Fluid Mech. **229** (1994), 291–310.

[Shi02] A. Shirikyan, *Analyticity of solutions for randomly forced two-dimensional Navier–Stokes equations*, Russian Math. Surveys **57** (2002), 785–799.

[VF88] M. I. Vishik and A. V. Fursikov, *Mathematical Problems in Statistical Hydromechanics*, Kluwer, Dordrecht, 1988.

[Yud63] V. I. Yudovich, *Non stationary flow of an ideal incompressible fluid (in Russ.)*, Zh. Vych. Mat. **3** (1963), 1032–1066.

Sergei B. Kuksin
Department of Mathematics
Heriot-Watt University
Edinburgh EH14 4AS
Scotland, UK
and
Steklov Institute of Mathematics
8 Gubkina St., 117966 Moscow
Russia

E-mail: kuksin@ma.hw.ac.uk

web-address: www.ma.hw.ac.uk/∼kuksin/