CONVERGENCE OF SPECTRA OF GRAPH-LIKE THIN MANIFOLDS

PAVEL EXNER AND OLAF POST

Abstract. We consider a family of compact manifolds which shrinks with respect to an appropriate parameter to a graph. The main result is that the spectrum of the Laplace-Beltrami operator converges to the spectrum of the (differential) Laplacian on the graph with Kirchhoff boundary conditions at the vertices. On the other hand, if the the shrinking at the vertex parts of the manifold is sufficiently slower comparing to that of the edge parts, the limiting spectrum corresponds to decoupled edges with Dirichlet boundary conditions at the endpoints. At the borderline between the two regimes we have a third possibility when the limiting spectrum can be described by a nontrivial coupling at the vertices.

1. Introduction

Graph models of quantum systems have a long history. Already half a century ago Ruedenberg and Scherr [RueS53] used this idea to calculate spectra of aromatic carbohydrate molecules, however, a real boom started from the late eighties when semiconductor graph-type structures became small and clean enough so that coherent effects in the corresponding quantum transport can play the dominating role. From the mathematical point of view these models were analyzed first thoroughly in [EŠ89], for recent developments and bibliography see [KoS99].

The free Hamiltonian of a graph model is the (differential) Laplacian on the graph. To define it properly one has to specify the boundary conditions which couple the wave functions at the vertices. They have to define a self-adjoint operator, however, this requirement itself does not specify the conditions uniquely: in a vertex joining n graph edges we have n^2 free parameters. A natural idea to remove this non-uniqueness is to regard the graph model as a limit case of a more realistic one with a unique Hamiltonian; an appropriate choice is a "thickened graph" composed of thin tubes which gives the original graph in the limit of a vanishing tube radius. Unfortunately, it is not easy to see what happens with spectral and/or scattering properties in such a limit. The spectral convergence when the "thick graph" is planar with Neumann boundary conditions has been solved recently by Kuchment–Zeng [KuZ01], and Rubinstein–Schatzman [RubS01]; Saito [Sa00] showed the convergence of the resolvent. Note that Colin de Verdière [CdV86] already established a similar result to prove that the first

Date: December 10, 2003.

non-zero eigenvalue of a compact manifold of dimension greater than 2 can have arbitrary high (finite) multiplicity. The physically more interesting situation with Dirichlet boundary represents a longstanding open problem.

These two cases do not exhaust all possible ways in which a family of manifolds can approach a graph. One more choice are manifolds without a boundary of codimension one in $\mathbb{R}^\nu$, $\nu \geq 3$, which encloses the graph like a system of "sleeves"[1], with the limit consisting of the sleeve diameter shrinking. It is particularly interesting from the viewpoint of recent efforts to build circuits based on carbon nanotubes. Recall that recently discovered techniques — see, e.g., [AMSC01, PRL$^+$00, TBG$^+$02] — allow to fabricate branched nanotubes and thus in principle objects very similar to the mentioned "sleeved graphs".

In this paper we consider a more abstract setting of the problem which covers the "strip graphs" of [KuZ01, RubS01] and their generalizations to higher dimensions, as well as the "sleeved graphs" described above.

Let us briefly describe the structure of the paper. In Section 2 we define the Laplacian on a graph and give an abstract eigenvalue comparision tool (Lemma 2.1). In Section 3 we define the graph like manifolds associated to a graph. In Subsection 3.E we motivate the four different limiting procedures on the vertex neighbourhoods discussed in Sections 5 – 8. In Section 4 we define the limit procedure of the edge neighbourhoods which remain the same in all cases. In the last section (Sec. 9 we give an application on the spectral convergence result in the case of periodic graphs.

2. Preliminaries

2.A. **Laplacian on a graph.** Suppose M_0 is a finite connected graph with vertices v_k, $k \in K$ and edges e_j, $j \in J$. Suppose furthermore that e_j has length $\ell_j > 0$, i.e., $e_j \cong I_j := [0, \ell_j]$. We clearly can make M_0 into a metric measure space with measure given by $p_j(x)dx$ on the edge e_j where $p_j \colon I_j \longrightarrow (0,\infty)$ is a smooth density function for each $j \in J$. We then have

$$L_2(M_0) = \bigoplus_{j\in J} L_2(I_j, p_j(x)dx)$$

$$\|u\|^2_{M_0} = \sum_{j\in J} \|u_j\|^2_{I_j} = \sum_{j\in J} \int_{I_j} |u_j(x)|^2 p_j(x)dx.$$

We let $\mathcal{H}^1(M_0)$ be the completion of

$$\{\, u \in C(M_0) \,|\, u_j := u\restriction_{e_j} \in C^1(I_j) \,\}$$

[1]Kuchment and Zeng [KuZ01] speak also about sleeves having meaning graph edges thickened into strips; what we have in mind here is rather a cylindrical surface with the graph edge as its axis.

where the closure is taken with respect to the norm

$$\|u\|^2_{1,M_0} := \sum_{j\in J}(\|u_j\|^2_{I_j} + \|u'_j\|^2_{I_j}).$$

Note that the weakly differentiable functions $\mathcal{H}^1(I_j)$ on an interval are continuous, therefore $\mathcal{H}^1(I_j) \subset C(I_j)$.

Next we associate with the graph a positive quadratic form,

$$\|u'\|^2_{M_0} := \sum_{j\in J} \|u'_j\|^2_{I_j}$$

for all $u \in \mathcal{H}^1(M_0)$. It allows us to define the *(differential) Laplacian on the (weighted) graph* M_0 as the unique self-adjoint and non-negative operator Δ_{M_0} associated with the closed form $u \mapsto \|u'\|^2_{M_0}$ (see [K66, Chapter VI], [RS-1] or [Dav96] for details on quadratic forms). In other words, the operator and the quadratic form are related by

$$\|u'\|^2_{M_0} = \langle \Delta_{M_0} u, u\rangle \tag{2.1}$$

for $u \in C^1(M_0)$ belonging to the domain of Δ_{M_0}. On the edge e_j, the operator Δ_{M_0} is given formally by

$$\Delta_{M_0} u = -\frac{1}{p_j(x)}(p_j(x)u'_j)'. \tag{2.2}$$

Note that the domain of Δ_{M_0} consists of all functions $u \in C(M_0)$ which are twice weakly differentiable on each edge. Furthermore, each function u satisfies (weighted) *Kirchhoff boundary conditions*[2] at each vertex v_k, i.e.,

$$\sum_{j,\ e_j \text{ meets } v_k} p_j(v_k)u'_j(v_k) = 0 \tag{2.3}$$

for all $k \in K$ where the derivative is taken on each edge in the direction away from the vertex. In particular, we assume Neumann boundary conditions at a vertex with only one edge emanating.[3] If we assume that p is continuous on M_0, we can omit the factors $p_j(v_k)$ in (2.3). Note that different values of $p_j(v_k)$ for j can correspond in our limiting result to different radii of the thickened edges which are attached to a vertex neighbourhood (see (4.2) below).

As we have mentioned in the introduction there are other self-adjoint operators which act according to (2.2) on the graph edges but satisfy different boundary conditions at the vertices — see [EŠ89, KoS99] for details. The corresponding

[2]This is the usual terminology, not quite a fortunate one. The name suggests that the probability current at the vertex obeys the conservation law analogous to Kirchhoff's law in an electric circuit. While this claim is valid, the current conservation requirement is equivalent to selfadjointness and thus also satisfied for the other operators mentioned below.

[3]This hypothesis is made for convenience only and our result will not change if it is replaced by any other boundary condition at the "loose ends", in particular, by Dirichlet or θ-periodic ones (cf. Subsection 9.A).

quadratic forms differ from (2.1) by an extra term. In general there are many admissible boundary conditions; a graph vertex joining n edges gives rise to a family with n^2 real parameters. An example is represented by the so-called δ coupling for which the corresponding domain consists of all functions $u \in C(M_0)$ which are twice weakly differentiable on each edge, and (2.3) is replaced by

$$\sum_{j,\ e_j \text{ meets } v_k} p_j(v_k) u_j'(v_k) = \kappa u(v_k) \tag{2.4}$$

with a fixed $\kappa \in \mathbb{R}$, where $u(v_k)$ is the common value of all the $u_j(v_k)$ at the vertex. One can ask naturally whether such graph Hamiltonians can be obtained from a family of graph-shaped manifolds. In Section 7 we will discuss a particular case of the limiting procedure leading to the spectrum which — although it does *not* correspond to a graph operator with the generalized boundary condition described above — is at least *similar* to that with a δ coupling. The difference is that in the boundary conditions (2.4) the coupling constant κ is replaced by a quantity dependent on the spectral parameter, the corresponding operator being defined not on $L^2(M_0)$ but on a slightly enlarged Hilbert space — cf. (7.1)–(7.4).

In Section 6 we obtain another limit operator due to a different limiting procedure. This operator is again no graph operator with boundary conditions as above, but decouples and the graph part corresponds to a fully decoupled operator with Dirichlet boundary conditons at each vertex.

The spectrum of Δ_{M_0} is purely discrete. We denote the corresponding eigenvalues by $\lambda_k(\Delta_{M_0}) = \lambda_k(M_0)$, $k \in \mathbb{N}$, written in the ascending order and repeated according to multiplicity. With this eigenvalue ordering, we can employ the *min-max principle* (in the present form it can be found, e.g., in [Dav96]): the k-th eigenvalue of Δ_{M_0} is expressed as

$$\lambda_k(M_0) = \inf_{L_k} \sup_{u \in L_k \setminus \{0\}} \frac{\|q_0(u)\|^2}{\|u\|^2} \tag{2.5}$$

where the infimum is taken over all k-dimensional subspaces L_k of $\mathcal{H}^1(M_0)$.

2.B. **Comparison of eigenvalues.** Let us now formulate a simple consequence of the min-max principle which will be crucial for the proof of our main results. Suppose that $\mathcal{H}$, $\mathcal{H}'$ are two separable Hilbert spaces with the norms $\|\cdot\|$ and $\|\cdot\|'$. We need to compare eigenvalues λ_k and λ_k' of non-negative operators Q and Q' with purely discrete spectra defined via quadratic forms q and q' on $\mathcal{D} \subset \mathcal{H}$ and $\mathcal{D}' \subset \mathcal{H}$. We set $\|u\|_{Q,n}^2 := \|u\|^2 + \|Q^{n/2}u\|^2$.

Lemma 2.1. *Suppose that $\Phi\colon \mathcal{D} \longrightarrow \mathcal{D}'$ is a linear map such that there exist constants $n_1, n_2 \geq 0$ and $\delta_1, \delta_2 \geq 0$ such that*

$$\|u\|^2 \leq \|\Phi u\|'^2 + \delta_1 \|u\|_{Q,n_1}^2 \tag{2.6}$$

$$q(u) \geq q'(\Phi u) - \delta_2 \|u\|_{Q,n_2}^2 \tag{2.7}$$

for all $u \in \mathcal{D}$ and that $\mathcal{D} \subset \operatorname{dom} Q^{\max\{n_1,n_2\}/2}$. Then to each k there is a positive function η_k given by (2.11) *satisfying $\eta_k := \eta(\lambda_k, \delta_1, \delta_2) \to 0$ as $\delta_1, \delta_2 \to 0$, such that*

$$\lambda_k \geq \lambda'_k - \eta_k.$$

Proof. Let $\varphi_1, \dots, \varphi_k$ be an orthonormal system of eigenvectors corresponding to the eigenvalues $\lambda_1, \dots, \lambda_k$. For u in the linear span E_k of $\varphi_1, \dots, \varphi_k$, we have

$$\|u\|^2_{Q,n} \leq (1 + \lambda_k^n)\|u\|^2 \tag{2.8}$$

and

$$\begin{aligned}\frac{q'(\Phi u)}{\|\Phi u\|'^2} - \frac{q(u)}{\|u\|^2} &= \frac{q(u)}{\|u\|^2}\frac{\|u\|^2 - \|\Phi u\|'^2}{\|\Phi u\|'^2} + \frac{q'(\Phi u) - q(u)}{\|\Phi u\|'^2} \\ &\leq \left(\frac{q(u)}{\|u\|^2}\delta_1\|u\|^2_{Q,n_1} + \delta_2\|u\|^2_{Q,n_2}\right)\frac{1}{\|\Phi u\|'^2} \\ &\leq (\lambda_k(1 + \lambda_k^{n_1})\delta_1 + (1 + \lambda_k^{n_2})\delta_2)\frac{\|u\|^2}{\|\Phi u\|'^2} \end{aligned} \tag{2.9}$$

where we have used (2.6) and (2.7) to get the first inequality and (2.8) to get the second one. From relation (2.6) we follow

$$(1 - (1 + \lambda_k^{n_1})\delta_1)\|u\|^2 \leq \|\Phi u\|'^2 \tag{2.10}$$

and thus we can estimate the r.h.s. of (2.9) by

$$\eta_k := \eta(\lambda_k, \delta_1, \delta_2) := \frac{\lambda_k(1 + \lambda_k^{n_1})\delta_1 + (1 + \lambda_k^{n_2})\delta_2}{1 - (1 + \lambda_k^{n_1})\delta_1} \tag{2.11}$$

provided $0 \leq \delta_1 < 1/(1 + \lambda_k^{n_1})$. From (2.10) we also conclude that $\|u\| = 0$ holds if $\|\Phi u\|' = 0$, i.e., that $\Phi(E_k)$ is k-dimensional. From the min-max principle applied to the quadratic form q' we obtain

$$\lambda'_k \leq \sup_{u \in E_k \setminus \{0\}} \frac{q'(\Phi u)}{\|\Phi u\|'^2} \leq \sup_{u \in E_k \setminus \{0\}} \frac{q(u)}{\|u\|^2} + \eta_k = \lambda_k + \eta_k$$

which is the desired result. □

3. Graph-like manifolds

3.A. **Laplacian on a manifold.** Throughout this paper we study manifolds of dimension $d \geq 2$. For a Riemannian manifold X (compact or not) without boundary we denote by $L_2(X)$ the usual L_2-space of square integrable functions on X with respect to the volume measure dX on X. In a chart, the volume measure has the density $(\det G)^{1/2}$ with respect to the Lebesgue measure, where $\det G$ is the determinant of the metric tensor $G := (g_{ij})$ in this chart. The norm

of $L_2(X)$ will be denoted by $\|\cdot\|_X$. For $u \in C_c^\infty(X)$, the space of compactly supported smooth functions, we set

$$\check{q}_X(u) := \|du\|_X^2 = \int_X |du|^2 dX.$$

Here the 1-form du denotes the exterior derivative of u whose squared norm in coordinates is given by

$$|du|^2 = \sum_{i,j} g^{ij} \partial_i u \, \partial_j \overline{u} = G^{-1} \nabla u \cdot \nabla \overline{u}$$

where (g^{ij}) is the component representation of the inverse matrix G^{-1}.

We denote the closure of the non-negative quadratic form $\check{q}_X$ by q_X. Note that the domain $\operatorname{dom} q_X$ of the closed quadratic form q_X consists of functions in $L_2(X)$ such that the weak derivative du is also square integrable, i.e., $q_X(u) < \infty$.

We define the *Laplacian* Δ_X (for a manifold without boundary) as the unique self-adjoint and non-negative operator associated with the closed quadratic form q_X, i.e., the operator and the quadratic form are related by

$$q_X(u) = \langle \Delta_X u, u \rangle$$

for $u \in C_c^\infty(X)$ (see again [K66, Chapter VI], [RS-1] or [Dav96]). Thus the Laplacian action on smooth functions u is given in a fixed chart by

$$\Delta_X u = -(\det G)^{-1/2} \sum_{i,j} \partial_i \big((\det G)^{1/2} g^{ij} \, \partial_j u \big). \tag{3.1}$$

If X is a compact manifold with piecewise smooth boundary $\partial X \neq \emptyset$ we can define the Laplacian with Neumann boundary condition via the closure q_X of the quadratic form $\check{q}_X$ defined on $C^\infty(X)$, the space of smooth functions with derivatives continuous up to the boundary of X. Note that the usual conditions on the normal derivative occurs only in the *operator* domain via the Gauss-Green formula. In a similar way other boundary conditions at ∂X may be introduced. The spectrum of Δ_X (with any boundary condition if $\partial X \neq \emptyset$) is purely discrete as long as X is compact and the boundary conditions are local. We denote the corresponding eigenvalues by $\lambda_k(\Delta_X) = \lambda_k(X)$, $k \in \mathbb{N}$, written in increasing order and repeated according to multiplicity.

3.B. **General estimates on manifolds.** We will employ (partial) averaging processes on edge and vertex neighbourhoods which correspond to projection onto the lowest (transverse) mode. We start with such a general Poincaré-type estimate:

Lemma 3.1. *Let X be a connected, compact manifold with smooth boundary ∂X. For $u \in \mathcal{H}^1(X)$ define the constant function $u_0(x) := \frac{1}{\operatorname{vol} X} \int_X u \, dX$. Then we have $\|u_0\|_X^2 \leq \|u\|_X^2$,*

$$\|u - u_0\|_X^2 \leq \frac{1}{\lambda_2^{\mathrm{N}}(X)} \|du\|_X^2 \qquad \text{and} \qquad \|u\|_X^2 - \|u_0\|_X^2 \leq \frac{1}{\delta \lambda_2^{\mathrm{N}}(X)} + \delta \|u\|_X^2$$

for $\delta > 0$.

Proof. The first inequality follows directly from Cauchy-Schwarz. For the second one, note that $u - u_0$ is orthogonal to the first eigenfunction of the Neumann Laplacian. By the min-max principle we obtain

$$\lambda_2^{\mathrm{N}}(X)\|u - u_0\|_X^2 \leq \|d(u - u_0)\|_X^2 = \|du\|_X^2,$$

because u_0 is constant on X. Since X is connected, we have $\lambda_2^{\mathrm{N}}(X) > 0$. The last inequality follows from

$$|\|u\|^2 - \|u_0\|^2| \leq 2\|u - u_0\|\,\|u\| \leq \frac{1}{\delta}\|u - u_0\|^2 + \delta\|u\|^2 \tag{3.2}$$

for all $\delta > 0$. □

Next, we often need the following continuity of the map which restricts a function on X to the boundary ∂X. To this aim we use standard Sobolev embedding theorems:

Lemma 3.2. *There exists a constant $c_1 > 0$ depending only on X and the metric g such that*

$$\|u\restriction_{\partial X}\|_{\partial X}^2 \leq c_1(\|u\|_X^2 + \|du\|_X^2)$$

for all $u \in \mathcal{H}^1(U)$.

Proof. See e.g. [Tay96, Ch. 4, Prop. 4.5]. An alternative proof similar to the proof of Lemma 6.7 exists, and follows easily from (6.20) together with a cut-off function. □

3.C. **Definition of the graph-like manifold.** For each $0 < \varepsilon \leq 1$ we associate with the graph M_0 a compact and connected Riemannian manifold M_ε of dimen-



FIGURE 1. The associated edge and vertex neighbourhoods with $F = \mathbb{S}^1$, i.e., $U_{\varepsilon,j}$ and $V_{\varepsilon,k}$ are 2-dimensional manifolds with boundary.

sion $d \geq 2$ equipped with a metric g_ε to be specified below. We suppose that M_ε is the union of compact subsets $U_{\varepsilon,j}$ and $V_{\varepsilon,k}$ such that the interiors of $U_{\varepsilon,j}$ and $V_{\varepsilon,k}$ are mutually disjoint for all possible combinations of $j \in J$ and $k \in K$. We think of $U_{\varepsilon,j}$ as the thickened edge e_j and of $V_{\varepsilon,k}$ as the thickened vertex v_k (see Figures 1 and 2). Note that the second picture describes the situation only rougly, since it assumes that M_ε is embedded in $\mathbb{R}^\nu$. More correctly, we should think of M_ε as an abstract manifold obtained by identifying the appropriate boundary parts of $U_{\varepsilon,j}$ and $V_{\varepsilon,k}$ via the connection rules of the graph M_0. This manifold



FIGURE 2. On the left, we have the graph M_0, on the right, the associated graph-like manifold (in this case, $F = \mathbb{S}^1$ and M_ε is a 2-dimensional manifold).

need not to be embedded, but the situation when M_ε is a submanifold of $\mathbb{R}^\nu$ ($\nu \geq d$) can be viewed also in this abstract context (see Remark 4.1).

As a matter of convenience we assume that $U_{\varepsilon,j}$ and $V_{\varepsilon,k}$ are independent of ε as manifolds, i.e., only their metric g_ε depend on ε. This can be achieved in the following way: for the edge regions we assume that $U_{\varepsilon,j}$ is diffeomorphic to $I_j \times F$ for all $0 < \varepsilon \leq 1$ where F denotes a compact and connected manifold (with or without a boundary) of dimension $m := d - 1$. For the vertex regions we assume that the manifold $V_{\varepsilon,k}$ is diffeomorphic to a ε-independent manifold V_k for $0 < \varepsilon \leq 1$. Pulling back the metrics to the diffeomorphic manifold we may assume that the underlying differentiable manifold is independent of ε. Therefore, $U_{\varepsilon,j} = U_j = I_j \times F$ and $V_{\varepsilon,k} = V_k$ with an ε-depending metric g_ε.

For further purposes, we need a decomposition of $e_j \cong I_j$ into two halves. We reverse the orientation of one such half so that each half is directed away from its adjacent vertex and collect all halves I_{jk} ending at the vertex v_k, i.e., $j \in J_k$, where

$$J_k := \{\, j \in J \,|\, e_j \text{ meets } v_k \,\}^4 \tag{3.3}$$

We denote $U_{jk} := I_{jk} \times F$ (and similar notation with subscript ε).

[4]For each loop e_j at v_k, i.e., each edge beginning *and* ending at v_k, we need to replace the label j by two distinct labels j_1, j_2 belonging to J_k in order to collect *both* halfs of the edge.

For further references, we denote the midpoint of the edge $e_j \cong I_j$ by x_j^* and the endpoint of I_j corresponding to the edge v_k by x_{jk}^0, e.g., $I_{jk} = [x_j^*, x_{jk}^0]$.

3.D. **Notation.** In the sequel, we are going to suppress the edge and vertex subscripts j and k unless a misunderstanding may occur. Similarly we set, e.g., $U := U_1$, in other words we omit the subscript ε if we only mean the underlying ε-independent manifold with metric g_1, i.e., if we fix $\varepsilon = 1$.

3.E. **Motivation for the different limit operators.** Let us shortly motivate why the limit operator of Δ_{M_ε} as $\varepsilon \to 0$ should depend on the volume decay of the vertex neighbourhoods $V_{\varepsilon,k}$ in comparison with $\operatorname{vol}_{d-1} \partial V_{\varepsilon,k}$ (or $\operatorname{vol}_d U_{\varepsilon,j}$, which is of the same order when $\varepsilon \to 0$ as we will see in Section 4). For simplicity, we assume that the radius of the transversal direction on the edge $U_{\varepsilon,j}$ is ε (i.e., $p_j \equiv 1$). The assumptions on the edge neighbourhoods will be specified in the next section. We stress that our aim in this subsection is to present a heuristic idea, not a proof (for a suitable reasoning cf. [RueS53] or [Ku02]).

Suppose $\varphi = \varphi_\varepsilon$ is an eigenfunction of Δ_{M_ε} w.r.t the eigenvalue $\lambda = \lambda_\varepsilon$. By the Gauss-Green formula, we have at the vertex $V_\varepsilon = V_{\varepsilon,k}$

$$\lambda \int_{V_\varepsilon} \varphi\, \overline{u}\, dV_\varepsilon = \int_{V_\varepsilon} \langle d\varphi, du\rangle\, dV_\varepsilon + \int_{\partial V_\varepsilon} \partial_{\mathrm{n}} \varphi\, \overline{u}\, d\partial V_\varepsilon \tag{3.4}$$

for all $u \in \mathcal{H}^1(M_\varepsilon)$. Assume that $\lambda_\varepsilon \to \lambda_0$ and $\varphi_\varepsilon \to \varphi_0 = (\varphi_{0,j})_j$.

If the vertex volume $\operatorname{vol}_d V_\varepsilon$ decays faster then the boundary area $\operatorname{vol}_{d-1} \partial V_\varepsilon$ only the boundary integral over ∂V_ε survives in the limit $\varepsilon \to 0$ and leads to

$$0 = \sum_{j \in J_k} \varphi'_{0,j}(v_k)$$

which is exactly the Kirchhoff boundary condition mentioned above in (2.3). This *fast decaying* vertex volume case will be treated in Section 5.

If the vertex volume decays slower than $\operatorname{vol}_{d-1} \partial V_\varepsilon$, the integrals over V_ε are dominant. In this case, $\operatorname{vol} V_{\varepsilon,k} \gg \operatorname{vol} U_{\varepsilon,j}$ and only slowly varying eigenfunctions on $V_{\varepsilon,k}$ lead to bounded eigenvalues $\lambda = \lambda_\varepsilon$. Since $\operatorname{vol} V_{\varepsilon,k} \gg \operatorname{vol} U_{\varepsilon,j}$, normalized eigenfunctions are nearly vanishing on $V_{\varepsilon,k}$ viewed from the scale on $U_{\varepsilon,j}$. This roughly explains, why we end up in a decoupled operator with Dirichlet boundary conditions on M_0 plus extra zero eigenmodes at the vertices (the zero eigenmodes also survive the limit $\varepsilon \to 0$). This *slow decaying* vertex volume case will be shortly be discussed in Section 6.

In the borderline case when $\operatorname{vol}_d V_\varepsilon \approx \operatorname{vol}_{d-1} \partial V_\varepsilon$, we also expect the eigenfunctions to vary slowly on $V_{\varepsilon,k}$ (since $\operatorname{vol}_d V_{\varepsilon,k} \to 0$), so the integral over $\langle d\varphi, du\rangle$ should tend to 0, and in the limit

$$\lambda_0 \varphi_0(v_k) = \sum_{j \in J_k} \varphi'_{0,j}(v_k).$$

This *borderline case* will be treated in Section 7.

If $\operatorname{vol} V_{\varepsilon,k}$ does not tend to 0, i.e., when $V_{\varepsilon,k}$ tends to a compact d-dimensional manifold $V_{0,k}$ without boundary (and *not* to a point as in the cases above), we still expect a decoupled operator with Dirichlet boundary conditions on the edges by the same arguments as in the slow decaying case. In addition, not only the lowest eigenmode of $V_{\varepsilon,k}$ but all eigenmodes survive, i.e., the limit operator should consist of the direct sum of all Dirichlet Laplacians on the edges plus the Laplacians on $V_{0,k}$, $k \in K$. This *non-decaying* vertex volume case will be treated in Section 8.

It needs an extra effort to prove rigorously the conclusions of the above reasoning; recall that we have assumed e.g. that $\lambda_\varepsilon \to \lambda_0$ (which we want to show in this paper) and $\|\varphi_\varepsilon\|_\infty, \|d\varphi_\varepsilon\|_\infty \le c$ which is in general not true for normalized (L_2-)eigenfunctions since $\operatorname{vol} M_\varepsilon \to 0$ as $\varepsilon \to 0$.

4. Edge neighbourhoods

4.A. **Definition of the thickened edges.** Suppose that $U = I \times F$ with metric g_ε, where I corresponds to some (part of an) edge and F denotes (as before) a compact and connected Riemannian manifold of dimension $m = d-1$ with metric h. Without loss of generality we may assume that $\operatorname{vol} F = 1$. We define another metric $\tilde{g}_\varepsilon$ on U_ε by

$$\tilde{g}_\varepsilon := dx^2 + \varepsilon^2 r_j^2(x)\, h(y), \qquad (x,y) \in U_j = I_j \times F \tag{4.1}$$

where

$$r_j(x) := (p_j(x))^{1/m} \tag{4.2}$$

defines a smooth function (specifying the radius of the fibre $\{x\} \times F$ at the point x), where p_j is the density function on the edge e_j introduced in Section 2.

We denote by G_ε and $\tilde{G}_\varepsilon$ the $d \times d$-matrices associated to the metrics g_ε and $\tilde{g}_\varepsilon$ with respect to the coordinates (x,y) (here y stands for suitable coordinates on F) and assume that the two metrics coincide up to an error term as $\varepsilon \to 0$, more specifically

$$G_\varepsilon = \tilde{G}_\varepsilon + \begin{pmatrix} o(1) & o(\varepsilon) \\ o(\varepsilon) & o(\varepsilon^2) \end{pmatrix} = \begin{pmatrix} 1+o(1) & o(\varepsilon) \\ o(\varepsilon) & \varepsilon^2 r_j\, H + o(\varepsilon^2) \end{pmatrix}, \tag{4.3}$$

i.e.,

$$g_{\varepsilon,xx} = 1 + o(1), \quad g_{\varepsilon,xy_\alpha} = o(\varepsilon), \quad g_{\varepsilon,y_\alpha y_\beta} = \varepsilon^2 r_j^2(x)\, h_{\alpha\beta}(y) + o(\varepsilon^2).$$

uniformly on U. To summarize, we assume that the metric g_ε is equal to the product metric $\tilde{g}_\varepsilon$ up to error terms.

Remark 4.1. This is a central assumption in our construction which describes how in fact the family of manifolds shrinks to the graph M_0. One of the reasons why we introduce a pair of metrics is the following. While our construction uses intrinsic metric properties of the manifolds only, we want it to be applicable to manifolds embedded into some Euclidean space (e.g. in $\mathbb{R}^{d+1}$ if $F = \mathbb{S}^1$). This is impossible if the cylindrical sleeves have the same length as the underlying graph

edges, but it can be achieved with the length modified by a factor of $o(1)$. It will be one of our aims to show that within the prescribed error margin such a "practically important" metric yields the same result as the product metric which is easier to handle.

In the next two examples we illustrate that a suitable smooth ε-neighbourhood of an embedded graph in $\mathbb{R}^\nu$ ($\nu \geq d$) as well as an embedded *curved* edge is covered by the above abtract setting of a pure product metric and an "almost" product metric.

Example 4.2. **Embedded graphs.** If we set $\nu = d = 2$ and $F = [-1, 1]$ (or more generally $\nu = d \geq 2$, $F := \{\, x \in \mathbb{R}^m \,|\, |x| \leq 1 \,\}$) and if M_ε is a suitable ε-neighbourhood of M_0, we recover the situation treated in [KuZ01].

Note that the error term $o(1)$ in the first component in (4.3) allows us to start with a graph M_0 embedded in some Euclidean space $\mathbb{R}^\nu$ as we have already mentioned in Remark 4.1. The edge neighbourhoods $U_{\varepsilon,j}$ of the embedded manifold M_ε have a length *smaller* than the length of the corresponding edge, with the error term $o(1)$. By a simple transformation of the variable x we return to our setting with a fixed length of $U_j = I_j \times F$.

Example 4.3. **Curved edges and variable transversal radius.** Suppose U_ε is the ε-neighbourhood of a smooth curve $\vec{\gamma} = \vec{\gamma}_j \colon I_j \longrightarrow \mathbb{R}^d$ parametrized by arc-length. If, e.g., $\nu = d = 2$ and $F = [-1, 1]$ a chart is given by

$$\Psi \colon I_j \times [-1, 1] \longrightarrow U_{\varepsilon,j}, \qquad (x, y) \mapsto \vec{\gamma}(x) + \varepsilon r_j(x) y \, \vec{n}(x),$$

i.e., we thicken the curve $\vec{\gamma}$ in its normal direction $\vec{n}(x)$ at the point $\vec{\gamma}(x)$ by the factor $\varepsilon r(x) = \varepsilon r_j(x)$. The corresponding metric in (x, y)-coordinates is given by

$$G_\varepsilon = \begin{pmatrix} (1 + \varepsilon\kappa y r)^2 + \varepsilon^2 y^2 \dot{r}^2 & \varepsilon^2 r \dot{r} y \\ \varepsilon^2 r \dot{r} y & \varepsilon^2 r^2 \end{pmatrix} = \begin{pmatrix} 1 + O(\varepsilon) & O(\varepsilon^2) \\ O(\varepsilon^2) & \varepsilon^2 r^2 \end{pmatrix}$$

where $\kappa := \dot{\gamma}_1 \ddot{\gamma}_2 - \dot{\gamma}_2 \ddot{\gamma}_2$ is the curvature of the generating curve $\vec{\gamma}$. Therefore, the error term $o(1)$ comes from the curvature of the embedded curve $\vec{\gamma}$ whereas the off-diagonal error terms come from the variable radius of the transversal direction (note that $\dot{r} = 0$ if $r(x)$ is constant). Furthermore, the metric density satisfies

$$\det G_\varepsilon^{\frac{1}{2}} = |1 + \varepsilon\kappa y r| \varepsilon r = (1 + O(\varepsilon)) \det \tilde{G}_\varepsilon^{\frac{1}{2}}.$$

We give a general treatment in the next lemma.

4.B. **Estimates on the thickened edges.** Following the philosophy explained in Remark 4.1, we start with pointwise estimates where we compare the product metric $\tilde{g}_\varepsilon$ with the original metric g_ε. Note that the assumption (4.3), while fully sufficient for our purposes, is optimal in a sense, i.e., that the following lemma ceases to be valid if we weaken its hypothesis even slightly.

Lemma 4.4. *Suppose that g_ε, $\tilde{g}_\varepsilon$ are given as in* (4.1) *and* (4.3)*, then*

$$(\det G_\varepsilon)^{\frac{1}{2}} = (1+o(1))(\det \tilde{G}_\varepsilon)^{\frac{1}{2}} \tag{4.4}$$

$$g_\varepsilon^{xx} := (G_\varepsilon^{-1})_{xx} = 1 + o(1) \tag{4.5}$$

$$|d_x u|^2 \leq (1+o(1))|du|^2_{g_\varepsilon} \tag{4.6}$$

$$|d_F u|^2_h \leq o(\varepsilon)|du|^2_{g_\varepsilon} \tag{4.7}$$

where d_x and d_F are the (exterior) derivative with respect to $x \in I$ and $y \in F$, respectively. All the estimates are uniform in (x,y) as $\varepsilon \to 0$.

Proof. The first equation follows from

$$\begin{aligned}\det(G_\varepsilon \tilde{G}_\varepsilon^{-1}) &= \det \begin{pmatrix} 1+o(1) & o(\varepsilon) \\ o(\varepsilon) & \varepsilon^2 H + o(\varepsilon^2) \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \varepsilon^{-2} H^{-1} \end{pmatrix} \\ &= \det \begin{pmatrix} 1+o(1) & o(\varepsilon^{-1}) \\ o(\varepsilon) & \mathbb{1} + o(1) \end{pmatrix} = 1 + o(1).\end{aligned}$$

For the second one, we consider the upper left component of

$$\begin{aligned} G_\varepsilon^{-1} - \tilde{G}_\varepsilon^{-1} &= -\tilde{G}_\varepsilon^{-1}(G_\varepsilon - \tilde{G}_\varepsilon)\tilde{G}_\varepsilon^{-1} + o(G_\varepsilon - \tilde{G}_\varepsilon) \\ &= \begin{pmatrix} 1 & 0 \\ 0 & O(\varepsilon^{-2}) \end{pmatrix} \begin{pmatrix} o(1) & o(\varepsilon) \\ o(\varepsilon) & o(\varepsilon^2) \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & O(\varepsilon^{-2}) \end{pmatrix} + o(1) \\ &= \begin{pmatrix} o(1) & o(\varepsilon^{-1}) \\ o(\varepsilon^{-1}) & o(\varepsilon^{-2}) \end{pmatrix}. \end{aligned}$$

Inequality (4.6) is equivalent to

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \leq (1+o(1)) G_\varepsilon^{-1}$$

in the sense of quadratic forms. This will be true if we show that

$$\begin{pmatrix} 1 & 0 \\ 0 & \delta\mathbb{1} \end{pmatrix} \leq (1+o(1)) G_\varepsilon^{-1}$$

for some $\delta > 0$, where $\mathbb{1}$ is the $m \times m$ unit matrix, which in turn means

$$(1+o(1)) \begin{pmatrix} 1 & 0 \\ 0 & \delta^{-1}\mathbb{1} \end{pmatrix} \geq G_\varepsilon.$$

However,

$$G_\varepsilon = \tilde{G}_\varepsilon + \begin{pmatrix} o(1) & o(\varepsilon) \\ o(\varepsilon) & o(\varepsilon^2) \end{pmatrix} = \begin{pmatrix} 1+o(1) & 0 \\ 0 & O(\varepsilon^2) \end{pmatrix} + \begin{pmatrix} 0 & o(\varepsilon) \\ o(\varepsilon) & 0 \end{pmatrix}$$

and the eigenvalues of the last matrix are of order $o(\varepsilon)$, so

$$G_\varepsilon \leq \begin{pmatrix} 1+o(1) & 0 \\ 0 & O(\varepsilon^2) \end{pmatrix} + o(\varepsilon)\mathbb{1} = \begin{pmatrix} 1+o(1) & 0 \\ 0 & o(\varepsilon) \end{pmatrix} \leq (1+o(1)) \begin{pmatrix} 1 & 0 \\ 0 & c\mathbb{1} \end{pmatrix}$$

for some constant $c > 0$, and therefore it is sufficient to choose $\delta < c^{-1}$. In the same way, inequality (4.7) is equivalent to

$$\begin{pmatrix} 0 & 0 \\ 0 & H^{-1} \end{pmatrix} \le o(\varepsilon) G_\varepsilon^{-1}.$$

We are looking for $\delta = \delta(\varepsilon)$ which tends to zero as $\varepsilon \to 0$ such that

$$\begin{pmatrix} \delta & 0 \\ 0 & H^{-1} \end{pmatrix} \le o(\varepsilon) G_\varepsilon^{-1}$$

which is equivalent to

$$o(\varepsilon) \begin{pmatrix} \delta^{-1} & 0 \\ 0 & H \end{pmatrix} \ge G_\varepsilon.$$

However, as before we have

$$G_\varepsilon \le \begin{pmatrix} 1+o(1) & 0 \\ 0 & O(\varepsilon^2)H \end{pmatrix} + o(\varepsilon)\mathbb{1} \le \begin{pmatrix} 1+o(1) & 0 \\ 0 & o(\varepsilon)H \end{pmatrix},$$

and consequently, $\delta(\varepsilon)^{-1} \ge (1+o(1))/o(\varepsilon)$, i.e., $\delta(\varepsilon) \le o(\varepsilon)/(1+o(1)) = o(\varepsilon)$. □

4.C. **Notation.** The tilde in a symbol refers always to the product metric $\tilde{g}_\varepsilon$. We denote, e.g., by $\tilde{U}_\varepsilon$ the manifold U_ε with metric $\tilde{g}_\varepsilon$ and (abusing the notation a little bit) employ the symbol U_ε for the manifold U_ε with the metric g_ε.

As a motivation for the above choice of the metrics, let us calculate the norm of $u \in L_2(\tilde{U}_\varepsilon)$ for a function u which is independent of the second argument $y \in F$, i.e. $u(x,y) = u(x)$. This yields

$$\begin{aligned} \|u\|^2_{\tilde{U}_\varepsilon} &= \int_{U_\varepsilon} |u|^2 d\tilde{U}_\varepsilon \\ &= \int_I \int_F |u(x)|^2 (\det \tilde{G}_\varepsilon)^{\frac{1}{2}}(x,y)\, dx\, dy \\ &= \varepsilon^m \int_I |u(x)|^2 r^m(x) dx \int_F (\det H)^{\frac{1}{2}}(y)\, dy \\ &= \varepsilon^m \|u\|^2_I \operatorname{vol}(F) = \varepsilon^m \|u\|^2_I. \end{aligned} \tag{4.8}$$

4.D. **Transversal averaging.** We will employ averaging processes on edge neighbourhoods $U_\varepsilon = U_{\varepsilon,j}$ which correspond to projection onto the lowest transverse mode:

$$Nu(x) = N_j u(x) := \int_F u(x,\cdot)\, dF \tag{4.9}$$

Note that $Nu(x)$ is well defined for $u \in \mathcal{H}^1(U_\varepsilon)$, and moreover,

$$\varepsilon^m \|Nu\|^2_I = \|Nu\|^2_{\tilde{U}_\varepsilon} \le \|u\|^2_{\tilde{U}_\varepsilon} = (1+o(1))\|u\|^2_{U_\varepsilon} \tag{4.10}$$

in view of eqs. (4.8), (4.4), and the Cauchy-Schwarz inequality.

In the following two lemmas we compare a function u and its derivative du with the normal averages Nu and d_xNu, respectively. Note that for the next lemma, (4.10) is not enough; we also need the opposite inequality:

Lemma 4.5. *For any $u \in \mathcal{H}^1(U_\varepsilon)$ we have*

$$\|u\|^2_{U_\varepsilon} - \|\varepsilon^{m/2} Nu\|^2_I \le o(\varepsilon^{1/2}) \left(\|u\|^2_{U_\varepsilon} + \|du\|^2_{U_\varepsilon}\right).$$

Proof. Applying Lemma 3.1 with $X = F$ we get

$$\|u(x,\cdot)\|^2_F - |Nu(x)|^2 \le \frac{1}{\delta \lambda_2^{\mathrm{N}}(F)} \|d_F u(x,\cdot)\|^2_F + \delta \|u(x,\cdot)\|^2_F. \tag{4.11}$$

Next we integrate over I_j and obtain

$$\|u\|^2_{\tilde{U}_\varepsilon} - \varepsilon^m \|Nu\|^2_{I_j} \le \frac{o(\varepsilon)}{\delta \lambda_2^{\mathrm{N}}(F)} \int_{\tilde{U}_{\varepsilon,j}} |du|^2_{g_\varepsilon} d\tilde{U}_{\varepsilon,j} + \delta \|u\|^2_{\tilde{U}_{\varepsilon,j}}$$

using estimate (4.7). We put $\delta := \sqrt{o(\varepsilon)}$ and apply (4.4) to obtain the result for the manifold $U_{\varepsilon,j}$. □

Lemma 4.6. *For any $u \in \mathcal{H}^1(U_\varepsilon)$ we have*

$$\|\varepsilon^{m/2} (Nu)'\|^2_I - \|du\|^2_{U_\varepsilon} \le o(1) \|du\|^2_{U_\varepsilon}.$$

Proof.

$$\begin{aligned} \|\varepsilon^{m/2} (Nu)'\|^2_I &= \|\varepsilon^{m/2} N(d_x u)\|^2_I \le (1 + o(1)) \|d_x u\|^2_{U_\varepsilon} \\ &= (1 + o(1)) \int_{U_\varepsilon} |d_x u|^2 \, dU_\varepsilon \le (1 + o(1)) \int_{U_\varepsilon} |du|^2_{g_\varepsilon} \, dU_\varepsilon = (1 + o(1)) \|du\|^2_{U_\varepsilon} \end{aligned}$$

holds in view of estimate (4.10) and (4.6). □

Next we need a pointwise estimate on the behaviour of Nu at the boundary.

Lemma 4.7. *We have*

$$|Nu(x^0)|^2 \le O(\varepsilon^{-m})(\|u\|^2_{U_\varepsilon} + \|du\|^2_{U_\varepsilon})$$

for all $u \in \mathcal{H}^1(U_\varepsilon)$, where $x^0 \in \partial I$.

Proof. One estimates

$$|Nu(x^0)|^2 \le \int_F |u(x^0, y)|^2 \, dF(y) \le c_1(\|u\|^2_U + \|d_x u\|^2_U) \le O(\varepsilon^{-m})(\|u\|^2_{U_\varepsilon} + \|du\|^2_{U_\varepsilon})$$

by Lemma 3.2 with $X = U$, (4.6) and (4.4). □

5. Fast decaying vertex volume

5.A. **Definition of the thickened vertices.** Remember that $V_{\varepsilon,k} = V_k$ as manifold, whereas g_ε denotes the ε-depending metric on $V_{\varepsilon,k}$. Let $g := g_1$, then we assume that

$$c_-\varepsilon^2 g \le g_\varepsilon \le c_+\varepsilon^{2\alpha} g \tag{5.1}$$

in the sense that there are constants $c_-, c_+ > 0$ such that

$$c_-\varepsilon^2 g(x)(v,v) \le g_\varepsilon(x)(v,v) \le c_+\varepsilon^{2\alpha} g(x)(v,v)$$

for all $v \in T_xV_k$ and all $x \in V_k$. The number α in the exponent is assumed to satisfy the inequalities

$$\frac{d-1}{d} < \alpha \le 1\,; \tag{5.2}$$

notice that $\alpha \le 1$ is needed for (5.1) to make sense with $0 < \varepsilon \le 1$. Thus the edge and vertex parts of the manifold need not shrink at the same rate but the vertex shrinking should not be too slower than that of the edges. This hypothesis expressed by (5.1) plays a central role here; other shrinking regimes will be discussed in the following sections.

Note that the manifold $V_{\varepsilon,k}$ shrinks at most as ε (in each direction) by the lower bound in (5.1). This ensures that a global *smooth* metric g_ε exists on M_ε with the requirements on $U_{\varepsilon,j}$ and $V_{\varepsilon,k}$. Therefore, we do not need an intermediate part (called *bottle neck*) between the edge and vertex neighbourhoods interpolating between the different scalings as in Sections 6 and 7 (see also Remark 5.11).

We easily obtain the following global estimates:

Lemma 5.1. *There are $c_1^\pm, c_2^\pm > 0$ such that*

$$c_1^-\varepsilon^d\|u\|_V^2 \le \|u\|_{V_\varepsilon}^2 \le c_1^+\varepsilon^{\alpha d}\|u\|_V^2 \tag{5.3}$$

$$c_2^-\varepsilon^{d-2\alpha}\|du\|_V^2 \le \|du\|_{V_\varepsilon}^2 \le c_2^+\varepsilon^{\alpha d-2}\|du\|_V^2 \tag{5.4}$$

for all $u \in \mathcal{H}^1(V_\varepsilon) = \mathcal{H}^1(V)$.

Proof. Using assumption (5.1) we obtain

$$c_-^{d/2}\varepsilon^d(\det G)^{1/2} \le (\det G_\varepsilon)^{1/2} \le c_+^{d/2}\varepsilon^{\alpha d}(\det G)^{1/2}$$

and

$$c_+^{-1}\varepsilon^{-2\alpha}G^{-1} \le G_\varepsilon^{-1} \le c_-^{-1}\varepsilon^{-2}G^{-1}$$

which implies the result with $c_1^\pm := c_\pm^{d/2}$ and $c_2^\pm := c_\pm^{d/2}c_\mp^{-1}$. □

5.B. **Convergence of the spectra.** The limit operator will concentrate only on the edge part in this case, therefore we define

$$\mathcal{H}_0 := L_2(M_0), \qquad \mathcal{D}_0 := \mathcal{H}^1(M_0), \qquad q_0(u) := \|u'\|_{M_0}^2 = \sum_j \|u_j'\|_{I_j}^2, \tag{5.5}$$

i.e., the limit operator Q_0 is Δ_{M_0} (see Def. (2.2)). With the above preliminaries we can finally formulate the main result of this section:

Theorem 5.2. *Under the stated assumptions* $\lambda_k(M_\varepsilon) \to \lambda_k(M_0)$ *as* $\varepsilon \to 0$.

Recall that the eigenvalues $\lambda_k(M_\varepsilon)$ are by assumption ordered in the ascending order, multiplicity taken into account, so the label of a particular eigenvalue curve may change as ε moves. The spectrum of the manifold is in general richer than that of the graph and a part of the eigenvalues escapes to $+\infty$ as $\varepsilon \to 0$; the proof presented below shows that this happens, roughly speaking, for all states with the transverse part of the eigenfunction orthogonal to the ground state.

Our aim is to find a two sided estimate on each eigenvalue $\lambda_k(M_\varepsilon)$ by means of $\lambda_k(M_0)$ with an error which is $o(1)$ w.r.t. the parameter ε.

5.C. **An upper bound.** The mentioned upper eigenvalue estimate now reads as follows:

Theorem 5.3. $\lambda_k(M_\varepsilon) \le \lambda_k(M_0) + o(1)$ *holds as* $\varepsilon \to 0$.

To prove it, we define the transition operator by

$$\Phi_\varepsilon u(z) := \begin{cases} \varepsilon^{-m/2} u(v_k) & \text{if } z \in V_k, \\ \varepsilon^{-m/2} u_j(x) & \text{if } z = (x,y) \in U_j \end{cases} \tag{5.6}$$

for any $u \in \mathcal{H}^1(M_0)$. Theorem 5.3 is then implied by Lemma 2.1 in combination with the following result.

Lemma 5.4. *We have* $\Phi_\varepsilon u \in \mathcal{H}^1(M_\varepsilon)$*, i.e.,* Φ_ε *maps the quadratic form domain of the Laplacian on the graph into the quadratic form domain of the Laplacian on the manifold. Furthermore, for* $u \in \mathcal{H}^1(M_0)$ *we have*

$$\|u\|^2_{M_0} - \|\Phi_\varepsilon u\|^2_{M_\varepsilon} \le o(1)\, \|u\|^2_{M_0} \tag{5.7}$$

$$\|d\,\Phi_\varepsilon u\|^2_{M_\varepsilon} - q_0(u) = o(1)\, q_0(u). \tag{5.8}$$

Proof. The first assertion is true since $\Phi_\varepsilon u$ is constant on each thickened vertex $V_{\varepsilon,k}$ and continuous on $\partial V_{\varepsilon,k}$. Clearly, $\Phi_\varepsilon u$ is weakly differentiable on each thickened edge $U_{\varepsilon,j}$. Moreover, we have

$$\begin{aligned} \|u\|^2_{M_0} - \|\Phi_\varepsilon u\|^2_{M_\varepsilon} &\le \sum_{j\in J} (\|u\|^2_{I_j} - \|\Phi_\varepsilon u\|^2_{U_{\varepsilon,j}}) \\ &= \sum_{j\in J} (\|u\|^2_{I_j} - (1+o(1))\|\Phi_\varepsilon u\|^2_{\tilde U_{\varepsilon,j}}) = o(1) \sum_{j\in J} \|u\|^2_{I_j} = o(1)\|u\|^2_{M_0} \end{aligned}$$

where we have neglected the contribution to the norm of $\Phi_\varepsilon u$ from the vertex parts of M_ε and employed eqs. (4.4) and (4.8). The second relation follows from

$$\begin{aligned} \|d\,\Phi_\varepsilon u\|^2_{M_\varepsilon} - q_0(u) &= \sum_{j\in J} ((1+o(1))\|g_\varepsilon^{xx} d_x \Phi_\varepsilon u\|^2_{\tilde U_{\varepsilon,j}} - \|u'\|^2_{I_j}) \\ &= \sum_{j\in J} ((1+o(1))\|u'\|^2_{I_j} - \|u'\|^2_{I_j}) = o(1)\, q_0(u) \end{aligned}$$

in the same way as above and with (4.5); recall that $\Phi_\varepsilon u$ is constant on $V_{\varepsilon,k}$ and independent of $y \in F$ on $U_{\varepsilon,j}$. □

Note that to get the upper bound we did not use any estimate for the metric on the vertex neighbourhoods $V_{\varepsilon,k}$.

5.D. **A lower bound.** The opposite estimate is more difficult. Here, we will also employ averaging processes on the vertex neigbourhoods $V_{\varepsilon,k}$ which correspond to projection onto the lowest (constant) mode:

$$Cu = C_k u := \frac{1}{\operatorname{vol} V_k} \int_{V_k} u \, dV_k. \tag{5.9}$$

Recall that $V = V_k$ denotes the manifold V_k with the metric $g = g_1$ (see Remark 5.7 for the reason why we use V_k instead of $V_{\varepsilon,k}$).

Lemma 5.5. *The inequality*

$$|C_k u - N_j u(x^0)|^2 \le O(\varepsilon^{2\alpha-d}) \|du\|^2_{V_{\varepsilon,k}}$$

holds for all $u \in \mathcal{H}^1(V_{\varepsilon,k})$ where the point $x^0 = x^0_{jk} \in \partial I_j$ corresponds to the vertex v_k.

Proof.

$$\begin{aligned} |C_k u - N_j u(x^0)|^2 &\le \int_F |C_k u - u(x^0, y)|^2 \, dF(y) \\ &\le c_1 \left(\|C_k u - u\|^2_{V_k} + \|du\|^2_{V_k} \right) \\ &\le c_1 \left(\frac{1}{\lambda^{\mathrm{N}}_2(V_k)} + 1 \right) \|du\|^2_{V_k} \\ &\le O(\varepsilon^{2\alpha-d}) \|du\|^2_{V_{\varepsilon,k}} \end{aligned}$$

holds by Lemma 3.2 and Lemma 3.1 with $X = V_k$ and metric $g = g_1$, and Lemma 5.1. □

Lemma 5.6. *We have*

$$\|u - Cu\|^2_{V_\varepsilon} \le O(\varepsilon^\beta) \|du\|^2_{V_\varepsilon}$$

for all $u \in \mathcal{H}^1(V_\varepsilon)$, where $\beta := (2+d)\alpha - d$.

Proof. Using again Lemmas 3.1 and 5.1 we infer

$$\|u - Cu\|^2_{V_\varepsilon} \le c_1^+ \varepsilon^{\alpha d} \|u - Cu\|^2_V \le c_1^+ \varepsilon^{\alpha d} \frac{1}{\lambda^{\mathrm{N}}_2(V)} \|du\|^2_V \le O(\varepsilon^{\alpha d - d + 2\alpha}) \|du\|^2_{V_\varepsilon}.$$

□

Notice that $\beta > 0$ is equivalent to $\alpha > d/(d+2)$ and the last inequality is satisfied due to (5.2) and the fact that $d \ge 2$ holds by assumption.

Remark 5.7. For Lemma 5.6, the "natural" averaging $C_\varepsilon u := \int_{V_\varepsilon} u \, dV_\varepsilon$ would yield the same result whereas Lemma 5.5 leads to the estimate $O(\varepsilon^{\beta-d})$ which is worse since $2\alpha > \beta$.

We conclude that in the fast decaying case the edge neighbourhoods lead to no spectral contribution in the limit $\varepsilon \to 0$:

Corollary 5.8. *The inequality*

$$\|u\|^2_{V_\varepsilon} \le O(\varepsilon^{\alpha d - m})(\|u\|^2_{U_\varepsilon \cup V_\varepsilon} + \|du\|^2_{U_\varepsilon \cup V_\varepsilon})$$

holds true for all $u \in \mathcal{H}^1(U_\varepsilon \cup V_\varepsilon)$.

Proof. We start from the telescopic estimate

$$\begin{aligned}\|u\|_{V_\varepsilon} &\le \|u - Cu\|_{V_\varepsilon} + \|Cu - Nu(x^0)\|_{V_\varepsilon} + \|Nu(x^0)\|_{V_\varepsilon} \\ &\le O(\varepsilon^{\beta/2})\|du\|_{V_\varepsilon} + (\operatorname{vol} V_\varepsilon)^{1/2} \left(O(\varepsilon^{(2\alpha-d)/2})\|du\|^2_{V_\varepsilon} + O(\varepsilon^{-m})(\|u\|^2_{U_\varepsilon} + \|du\|^2_{U_\varepsilon}\right)^{1/2} \\ &= O(\varepsilon^{(\alpha d - m)/2})(\|u\|^2_{U_\varepsilon \cup V_\varepsilon} + \|du\|^2_{U_\varepsilon \cup V_\varepsilon})^{1/2}\end{aligned}$$

where we have used Lemmas 5.6, 5.5, and 4.7, and furthermore the inequality (5.3) to obtain $\operatorname{vol} V_\varepsilon = O(\varepsilon^{\alpha d})$. Finally, note that $\beta = (d+2)\alpha - d > \alpha d - m > 0$ and that $\alpha d - m > 0$ is equivalent to assumption (5.2). □

Now we define the transition operator by

$$(\Psi_\varepsilon u)_j(x) := \varepsilon^{m/2}(N_j u(x) + \rho(x)(C_k u - N_j u(x^0)) \quad \text{for } x \in I_{jk} \tag{5.10}$$

where $\rho\colon \mathbb{R} \longrightarrow [0,1]$ is a smooth function such that

$$\rho(x^0) = 1 \quad \text{and} \quad \rho(x) = 0 \quad \text{for all } |x - x^0| \ge \frac{1}{2} \min_{j \in J} \ell(I_j) \tag{5.11}$$

where $\ell(I_j)$ denotes the length of the edge $e_j \cong I_j$. Furthermore, $x^0 = x^0_{jk} \in \partial I_j$ is the edge point which can be identified with the vertex v_k. Recall that I_{jk} denotes the (closed) half of the interval $I_j \cong e_j$ adjacent with the vertex v_k and directed away from v_k.

Lemma 5.9. *We have* $\Psi_\varepsilon u \in \mathcal{H}^1(M_0)$ *if* $u \in \mathcal{H}^1(M_\varepsilon)$. *Furthermore,*

$$\|u\|^2_{M_\varepsilon} - \|\Psi_\varepsilon u\|^2_{M_0} \le o(1)(\|u\|^2_{M_\varepsilon} + \|du\|^2_{M_\varepsilon}) \tag{5.12}$$

$$q_0(\Psi_\varepsilon u) - \|du\|^2_{M_\varepsilon} \le o(1)(\|u\|^2_{M_\varepsilon} + \|du\|^2_{M_\varepsilon}) \tag{5.13}$$

for all $u \in \mathcal{H}^1(M_\varepsilon)$.

Proof. The first assertion follows from $(\Psi_\varepsilon u)_j(x^0_{jk}) = C_k u$. Furthermore, we have

$$\begin{aligned}
&\|u\|^2_{M_\varepsilon} - \|\Psi_\varepsilon u\|^2_{M_0} \\
&\quad\le \sum_{k\in K}\Big(\|u\|^2_{V_{\varepsilon,k}} + \sum_{j\in J_k}\big(\|u\|^2_{U_{\varepsilon,jk}} - \varepsilon^m\|Nu + \rho\cdot(Cu - Nu(x^0))\|^2_{I_{jk}}\big)\Big) \\
&\qquad\le \sum_{k\in K}\Big(\|u\|^2_{V_{\varepsilon,k}} + \sum_{j\in J_k}\big(\|u\|^2_{U_{\varepsilon,jk}} - \|\varepsilon^{m/2}Nu\|^2_{I_{jk}}\big) \\
&\qquad\qquad + \sum_{j\in J_k}\big(\delta\|\varepsilon^{m/2}Nu\|^2_{I_{jk}} + \varepsilon^m\delta^{-1}\|\rho\|^2_{I_{jk}}|Cu - Nu(x^0)|^2\big)\Big)
\end{aligned}$$

where we have used the inequality

$$(a+b)^2 \ge (1-\delta)a^2 - \frac{1}{\delta}b^2, \qquad \delta > 0. \tag{5.14}$$

The last term in the sum can be estimated by $O(\varepsilon^m)\delta^{-1}|Cu - Nu(x^0)|^2$. Applying Lemma 5.5 we arrive at the bound by $O(\varepsilon^{m+2\alpha-d})\delta^{-1}(\|u\|^2_{M_\varepsilon} + \|du\|^2_{M_\varepsilon})$. Note that $m + 2\alpha - d = 2\alpha - 1 > 0$ since $\alpha > 1/2$. Set $\delta := \varepsilon^{(2\alpha-1)/2}$. The remaining terms can be estimated by Corollary 5.8, Lemma 4.5, and estimate (4.10).

The second inequality can be proven in the same way, namely

$$\begin{aligned}
&q_0(\Psi_\varepsilon u) - \|du\|^2_{M_\varepsilon} \le \sum_{\substack{k\in K\\ j\in J_k}}\Big(\varepsilon^m\|(Nu)' + \rho'\cdot(Cu - Nu(x^0))\|^2_{I_{jk}} - \|du\|^2_{U_{\varepsilon,jk}}\Big) \\
&\le \sum_{\substack{k\in K\\ j\in J_k}}\big(\|\varepsilon^{m/2}(Nu)'\|^2_{I_{jk}} - \|du\|^2_{U_{\varepsilon,jk}} + \delta\|\varepsilon^{m/2}(Nu)'\|^2_{I_{jk}} + \frac{2\varepsilon^m}{\delta}\|\rho'\|^2_{I_{jk}}|Cu - Nu(x^0)|^2\big)
\end{aligned}$$

where we have used

$$(a+b)^2 \le (1+\delta)a^2 + \frac{2}{\delta}b^2, \qquad 0 < \delta \le 1, \tag{5.15}$$

with $\delta := \varepsilon^{(2\alpha-1)/2}$. Since the norm involving ρ' is a fixed constant, the result follows from Lemma 4.6 and Lemma 5.5. □

Using Lemma 5.9 we arrive at the sought lower bound. Note that the error term η_k in (2.11) can be estimated by some ε-independent quantity because $\lambda_k = \lambda_k(M_\varepsilon) \le c_k$ by the upper bound given in Theorem 5.3.

Theorem 5.10. *We have* $\lambda_k(M_0) \le \lambda_k(M_\varepsilon) + o(1)$.

Theorem 5.2 now follows easily by combining the last result with Theorem 5.3.

Remark 5.11. If we allow a maximal shrinking factor $\alpha'' > 1$ in (5.1), i.e.,

$$c_-\varepsilon^{2\alpha''}g \le g_\varepsilon \le c_+\varepsilon^{2\alpha}g$$

we would have to introduce the bottle necks already in this section and we would need

$$\alpha \le \alpha'' < \frac{2}{d}\alpha + \frac{m}{d};$$

the second inequality is due to Lemma 5.5 since we would need $m+(2\alpha-\alpha'' d) > 0$ in the proof of Lemma 5.9. We have omitted this general setting to keep the previous section simple.

6. Slowly decaying vertex volume

If the volume of the vertex region decays significantly slower than the volume of the edge neighbourhoods, the limit operator is different. At the ends of the edges we have Dirichlet boundary conditions, whereas for each vertex v_k, $k \in K$, we obtain an additional eigenmode. In other words, we add a point measure at each vertex to the given measure on the graph M_0; the corresponding Hilbert space and quadratic form (domain) is therefore given by

$$\mathcal{H}_0 := L_2(M_0) \oplus \mathbb{C}^K, \qquad \mathcal{D}_0 := \bigoplus_j \mathring{\mathcal{H}}^1(I_j) \oplus \mathbb{C}^K, \qquad q_0(u) := \sum_j \|u_j'\|^2_{I_j}. \quad (6.1)$$

For elements of $\mathcal{H}_0$ we write $u = ((u_j)_{j\in J}, (u_k)_{k\in K})$ where $u_j \in L_2(I_j, p_j(x)dx)$ and $u_k \in \mathbb{C}$. We sometimes omit the indices and simply write u instead of u_j. Note that the point contributions u_k do not occur in the quadratic form, i.e., the additional eigenmodes have zero energy. Furthermore, the associated operator

$$Q_0 := \bigoplus_{j\in J} \Delta^{\mathrm{D}}_{I_j} \oplus \mathbf{0}$$

corresponds to a fully decoupled graph, i.e., a collection of independent edges, and its spectrum consists of all Dirichlet eigenvalues of the intervals I_j and 0. Here, $\mathbf{0}$ corresponds to the zero operator on $\mathbb{C}^K$.

In order to define assumptions such that a smooth metric g_ε exists globally with different length scalings on the vertex and edge neighbourhoods, we need to introduce some additional notation (see Figure 3): let V_k^- be a closed submanifold of V_k of the same dimension with a positive distance from all adjacent edge neighbourhoods U_{jk}, $j \in J_k$. Furthermore, we assume that the cylindrical structure of the half vertex neighbourhood U_{jk} extends to the component of $V_k \setminus V_k^-$ where U_{jk} meets V_k, i.e., the closure of $V_k \setminus V_k^-$ is diffeomorphic to the disjoint union of cylinders $[0,1] \times F$. We denote the extended cylinder containing U_{jk} together with the corresponding cylindrical end (the *bottle neck*) of V_k by $U_{jk}^+ = I_{jk}^+ \times F$ and the bottle neck alone by $A_{jk} = I_{jk}^0 \times F$. Note that $A_{jk} = U_{jk}^+ \cap V_k$ and that $I_{jk}^+ = I_j \cup I_{jk}^0$.

Again, we use the subscript ε to indicate the corresponding Riemannian manifold with metric g_ε.



FIGURE 3. The decomposition with the different scaling areas.

6.A. **Assumption on the smaller vertex neighbourhood.** We first fix the scaling behaviour on the smaller vertex neighbourhood V_k^-. Here, we assume that

$$c_-\varepsilon^{2\alpha} g \le g_\varepsilon \le c_+\varepsilon^{2\alpha'} g \qquad \text{on } V_k^- \tag{6.2}$$

(for the notation see (5.1)) where

$$0 < \alpha < \frac{d-1}{d}, \tag{6.3}$$

i.e., $V^-_{\varepsilon,k}$ scales at most as ε^α in each direction and at least as $\varepsilon^{\alpha'}$ where

$$\frac{d}{d+2}\alpha < \alpha' \le \alpha, \tag{6.4}$$

e.g., a homogenious scaling ($\alpha' = \alpha$) would do. Note that $\alpha' \le \alpha$ is necessary in order that (6.2) makes sense whereas $\alpha d/(d+2) < \alpha'$ ensures that the second Neumann eigenvalue of V_ε^- tends to ∞ as we will need in Lemma 6.5.

6.B. **Assumptions on the bottle neck.** Roughly speaking, we have to avoid that the bottle neck has more than a single neck separating $V_{\varepsilon,k}$ in more than one part as $\varepsilon \to 0$. In that case more than one zero eigenmode occur in the limit. Such a counterexample will be given in Remark 6.12.

We use the same notation as in Section 4 for the metric g_ε on the bottle neck $A = A_{jk}$ and set

$$\tilde{g}_\varepsilon := a_\varepsilon^2(x)dx^2 + r_\varepsilon^2(x)h(y), \qquad (x,y) \in A = I^0 \times F \tag{6.5}$$

for the (pure) product metric on A. Here, $a_\varepsilon = a_{\varepsilon,jk}$ and $r_\varepsilon = r_{\varepsilon,jk}$ are strictly positive smooth functions. Note that r_ε defines the radius of the fibre $\{x\} \times F$ at the point x. Again, we denote by G_ε and $\tilde{G}_\varepsilon$ the $d \times d$-matrices associated to the metrics g_ε and $\tilde{g}_\varepsilon$ with respect to the coordinates $(x, y) \in I^0 \times F$) and assume that the two metrics coincide up to an error term as $\varepsilon \to 0$, more specifically

$$G_\varepsilon = \tilde{G}_\varepsilon + \begin{pmatrix} o(a_\varepsilon^2) & o(a_\varepsilon r_\varepsilon) \\ o(a_\varepsilon r_\varepsilon) & o(r_\varepsilon^2) \end{pmatrix} = \begin{pmatrix} (1+o(1))a_\varepsilon^2 & o(a_\varepsilon r_\varepsilon) \\ o(a_\varepsilon r_\varepsilon) & (H+o(1))r_\varepsilon^2 \end{pmatrix}, \tag{6.6}$$

uniformly on A.

We prove the following lemma in the same way as Lemma 4.4:

Lemma 6.1. *Suppose that g_ε, $\tilde{g}_\varepsilon$ are given as above then*

$$(\det G_\varepsilon)^{\frac{1}{2}} = (1+o(1))\,(\det \tilde{G}_\varepsilon)^{\frac{1}{2}} \tag{6.7}$$

$$g_\varepsilon^{xx} := (G_\varepsilon^{-1})_{xx} = a_\varepsilon^{-2}(1+o(1)) \tag{6.8}$$

$$a_\varepsilon^{-2}|d_x u|^2 \le O(1)\,|du|^2_{g_\varepsilon} \tag{6.9}$$

where d_x denotes the partial derivative with respect to x.

To make a smooth junction between the metrics on U_j and V_k^- possible, we assume that

$$\begin{aligned} &a_\varepsilon(x) = \varepsilon^\alpha, && r_\varepsilon(x) = \varepsilon^\alpha && \text{near } x^+ \\ &a_\varepsilon(x) = 1, && r_\varepsilon(x) = \varepsilon r_- && \text{near } x^0 \end{aligned}$$

where $x \in I^0 = [x^+, x^0]$ and $r_- := r_j(x^0)$ (the radius of the fibre at x^0, see also equation (4.1)).

Furthermore, we assume that

$$a_\varepsilon(x) \le \begin{cases} \varepsilon^\alpha & \text{on } [x^+, x^0 - \delta_0,] \\ 1 & \text{on } [x^0 - \delta_0, x^0] \end{cases} \qquad \varepsilon r_- \le r_\varepsilon(x) \le \begin{cases} \varepsilon^\alpha & \text{on } [x^+, x^+ + \delta_+] \\ \varepsilon r_+ & \text{on } [x^+ + \delta_+, x^0] \end{cases} \tag{6.10}$$

for some constant $r_+ \ge r_-$, where $\delta_0 = \varepsilon^\alpha$ and $\delta_+ = \varepsilon^{(1-\alpha)m} = \varepsilon^\alpha \varepsilon^{m-\alpha d}$. These



FIGURE 4. The functions a_ε and r_ε in its allowed area (in grey).

assumptions are needed in Lemma 6.7, e.g. to assure that the eigenfunctions of M_ε do not concentrate on $A_{\varepsilon,jk}$ (i.e., (6.18) holds).

6.C. **Convergence of the spectra.** With the above prerequisites we can finally formulate the main result of this section:

Theorem 6.2. *Under the stated assumptions $\lambda_k(M_\varepsilon) \to \lambda_k(Q_0)$ as $\varepsilon \to 0$. More precisely, the first $|K|$ eigenvalues tend to 0, while the remaining bounded eigenvalue branches tend to Dirichlet eigenvalues of the intervals I_j, i.e.,*

$$\lambda_k(M_\varepsilon) \to 0 \qquad \text{if } 1 \le k \le |K| \tag{6.11}$$

$$\lambda_k(M_\varepsilon) \to \lambda^{\mathrm{D}}_{k-|K|}(\dot{\bigcup}_{j\in J} I_j) \qquad \text{if } k > |K|, \tag{6.12}$$

where $\lambda^{\mathrm{D}}_n(\dot{\bigcup}_{j\in J} I_j)$ denotes the Dirichlet eigenvalues $\lambda^{\mathrm{D}}_l(\Delta_{I_j})$ of the operators on I_j ($j \in J$) defined as in (2.2), reordered with respect to multiplicity. In particular, if the length of all the edges I_j is ℓ and $p_j(x) = 1$ for all j, we have

$$\lambda_k(M_\varepsilon) \to \lambda^{\mathrm{D}}_m([0,\ell]) = \pi^2 m^2/\ell^2 \qquad \text{if } k = (m-1)|J| + 1, \dots, m|J|. \tag{6.13}$$

Again, our aim is to find a two sided estimate on each eigenvalue $\lambda_k(M_\varepsilon)$ by means of $\lambda_k(Q_0)$ with an error which is $o(1)$ w.r.t. the parameter ε.

6.D. **An upper bound.** The following upper eigenvalue estimate is slightly more difficult to show than in the previous section:

Theorem 6.3. *$\lambda_k(M_\varepsilon) \le \lambda_k(Q_0) + o(1)$ holds as $\varepsilon \to 0$.*

To prove it, we define the transition operator by

$$\Phi_\varepsilon u(z) := \begin{cases} (\operatorname{vol} V^-_{\varepsilon,k})^{-1/2} u_k & \text{if } z \in V_k, \\ \varepsilon^{-m/2} u_j(x) + (\operatorname{vol} V^-_{\varepsilon,k})^{-1/2} \rho(x) u_k & \text{if } z = (x,y) \in U_j \end{cases}$$

for any $u \in \mathcal{D}_0$, where ρ is a smooth function as in (5.11) and $x^0 = x^0_{jk}$ denotes the endpoint of the half-edge I_{jk} corresponding to the vertex v_k. Theorem 6.3 is then implied by Lemma 2.1 in combination with the following result.

Lemma 6.4. *We have $\Phi_\varepsilon u \in \mathcal{H}^1(M_\varepsilon)$, i.e., Φ_ε maps the quadratic form domain $\mathcal{D}_0$ into the quadratic form domain of the Laplacian on the manifold. Furthermore,*

$$\|u\|^2_{\mathcal{H}_0} - \|\Phi_\varepsilon u\|^2_{M_\varepsilon} \le o(1)\, \|u\|^2_{\mathcal{H}_0} \tag{6.14}$$

$$\|d\,\Phi_\varepsilon u\|^2_{M_\varepsilon} - q_0(u) \le o(1)\, (\|u\|^2_{\mathcal{H}_0} + q_0(u)) \tag{6.15}$$

Proof. Since $u_j\restriction_{\partial I_j} = 0$, the function $\Phi_\varepsilon u$ agrees on $\partial V_{\varepsilon,k}$ for both definitions. Clearly, $\Phi_\varepsilon u$ is weakly differentiable on each thickened edge $U_{\varepsilon,j}$. Moreover, we

have

$$\|u\|^2_{\mathcal{H}_0} - \|\Phi_\varepsilon u\|^2_{M_\varepsilon} = \sum_{k\in K}\biggl(\sum_{j\in J_k}\Bigl(\|u\|^2_{I_{jk}} - \|\Phi_\varepsilon u\|^2_{U_{\varepsilon,jk}}\Bigr) + \Bigl(|u_k|^2 - \|\Phi_\varepsilon u\|^2_{V_{\varepsilon,k}}\Bigr)\biggr)$$

$$\le \sum_{\substack{k\in K\\ j\in J_k}}\biggl(\Bigl(\|u\|^2_{I_{jk}} - (1+o(1))\Bigl\|\varepsilon^{-m/2}u + (\operatorname{vol} V^-_{\varepsilon,k})^{-1/2}\rho\, u_k\Bigr\|^2_{\tilde U_{\varepsilon,jk}}\Bigr)\biggr) + \sum_{k\in K}\Bigl(|u_k|^2 - \|\Phi_\varepsilon u\|^2_{V^-_{\varepsilon,k}}\Bigr)$$

where we have used equation (4.4) and that $V^-_{\varepsilon,k}\subset V_{\varepsilon,k}$. Note that the latter sum in the last line is equal to 0. To estimate the remaining sum, remember that $\Phi_\varepsilon u$ is independent of y on $U_{\varepsilon,jk}$. Therefore we can apply equation (4.8), and inequality (5.14) with $\delta := \varepsilon^{(m-\alpha d)/2}$ yields the upper estimate

$$\sum_{\substack{k\in K\\ j\in J_k}}\biggl(\|u\|^2_{I_{jk}} - (1+o(1))\Bigl((1-\delta)\|u\|^2_{I_{jk}} - \frac{\varepsilon^m}{\delta\operatorname{vol} V^-_{\varepsilon,k}}\|\rho\|^2_{I_{jk}}|u_k|^2\Bigr)\biggr)$$

$$\le \sum_{\substack{k\in K\\ j\in J_k}}\Bigl((\delta+o(1))\|u\|^2_{I_{jk}} + \frac{O(\varepsilon^{m-\alpha d})}{\delta}|u_k|^2\Bigr) = o(1)\|u\|^2_{\mathcal{H}_0}.$$

In the last inequality, we have used the estimate $(\operatorname{vol} V^-_{\varepsilon,k})^{-1}\le O(\varepsilon^{-\alpha d})$ which follows from the lower bound of (6.2). Note that $\delta = o(1)$ by assumption (6.3). The second relation follows from

$$\|d\,\Phi_\varepsilon u\|^2_{M_\varepsilon} - q_0(u) = \sum_{\substack{k\in K\\ j\in J_k}}\Bigl(\bigl(1+o(1)\bigr)\|g^{xx}_\varepsilon d_x\Phi_\varepsilon u\|^2_{\tilde U_{\varepsilon,jk}} - \|u'\|^2_{U_{\varepsilon,jk}}\Bigr)$$

$$= \sum_{\substack{k\in K\\ j\in J_k}}\Bigl(\bigl(1+o(1)\bigr)\varepsilon^m\|\varepsilon^{-m/2}u' + (\operatorname{vol} V^-_{\varepsilon,k})^{-1/2}\rho'\,u_k\|^2_{I_{jk}} - \|u'\|^2_{I_{jk}}\Bigr)$$

$$\le \sum_{\substack{k\in K\\ j\in J_k}}\bigl(1+o(1)\bigr)\Bigl(\delta\|u'\|^2_{I_{jk}} + \frac{2\varepsilon^m}{\delta\operatorname{vol} V^-_{\varepsilon,k}}\|\rho'\|^2_{I_{jk}}|u_k|^2\Bigr)$$

$$\le o(1)\bigl(\|u\|^2_{\mathcal{H}_0} + q_0(u)\bigr)$$

in the same way as above together with (4.5) for the second equality and (5.15) in the last line; recall that $\Phi_\varepsilon u$ is constant on $V_{\varepsilon,k}$. □

Note that we need a counterpart to $\|u\|^2_{V_{\varepsilon,k}}$ on the limit problem Hilbert space $\mathcal{H}_0$. In the case of a fast decaying vertex volume in the previous section, the correponding norm vanished (see Corollary 5.8), but here we need the additional subspace $\mathbb{C}^K$ in $\mathcal{H}_0$ coming from extra point measures at the vertices.

Furthermore, note that the upper bound estimate on $\lambda_k(M_\varepsilon)$ already proven in Lemma 5.4 remains valid in this setting, but it is too rough for the present purpose.

6.E. **A lower bound.** Again, the opposite estimate is more difficult. We will employ averaging processes also on the vertex neigbourhoods; this time with the ε-scaled manifold $V^-_{\varepsilon,k}$:

$$C^-_\varepsilon u = C^-_{\varepsilon,k} u := \frac{1}{\operatorname{vol} V^-_{\varepsilon,k}} \int_{V^-_{\varepsilon,k}} u \, dV^-_{\varepsilon,k}. \tag{6.16}$$

In the first lemma, we prove an estimate similar to the one in Lemma 5.6. Note that $\|C^-_\varepsilon u\| \le \|u\|$ by Cauchy-Schwarz, but we need the opposite inequality.

Lemma 6.5. *For any $u \in \mathcal{H}^1(V^-_\varepsilon)$ we have*

$$\|u\|^2_{V^-_\varepsilon} - \|C^-_\varepsilon u\|^2_{V^-_\varepsilon} \le o(1)\big(\|u\|^2_{V^-_\varepsilon} + \|du\|^2_{V^-_\varepsilon}\big).$$

Proof. Apply Lemma 3.1 with $X = V^-_\varepsilon$ and $\delta = \varepsilon^{((d+2)\alpha' - d\alpha)/2}$. From the min-max principle we obtain $\lambda^{\mathrm{N}}_2(V^-_\varepsilon) \ge O(\varepsilon^{d(\alpha-\alpha')-2\alpha'})$. Note that $\delta = o(1)$ since $(d+2)\alpha' - d\alpha > 0$ by (6.4). □

The next three results are valid independently of the assumptions on α given in (6.3) and (6.4). We will need these results also for the borderline case $\alpha = (d-1)/d$ in the next section.

We need an estimate on the average $|Nu(x^0)|^2$. Since on the bottle neck $A_{\varepsilon,jk}$, the estimates are quite delicate, we first prove the result for $|Nu(x^+)|^2$ (i.e., on $\partial V^-_{\varepsilon,k}$ where the scaling of the metric is of the right order. The error is controlled by (6.17). Note that this estimate is a counterpart to the estimate in Lemma 4.7 where we extended the function to the *edge* neighbourhood $U_{\varepsilon,j}$ (useful in the case of fast decaying vertex volume, $\alpha d - m > 0$). This is not possible here, since $\alpha d - m < 0$. Therefore, we extend the function to the *vertex* neighbourhood $V^-_{\varepsilon,k}$.

Lemma 6.6. *The inequality*

$$|Nu(x^+)|^2 \le \|u(x^+,\cdot)\|^2_F \le O(\varepsilon^{-\alpha d}) \left(\|u\|^2_{V^-_\varepsilon} + \|du\|^2_{V^-_\varepsilon}\right)$$

holds for any $u \in \mathcal{H}^1(V^-_\varepsilon)$.

Proof. We have

$$\begin{aligned} |Nu(x^+)|^2 &\le \int_F |u(x^+,y)|^2 dF(y) \\ &\le c_1\big(\|u\|^2_{V^-} + \|du\|^2_{V^-}\big) \le O(\varepsilon^{-\alpha d})\big(\|u\|^2_{V^-_\varepsilon} + \|du\|^2_{V^-_\varepsilon}\big) \end{aligned}$$

by Lemma 3.2 with $X = V^-$ and the lower bound in assumption (6.3). □

The next lemma is the key ingredient in dealing with the bottle neck. Here, we prove two Poincaré-like estimates. Since we want to avoid a cut-off function (leading to divergent terms when being differentiated) we only prove an estimate on the *difference* and not on $Nu(x^0)$ itself in (6.17). For the same reason, an integral over F remains in (6.18). Note that $I^+_\varepsilon(x^0) = \operatorname{vol} A_\varepsilon$.

Lemma 6.7. *There is a constant $C > 0$ such that*

$$|Nu(x^0) - Nu(x^+)|^2 \le C I^-_\varepsilon(x^0) \, \|du\|^2_{A_\varepsilon}, \tag{6.17}$$

$$\|u\|^2_{A_\varepsilon} \le 4 I^+_\varepsilon(x^0) \, \|u(x^+,\cdot)\|^2_F + 4C I^{+-}_\varepsilon \, \|du\|^2_{A_\varepsilon} \tag{6.18}$$

for all $u \in \mathcal{H}^1(A_\varepsilon)$ where

$$I^\pm_\varepsilon(x) := \int_{x+}^{x} a_\varepsilon(x') r^{\pm m}_\varepsilon(x') \, dx' \quad \text{and} \quad I^{+-}_\varepsilon := \int_{x^+}^{x^0} a_\varepsilon(x) r^m_\varepsilon(x) I^-_\varepsilon(x) \, dx.$$

Furthermore, under the assumption (6.10)*, we have*

$$I^-_\varepsilon(x^0) = o(\varepsilon^{-m}), \qquad I^+_\varepsilon(x^0) = o(\varepsilon^{\alpha d}) \qquad \text{and} \qquad I^{+-}_\varepsilon = o(1).$$

Proof. For a smooth function u we have

$$u(x,y) - u(x^+,y) = \int_{x^+}^{x} \partial_x u(x',y) \, dx'. \tag{6.19}$$

For the first assertion, we set $x = x^0$, foremost integrate over $y \in F$ and then apply Cauchy-Schwarz

$$|Nu(x^0) - Nu(x^+)|^2 \le \int_F \int_{x^+}^{x^0} a^2_\varepsilon(x') \det G_\varepsilon(x',y)^{-\frac12} \, dx' \times \\ \times \int_{x^+}^{x^0} a^{-2}_\varepsilon(x') |\partial_x u(x',y)|^2 \det G_\varepsilon(x',y)^{\frac12} \, dx' \, dF(y).$$

The first integrand over x' can be estimated by $C a_\varepsilon(x') r^{-m}_\varepsilon$ applying (6.7). Therefore, the first integral is smaller than $C I^-_\varepsilon(x^0)$. The second integral together with the integral over F can be estimated by $O(1) \, \|du\|^2_{A_\varepsilon}$ applying (6.9).

For the second assertion, we first apply Cauchy-Schwarz (and (5.15) with $\delta = 1$) to (6.19) and than integrate over $y \in F$ to obtain

$$\int_F |u(x,y)|^2 \, dF(y) \le 2 \int_F |u(x^+,y)|^2 \, dF(y) + 2 \int_F \int_{x^+}^{x} a^2_\varepsilon(x') \det G(x',y)^{-\frac12} \, dx' \times \\ \times \int_{x^+}^{x} a^{-2}_\varepsilon(x') \, |\partial_x u(x',y)|^2 \det G(x',y)^{\frac12} \, dx' \, dF(y). \tag{6.20}$$

The first integral over x' can be estimated as before by $C I^-_\varepsilon(x)$. Finally, multiplying with $a_\varepsilon(x) r^m_\varepsilon(x)$ and integrating over $x \in I^0$ yields

$$\|u\|^2_{\tilde{A}_\varepsilon} \le 2 I^+_\varepsilon(x_0) \, \|u(x^+,\cdot)\|^2_F + 2C I^{+-}_\varepsilon \, \|du\|^2_{A_\varepsilon}.$$

Applying (6.7) once more we obtain the desired estimate over A_ε instead of $\tilde{A}_\varepsilon$ (note that $2/(1+o(1)) \le 4$ provided ε is small enough). The general case of non-smooth functions can easily shown with approximation arguments.

The integral estimates follow from

$$I_\varepsilon^-(x^0) \le \int_{x^+}^{x^0-\delta_0} \varepsilon^\alpha (\varepsilon r_-)^{-m}\,dx + \int_{x^0-\delta_0}^{x^0} (\varepsilon r_-)^{-m}\,dx = (\varepsilon^\alpha + \delta_0) O(\varepsilon^{-m}).$$

Since $\delta_0 = \varepsilon^\alpha$, we have $I_\varepsilon^-(x^0) \le O(\varepsilon^{\alpha-m})$. Next, we have

$$\begin{aligned} I_\varepsilon^+(x^0) \le \int_{x^+}^{x^++\delta_+} \varepsilon^\alpha \varepsilon^{\alpha m}\,dx + \int_{x^++\delta_+}^{x^0-\delta_0} \varepsilon^\alpha (\varepsilon r_+)^m\,dx \\ + \int_{x^0-\delta_0}^{x^0} (\varepsilon r_+)^m\,dx = \delta_+ O(\varepsilon^{\alpha d}) + (\varepsilon^\alpha + \delta_0) O(\varepsilon^m) \end{aligned}$$

and therefore $I_\varepsilon^+(x^0) \le O(\varepsilon^{\alpha+m}) = O(\varepsilon^{\alpha+m-\alpha d}) O(\varepsilon^{\alpha d})$ since $\delta_+ = \varepsilon^\alpha \varepsilon^{m-\alpha d}$. The last assertion follows from $I_\varepsilon^{+-} \le I_\varepsilon^-(x_0)\, I_\varepsilon^+(x_0) \le O(\varepsilon^{2\alpha})$. □

The following corollary is again independent of the assumption we made about α in (6.3) and (6.4), in particular, it is also valid in the setting of the borderline case of Section 7.

Corollary 6.8. *For all $u \in \mathcal{H}^1(V_\varepsilon)$ we have*

$$\|u\|_{A_\varepsilon}^2 \le o(1)\,\big(\|u\|_{V_\varepsilon}^2 + \|du\|_{V_\varepsilon}^2\big).$$

Proof. We only have to put together (6.18) and Lemma 6.6. □

We now formulate a consequence of the preceding lemmas under the assumption (6.3).

Corollary 6.9. *Suppose $0 < \alpha < m/d = (d-1)/d$. Then we have*

$$|Nu(x^0)|^2 \le o(\varepsilon^{-m})\,\big(\|u\|_{V_\varepsilon}^2 + \|du\|_{V_\varepsilon}^2\big)$$

for all $u \in \mathcal{H}^1(V_\varepsilon)$.

Proof. Applying (5.14) with $\delta = 1/2$ to (6.17) we obtain

$$|Nu(x^0)|^2 \le o(\varepsilon^{-m}) \|du\|_{A_\varepsilon}^2 + 4|Nu(x^+)|^2.$$

The second term is of order $O(\varepsilon^{-\alpha d})$ by Lemma 6.6 and therefore also of order $o(\varepsilon^{-m})$ by the assumption on α. □

In this section, we define the transition operator by

$$\begin{aligned} (\Psi_\varepsilon u)_j(x) &:= \varepsilon^{m/2} N_j u(x) - \rho(x) N_j u(x^0) \quad \text{for } x \in I_{jk} \\ (\Psi_\varepsilon u)_k &:= (\operatorname{vol} V_{\varepsilon,k}^-)^{1/2} C_{\varepsilon,k}^- u \end{aligned} \tag{6.21}$$

where ρ is a smooth function as in (5.11) and $x^0 = x_{jk}^0$ denotes the endpoint of the half-edge I_{jk} corresponding to the vertex v_k.

Lemma 6.10. *We have $\Psi_\varepsilon u \in \mathcal{D}_0$ if $u \in \mathcal{H}^1(M_\varepsilon)$. Furthermore,*

$$\|u\|^2_{M_\varepsilon} - \|\Psi_\varepsilon u\|^2_{\mathcal{H}_0} \le o(1)\big(\|u\|^2_{M_\varepsilon} + \|du\|^2_{M_\varepsilon}\big) \tag{6.22}$$

$$q_0(\Psi_\varepsilon u) - \|du\|^2_{M_\varepsilon} \le o(1)\big(\|u\|^2_{M_\varepsilon} + \|du\|^2_{M_\varepsilon}\big) \tag{6.23}$$

for all $u \in \mathcal{H}^1(M_\varepsilon)$.

Proof. The first assertion follows from the fact that $(\Psi_\varepsilon u)_j(x^0) = 0$. Furthermore, we have

$$\|u\|^2_{M_\varepsilon} - \|\Psi_\varepsilon u\|^2_{\mathcal{H}_0} \le \sum_{k\in K}\bigg(\big(\|u\|^2_{V^-_{\varepsilon,k}} - \|C^-_\varepsilon u\|^2_{V^-_{\varepsilon,k}}\big) + \sum_{j\in J_k}\big(\|u\|^2_{A_{\varepsilon,jk}} + \|u\|^2_{U_{\varepsilon,jk}} - \varepsilon^m\big\|Nu - \rho\cdot Nu(x^0)\big\|^2_{I_{jk}}\big)\bigg).$$

The first difference is of the desired form by Lemma 6.5. Furthermore, the integral over the "bottle necks" $A_{\varepsilon,jk}$ can be estimated in the needed way by Corollary 6.8. Applying (5.14) to the remaining difference in the last sum we obtain the upper estimate by

$$\big(\|u\|^2_{U_{\varepsilon,jk}} - \varepsilon^m\|Nu\|^2_{I_{jk}}\big) + \delta\,\varepsilon^m\|Nu\|^2_{I_{jk}} + \frac{\varepsilon^m}{\delta}\|\rho\|^2_{I_{jk}}|Nu(x^0)|^2 \tag{6.24}$$

For the first two terms we obtain the sought bound by virtue of Lemma 4.5 and (4.10); for the remaining term one has to apply Corollary 6.9.

The second inequality can be proven in the same way, namely

$$q_0(\Psi_\varepsilon u) - \|du\|^2_{M_\varepsilon} = \sum_{k\in K}\Big(- \|du\|^2_{V^-_{\varepsilon,k}} + \sum_{j\in J_k}\big(\varepsilon^m\big\|(Nu)' - \rho'\,Nu(x^0)\big\|^2_{I_{jk}} - \|du\|^2_{U_{\varepsilon,jk}}\big)\Big)$$

We omit the norm contribution from $V^-_{\varepsilon,k}$ and estimate the remaining difference with (5.15) and obtain (up to the summation)

$$\big(\varepsilon^m\|(Nu)'\|^2_{I_{jk}} - \|du\|^2_{U_{\varepsilon,jk}}\big) + \delta\,\varepsilon^m\|(Nu)'\|^2_{I_{jk}} + 2\frac{\varepsilon^m}{\delta}\|\rho'\|^2_{I_{jk}}|Nu(x^0)|^2.$$

For the first difference we obtain the needed estimate by virtue of Lemma 4.6. An upper bound for the remaining term is of the same form as before. □

Using Lemma 6.10 we arrive at the sought lower bound. Note that the error term η_k in (2.11) can be estimated by some ε-independent quantity because $\lambda_k = \lambda_k(M_\varepsilon) \le c_k$ by Theorem 6.3.

Theorem 6.11. *We have $\lambda_k(Q_0) \le \lambda_k(M_\varepsilon) + o(1)$.*

Theorem 6.2 now follows easily by combining the last result with Theorem 6.3.

Remark 6.12. Without assumption (6.10) on the metric on the bottle neck A_{jk}, the second Neumann eigenvalue of $V_{\varepsilon,k}$ could tend to 0 (and not to ∞, as required in the proof of Lemma 6.5), for example, if $V_{\varepsilon,k}$ separates into more than one part as $\varepsilon \to 0$ (i.e., $V_{\varepsilon,k}$ has an additional "throat").

7. The borderline case

7.A. **Definition of the thickened vertices.** If the volume of the vertex region decays at the same rate as the volume of the edge neighbourhoods, the limit operator acts again in the extended Hilbert space introduced in the previous section but it is not decoupled anymore. Thus it is not supported by the graph alone, in particular, it is not the Hamiltonian with the boundary conditions (2.4).

We start with the definition of the limit operator. The corresponding Hilbert space and quadratic form are given by

$$\mathcal{H}_0 := L_2(M_0) \oplus \mathbb{C}^K, \qquad q_0(u) := \sum_j \|u_j'\|^2_{I_j}\,, \tag{7.1}$$

where the form domain $\mathcal{D}_0$ of q_0 is given by those functions $u = ((u_j)_{j\in J}, (u_k)_{k\in K})$ such that

$$u \in \mathcal{H}^1(M_0) \oplus \mathbb{C}^K \qquad \text{and} \qquad (\operatorname{vol} V_k^-)^{1/2} u_j(v_k) = u_k \tag{7.2}$$

for all $j \in J_k$ and $k \in K$, i.e., values of the functions at the edge endpoints $v_k \equiv x^0_{jk}$ are now coupled with the additional wave function components; recall that V_k^- denotes the manifold $V^-_{\varepsilon,k}$ with $\varepsilon = 1$. The corresponding operator Q_0 is given by

$$Q_0 u = \left(\left(-\frac{1}{p_j}(p_j u_j')' \right)_j, \left(-(\operatorname{vol} V_k^-)^{-\frac{1}{2}} \sum_{j\in J_k} p_j(v_k) u_j'(v_k) \right)_k \right); \tag{7.3}$$

it depends parametrically on $\operatorname{vol}(V_k^-)$ but we refrain from marking this fact explicitly. Again, this operator has a purely discrete spectrum provided the graph M_0 is finite.

As we have said, Q_0 is not a graph operator with the conditions (2.4). Nevertheless, there is a similarity between the two noticed by Kuchment and Zeng in [KuZ03]. To solve the spectral problem $Q_0 u = \lambda u$ one has to find $(u_j)_{j\in J}$ such that $-(p_j u_j')'/p_j = \lambda u_j$ and at the vertices the functions satisfy the conditions

$$\sum_{j\in J_k} p_j(v_k) u_j'(v_k) = -\lambda (\operatorname{vol} V_k^-)^{\frac{1}{2}} u_k\,. \tag{7.4}$$

This looks like (2.4), the difference is that the coefficient at the right-hand side is not a constant but a multiple of the spectral parameter; in physical terms one may say that the coupling strength at a vertex is proportional to the energy.

After this digression let us return to the limiting properties. We adopt again the assumption (6.10) in this section. Instead of (6.2) we suppose now that on

the vertex neighbourhood the metric satisfies the relation

$$g_\varepsilon = \varepsilon^{2\alpha} g + o(\varepsilon^{2\alpha}) \qquad \text{on } V_k^- \tag{7.5}$$

with

$$\alpha = \frac{d-1}{d}\,, \tag{7.6}$$

which corresponds to the above mentioned equal decay rate for the volume of the edge and vertex neighbourhoods. In particular, we have

$$\|u\|^2_{V_\varepsilon^-} = \varepsilon^{\alpha d}(1+o(1))\|u\|^2_{V^-} \quad \text{and} \quad \|du\|^2_{V_\varepsilon^-} = \varepsilon^{\alpha(d-2)}(1+o(1))\|du\|^2_{V^-} \tag{7.7}$$

and

$$\operatorname{vol}(V_\varepsilon^-) = \varepsilon^{\alpha d}(1+o(1))\operatorname{vol}(V^-) \tag{7.8}$$

for each $V^- = V_k^-$ as in Lemmas 4.4 and 5.1.

7.B. **Convergence of the spectra.** With the above prerequisites we can finally formulate the main result of this section:

Theorem 7.1. *Under the stated assumptions $\lambda_k(M_\varepsilon) \to \lambda_k(Q_0)$ as $\varepsilon \to 0$.*

To prove it, our aim is again to find a two sided estimate on each eigenvalue $\lambda_k(M_\varepsilon)$ by means of $\lambda_k(Q_0)$ with an error which is $o(1)$ w.r.t. the parameter ε.

7.C. **An upper bound.** Again, we first show the easier upper eigenvalue estimate:

Theorem 7.2. *$\lambda_k(M_\varepsilon) \le \lambda_k(Q_0) + o(1)$ holds as $\varepsilon \to 0$.*

We define the transition operator by

$$\Phi_\varepsilon u(z) := \begin{cases} \operatorname{vol}(V^-_{\varepsilon,k})^{-1/2} u_k & \text{if } z \in V_k, \\ \varepsilon^{-m/2} u_j(x) + \rho(x) \times & \\ \qquad \times \big(\operatorname{vol}(V^-_{\varepsilon,k})^{-1/2} u_k - \varepsilon^{-m/2} u_j(x^0)\big) & \text{if } z = (x,y) \in U_j \end{cases} \tag{7.9}$$

for any $u \in \mathcal{D}_0$, where ρ is a smooth function as in (5.11) and $x^0 = x^0_{jk}$ denotes the endpoint of the half-edge I_{jk} away from the vertex v_k. Theorem 7.2 is then implied by Lemma 2.1 in combination with the following result.

Lemma 7.3. *We have $\Phi_\varepsilon u \in \mathcal{H}^1(M_\varepsilon)$, i.e., Φ_ε maps the quadratic form domain $\mathcal{D}_0$ into the quadratic form domain of the Laplacian on the manifold. Furthermore,*

$$\|u\|^2_{\mathcal{H}_0} - \|\Phi_\varepsilon u\|^2_{M_\varepsilon} \le o(1)\,\|u\|^2_{\mathcal{H}_0} \tag{7.10}$$

$$\|d\,\Phi_\varepsilon u\|^2_{M_\varepsilon} - q_0(u) \le o(1)\,(\|u\|^2_{\mathcal{H}_0} + q_0(u)) \tag{7.11}$$

Proof. The argument is analogous to the proof of Lemma 6.4. The only difference is that we need the following estimate

$$\varepsilon^m \big|\operatorname{vol}(V^-_{\varepsilon,k})^{-1/2} u_k - \varepsilon^{-m/2} u_j(x^0)\big|^2 = \big|\varepsilon^{m/2}(\operatorname{vol} V^-_{\varepsilon,k})^{-1/2} - (\operatorname{vol} V_k^-)^{-1/2}\big|^2\,|u_k|^2$$

since $u \in \mathcal{D}_0$. The last difference is of order $o(1)$ by (7.8). □

7.D. **A lower bound.** The estimate on $\lambda_k(M_\varepsilon)$ from below can be found in analogy with the slow-decay case in Section 6. Furthermore, we need the following averaging operator

$$C_k^- u := \frac{1}{\operatorname{vol} V_k^-} \int_{V_k^-} u \, dV_k^-.$$

Since we have an exact scaling of the metric of order ε^α by (7.6), we also could use the ε-depending manifold $V_{\varepsilon,k}^-$ here (cf. also Remark 5.7).

Lemma 7.4. *For all $u \in \mathcal{H}^1(V_{\varepsilon,k}^-)$ we have*

$$\left|C_k^- u - N_j u(x^0)\right|^2 \le o(\varepsilon^{-m}) \, \|du\|^2_{V_{\varepsilon,k}}$$

Proof. We have

$$\left|C_k^- u - N_j u(x^0)\right| \le \left|C_k^- u - N_j u(x^+)\right| + \left|N_j u(x^+) - N_j u(x^0)\right|.$$

The first difference can be estimated in the same way as Lemma 5.5 (replacing V_k by V_k^- and using estimate (7.7), i.e., we arrive at

$$\left|C_k^- u - N_j u(x^+)\right|^2 \le O(\varepsilon^{-(d-2)\alpha}) \, \|du\|^2_{V_\varepsilon^-};$$

recall that now we have $\alpha d = m$. For the second difference, use (6.17). □

Lemma 7.5. *For all $u \in \mathcal{H}^1(V_\varepsilon^-)$, we have*

$$\|u\|^2_{V_\varepsilon^-} - \|C^- u\|^2_{V_\varepsilon^-} \le O(\varepsilon^\alpha)(\|u\|^2_{V_\varepsilon^-} + \|du\|^2_{V_\varepsilon^-}).$$

Proof. The proof is similar to the proof of Lemma 6.5: inequality (3.2) together with (7.7) and Lemma 3.1 for $X = V^-$ implies

$$\begin{aligned}\|u\|^2_{V_\varepsilon^-} - \|C^- u\|^2_{V_\varepsilon^-} &\le \frac{1}{\delta}\varepsilon^{\alpha d}(1+o(1))\|u - C^- u\|^2_{V^-} + \delta(\|u\|^2_{V_\varepsilon^-} + \operatorname{vol} V_\varepsilon^- |C^- u|^2) \\ &\le \frac{\varepsilon^{2\alpha}(1+o(1))}{\delta \lambda_2^{\mathrm{N}}(V^-)} \|du\|^2_{V_\varepsilon^-} + \delta(\|u\|^2_{V_\varepsilon^-} + \operatorname{vol} V_\varepsilon^- \|u\|^2_{V^-})\end{aligned}$$

Applying (7.8) and (7.7) once more, the result follows setting $\delta = \varepsilon^\alpha$. □

Now we define the transition operator by

$$\begin{aligned}(\Psi_\varepsilon u)_j(x) &:= \varepsilon^{m/2}\Big(N_j u(x) + \rho(x)\big(C_k^- u - N_j u(x^0)\big)\Big) \quad \text{for } x \in I_{jk} \\ (\Psi_\varepsilon u)_k &:= \varepsilon^{m/2}(\operatorname{vol} V_k^-)^{1/2} C_k^- u\end{aligned} \tag{7.12}$$

where ρ is a smooth function as in (5.11) and $x^0 = x_{jk}^0$ denotes the endpoint of the half-edge I_{jk} corresponding to the vertex v_k.

Lemma 7.6. *We have $\Psi_\varepsilon u \in \mathcal{D}_0$ if $u \in \mathcal{H}^1(M_\varepsilon)$. Furthermore,*

$$\|u\|^2_{M_\varepsilon} - \|\Psi_\varepsilon u\|^2_{\mathcal{H}_0} \le o(1)\big(\|u\|^2_{M_\varepsilon} + \|du\|^2_{M_\varepsilon}\big) \tag{7.13}$$

$$q_0(\Psi_\varepsilon u) - \|du\|^2_{M_\varepsilon} \le o(1)\big(\|u\|^2_{M_\varepsilon} + \|du\|^2_{M_\varepsilon}\big) \tag{7.14}$$

for all $u \in \mathcal{H}^1(M_\varepsilon)$.

Proof. The arguments are the same as in the proof of Lemma 6.10. For the vertex contribution, we need the estimate

$$\|u\|^2_{V^-_{\varepsilon,k}} - \varepsilon^m (\operatorname{vol} V_k^-)|C_k^- u|^2 = (\|u\|^2_{V^-_{\varepsilon,k}} - \|C_k^- u\|^2_{V^-_{\varepsilon,k}}) + \Big(\frac{\operatorname{vol} V^-_{\varepsilon,k}}{\varepsilon^m \operatorname{vol} V_k^-} - 1\Big)\varepsilon^m \|C_k^- u\|^2_{V_k^-}.$$

The first difference can be treated with Lemma 7.5 and leads to an error term $O(\varepsilon^\alpha)$. The second term is of order $o(1)\|u\|^2_{V^-_{\varepsilon,k}}$ by (7.7), (7.8) and Cauchy-Schwarz. Furthermore, Corollary 6.8 is also true in this setting (independent on the particular α). We also need Lemma 7.4. □

Using Lemma 7.6 we arrive at the sought lower bound. Again, the error term η_k in (2.11) can be estimated by some ε-independent quantity because $\lambda_k = \lambda_k(M_\varepsilon) \le c_k$ by Theorem 7.2.

Theorem 7.7. *We have* $\lambda_k(Q_0) \le \lambda_k(M_\varepsilon) + o(1)$.

Theorem 7.1 now follows easily by combining the last result with Theorem 7.2.

8. Non-decaying vertex volume

In this section, we treat the case when the vertex volume does not tend to 0. In some sense, this case corresponds to $\alpha = 0$ in the previous notation but we need more assumptions to precise the convergence of the manifold $V_{\varepsilon,k}$ to a manifold $V_{0,k}$ as $\varepsilon \to 0$. We cite only the result here since it has already been presented in [P03] or with a more detailed proof in [P00]. A related result corresponding to the embedded case (see Example 4.2) as in [KuZ01] was proven by Jimbo and Morita in [JM92] or for manifolds (with non-smooth junctions between edge and vertex neighbourhoods) by Anné and Colbois in [AC95].

In this section, we assume that the transversal direction is a sphere, i.e., $F = \mathbb{S}^m$. Let $V_{0,k}$ be a compact d-dimensional manifold without boundary for $k \in K$. To each edge $j \in J_k$ eminating from the vertex v_k, we associate a point $x^0_{jk} \in V_{0,k}$ such that x^0_{jk} ($j \in J_k$) are mutually distinct points with lower bound $2\varepsilon_0 > 0$ on their distance to each other. We assume for simplicity that the metric at $x^0 = x^0_{jk}$ is locally flat within a distance ε_0 from x^0 (the general case can be found in [P03]). Then the metric in polar coordinates $(x, y) \in (0, \varepsilon_0) \times \mathbb{S}^m$ looks locally like

$$g = dx^2 + x^2\, h_y$$

where h_y is the standard metric on $\mathbb{S}^m$. Modifying the factor before h_y, we define a new metric by

$$g_\varepsilon = dx^2 + r_\varepsilon^2(x)\, h_y$$

with a smooth monotone function $r_\varepsilon \colon (0, \varepsilon_0) \longrightarrow (0, \infty)$ such that

$$r_\varepsilon(x) = \begin{cases} \varepsilon & \text{for } 0 < x < \varepsilon/2 \\ x & \text{for } 2\varepsilon < x < \varepsilon_0. \end{cases}$$

We denote the (completion of the) manifold $(V_{0,k} \setminus \bigcup_{j\in J_k} x^0_{jk}, g_\varepsilon)$ by $V_{\varepsilon,k}$. Note that this manifold has $|J_k|$ attached cylindrical ends of order ε at each point x^0_{jk}. Now we can construct the graph-like manifold M_ε as in Section 3.

As in the slow decaying case of Section 6 the limit operator

$$Q_0 := \bigoplus_{j\in J} \Delta^{\mathrm{D}}_{I_j} \oplus \bigoplus_{k\in K} \Delta_{V_{0,k}}$$

decouples and the next result follows (cf. [P03, Theorem 1.2] or [P00]):

Theorem 8.1. *We have $\lambda_k(M_\varepsilon) \to \lambda_k(Q_0)$ as $\varepsilon \to 0$.*

9. Applications

Finally we comment on consequences of the spectral convergence. We begin with a general remark stating that we only have uniform control over a *compact* spectral interval:

Remark 9.1. Note that the convergence $\lambda_k(M_\varepsilon) \to \lambda_k(M_0)$ cannot be uniform in $k \in \mathbb{N}$: if this were the case, the theta-function

$$\Theta_\varepsilon(t) := \operatorname{tr} \mathrm{e}^{-t\Delta_{M_\varepsilon}} = \sum_k \mathrm{e}^{-t\lambda_k(M_\varepsilon)}$$

would converge to $\Theta_0(t)$. But Weyl asymtotics are different in the two cases,

$$\Theta_\varepsilon(t) \sim \frac{\operatorname{vol}_d M_\varepsilon}{(4\pi t)^{d/2}}, \qquad \text{whereas} \qquad \Theta_0(t) \sim \frac{\operatorname{vol}_1 M_0}{(4\pi t)^{1/2}}$$

as $t \to 0$ (cf. [Cha84, Sec. VI.4] and [Ro84, Thm. 1]). Recall that $d \geq 2$ and $\operatorname{vol}_1 M_0 := \sum_j \ell(I_j)$, i.e. the sum over the length of each edge.

9.A. **Periodic graphs.** Suppose we have an infinite graph X_0 on which a discrete, finitely generated group Γ operates such that the quotient $M_0 := X_0/\Gamma$ is a finite graph. In the same way as in the previous sections, we can associate a family of graph-like compact manifolds M_ε to the graph M_0. By a lifting procedure we obtain a (non-compact) covering manifold X_ε of M_ε with deck transformation group Γ, i.e., M_ε is isometric to X_ε/Γ. Furthermore, X_ε is a graph-like manifold collapsing to the infinite graph X_0.

We are interested in spectral properties of the non-compact manifolds X_ε. Assuming that Γ is abelian, we can apply Floquet theory (a non-commutative version is work in progress, cf. [LP03]). Instead of investigating Δ_{X_ε} we analyze a family of operators $\Delta^\theta_{M_\varepsilon}$, $\theta \in \hat{\Gamma}$, where $\hat{\Gamma}$ is the dual group, i.e., the group of homomorphisms from Γ into the unit circle $\mathbb{T}^1$. The operator $\Delta^\theta_{M_\varepsilon}$ acts on a complex line bundle over the compact manifold M_ε, or equivalently, over the closure of a fundamental domain $D_\varepsilon \subset X_\varepsilon$ with θ-periodic boundary conditions.

We call the closure $\overline{D}_\varepsilon$ a *period cell* and denote it also by M_ε (for details see e.g. [RS-4] or [P03]). The direct integral decomposition implies

$$\operatorname{spec}\Delta_{X_\varepsilon} = \bigcup_{k\in\mathbb{N}} B_k(\varepsilon), \qquad B_k(\varepsilon) := \{\, \lambda_k^\theta(M_\varepsilon) \,|\, \theta\in\hat{\Gamma} \,\}$$

where $B_k(\varepsilon)$ is a compact subset of $[0,\infty)$, called the *k-th band*.[5] A similar assertion holds for the limit operator on X_ε.

9.B. **Spectral gaps.** We are interested in the existence of *spectral gaps* of the operator Δ_{X_ε}, i.e., the existence of an interval $[a,b]$, $0<a<b$, such that $\operatorname{spec}\Delta_{X_0}\cap[a,b]=\emptyset$. Note that $\operatorname{spec}\Delta_{X_\varepsilon}$ is purely essential.

Theorem 9.2. *We have $\lambda_k^\theta(M_\varepsilon)\to\lambda_k^\theta(Q_0)$ for $\varepsilon\to 0$* uniformly *in $\theta\in\hat{\Gamma}$. Furthermore,*

$$B_k(\varepsilon)\cap B_{k+1}(\varepsilon)=\emptyset \qquad \text{if} \qquad B_k(0)\cap B_{k+1}(0)=\emptyset$$

provided ε is small enough. In particular, an arbitrary (but finite) number of gaps open up in the spectrum of Δ_{X_ε} provided the limit operator Q_0 has enough gaps and ε is small enough.

Proof. The spectral convergence can be proven in the same way as in the previous sections. Note that the error terms converge *uniformly* in $\theta\in\hat{\Gamma}$ since all error bounds are independent of θ. The only point where θ enters is the error estimate (2.11) for the lower eigenvalue estimate. In this case, we argue as follows: we have $\lambda_k^\theta(M_\varepsilon)\le\lambda_k^{\mathrm{D}}(M_\varepsilon)$, i.e., the Dirichlet Laplacian eigenvalues form an upper bound on the θ-periodic eigenvalues. Here, we pose Dirichlet boundary conditions on the boundary of the period cell. Furthermore, $\lambda_k^{\mathrm{D}}(M_\varepsilon)\to\lambda_k^{\mathrm{D}}(M_0)$ by the same arguments as in the previous sections. Therefore, we can choose $\lambda_k=\lambda_k^\theta(M_\varepsilon)\le\lambda_k^{\mathrm{D}}(M_\varepsilon)\le 2\lambda_k^{\mathrm{D}}(M_0)$ in (2.11) *independently* of θ. □

Note that we cannot expect to show the existence of *infinitely many* gaps in X_ε even if $\operatorname{spec}\Delta_{X_0}$ has infinitely many gaps since the convergence is not uniform in k (cf. Remark 9.1). This is related to the deep open problem about the validity of Bohr-Sommerfeld conjecture on such periodic manifolds.

Remark 9.3. If two neighboured bands $B_k(0)$ and $B_{k+1}(0)$ *overlap*, i.e., intersect in a set of positive length, the same is true for $B_k(\varepsilon)$ and $B_{k+1}(\varepsilon)$ provided ε is small enough. In contrast, if the bands intersect only in *one* point, i.e., if they *touch* each other, we cannot say anything about the (non-)existence of gaps in the spectrum of Δ_{X_ε}.

[5]Note that $\hat{\Gamma}$ is connected iff Γ is torsion free, e.g., if $\Gamma=\mathbb{Z}\times\mathbb{Z}_2$ then $\hat{\Gamma}\cong\mathbb{T}^1\times\mathbb{Z}_2$ which is homeomorphic to two disjoint copies of the unit circle $\mathbb{T}^1$. Therefore, the bands $B_k(\varepsilon)$ being the continuous image of $\hat{\Gamma}$ under the map $\theta\mapsto\lambda_k^\theta(M_\varepsilon)$ need not to be intervals.

9.C. **Decoupling limit operators.** Suppose that our graph-like periodic manifold X_ε is constructed as in Sections 6 or 8. In this case, the limit operator is a direct sum of the limit operator on the quotient M_0 since the limit operator *decouples*. Therefore, the bands $B_k(0)$ degenerate to the points $\lambda_k(Q_0)$ where Q_0 is given as in Sections 6 or 8 and the limit operator on X_0 has infinitely many gaps. This means, in particular, that the limit spectrum is not absolutely continuous, while those of the approximating operators may be. Furthermore, Theorem 9.2 applies in this case.

9.D. **Cayley graphs and Kirchhoff boundary conditions.** In the following, we want to construct examples of graph-like manifolds with fast decaying vertex volume as constructed in Section 5 such that Δ_{X_ε} has spectral gaps. In this case, the limit operator is the Laplacian Δ_{X_0} on the graph X_0 with Kirchhoff boundary conditions as in (2.3). We want to calculate the spectrum of Δ_{X_0} for certain graphs X_0. For simplicity, we assume that $p_j \equiv 1$.

Suppose that Γ is an abelian, finitely generated discrete group. Therefore,

$$\Gamma \cong \mathbb{Z}^{r_0} \times \mathbb{Z}_{p_1}^{r_1} \times \cdots \times \mathbb{Z}_{p_a}^{r_a}$$

where $\mathbb{Z}_p$ is the cyclic group of order p. Furthermore, $r_0 > 0$ since X_0 is non-compact and X_0/Γ is compact. Denote $r := r_0 + r_1 + \cdots + r_a$.

We assume that X_0 is the (metric) Cayley graph accociated to Γ w.r.t. the canonical generators $\varepsilon_1, \ldots, \varepsilon_r$ (ε_j equals 1 at the j-th component and 0 otherwise), i.e., the set of vertices is Γ and two vertices γ_1, γ_2 are connected iff $\gamma_2 = \varepsilon_j \gamma_1$ for some $1 \le j \le r$ (see Figure 5). For simplicity, we assume that each edge has length 1. Note that X_0 is $2r$-regular, i.e., each vertex meets $2r$ edges. We want to



FIGURE 5. The Cayley graph associated to the group $\Gamma = \mathbb{Z} \times \mathbb{Z}_2$ and the corresponding period cell. Note that Δ_{X_0} has no spectral gaps

calculate the eigenfunctions and eigenvalues of the θ-periodic operator $\Delta_{M_0}^\theta$, i.e., functions u_j on $I_j \cong [0,1]$ satisfying $-u_j'' = \lambda u_j$ with the boundary conditions

$$u_j(0) = u(0), \quad \mathrm{e}^{-\mathrm{i}\theta_j} u_j(1) = u(0) \quad \text{and} \quad \sum_{k=1}^{r} \bigl(\mathrm{e}^{-\mathrm{i}\theta_k} u_k'(1) - u_k'(0)\bigr) = 0 \qquad (9.1)$$

for all $j = 1, \ldots, r$. Here, $\theta \in \mathbb{T}^{r_0} \times \mathbb{T}_{p_1}^{r_1} \times \cdots \times \mathbb{T}_{p_a}^{r_a}$ where $\mathbb{T}_p := \{\, \xi \in \mathbb{R}/\mathbb{Z} \,|\, \mathrm{e}^{\mathrm{i}\xi p} = 1 \,\}$ is the group of p-th unit roots (isomorphic to $\mathbb{Z}_p$). Note that we have identified $\theta \in \mathbb{T}^r$ with $\gamma \mapsto \mathrm{e}^{\mathrm{i}\theta\cdot\gamma} \in \hat{\Gamma}$.

If $\lambda = \omega^2 > 0$ (and $\omega > 0$) we make the Ansatz

$$u_j(x) := Z\cos(\omega x) + A_j \sin(\omega x)$$

and arrive at the coefficient matrix $M(\omega)$ given by

$$\begin{pmatrix} \mathrm{e}^{-\mathrm{i}\theta_1}\sin\omega & 0 & \cdots & 0 & (\mathrm{e}^{-\mathrm{i}\theta_1}\cos\omega - 1) \\ 0 & \mathrm{e}^{-\mathrm{i}\theta_2}\sin\omega & \cdots & 0 & (\mathrm{e}^{-\mathrm{i}\theta_2}\cos\omega - 1) \\ \vdots & & & 0 & \vdots \\ 0 & 0 & \cdots & \mathrm{e}^{-\mathrm{i}\theta_r}\sin\omega & (\mathrm{e}^{-\mathrm{i}\theta_r}\cos\omega - 1) \\ (\mathrm{e}^{-\mathrm{i}\theta_1}\cos\omega - 1) & (\mathrm{e}^{-\mathrm{i}\theta_2}\cos\omega - 1) & \cdots & (\mathrm{e}^{-\mathrm{i}\theta_r}\cos\omega - 1) & (-\sin\omega \sum_{k=1}^{r} \mathrm{e}^{-\mathrm{i}\theta_k}) \end{pmatrix}$$

for the variables $A_1, \ldots, A_r, Z$. A direct calculation shows that

$$\det M(\omega) = 2(\sin^{r-1}\omega)\mathrm{e}^{-\mathrm{i}\sum_k \theta_k} \sum_k (\cos\omega - \cos\theta_k).$$

Non-trivial solutions of the eigenvalue problem exist iff $\det M(\omega) = 0$, i.e. if $\omega = \ell\pi$, $\ell \in \mathbb{N}$, or

$$\cos\omega = \frac{1}{r}\sum_{k=1}^{r} \cos\theta_k. \tag{9.2}$$

The solutions $\omega = \ell\pi$ correspond to Dirichlet eigenfunctions on each edge and produce therefore bands degenerated to a point $\{(\ell\pi)^2\}$. The multiplicity is $r-1$ provided $\theta \neq 0$ (if ℓ is even) resp. $\theta \neq \pi$ (if ℓ is odd) and $r+1$ if $\theta = 0$ resp. $\theta = \pi$ (modolo 2π). If $\omega \neq \ell\pi$, the eigenvalues are simple. Note that the bands at $\omega^2 = (\ell\pi)^2$ do not overlap, but *touch* each other.

For $\omega = 0$, we need a special Ansatz. The only possibility is the case of periodic boundary conditions ($\theta = 0$); the eigenvalue is simple.

Theorem 9.4. *If one of the orders $p_1, \ldots, p_a$ is odd, the operator Δ_{X_0} has infinitely many spectral gaps. In particular, Theorem 9.2 applies.*

If all orders $p_1, \ldots, p_a$ are even, we have $\operatorname{spec}\Delta_{X_0} = [0, \infty)$.

Proof. We analyze the behaviour of ω in dependence of the continuous parameters $\theta_1, \ldots, \theta_{r_0} \in \mathbb{T}^{r_0}$ given by the relation (9.2). We have gaps iff $\frac{1}{r}\sum_{k=1}^{r}\cos\theta_k$ in (9.2) does not cover the whole interval $[-1, 1]$. We reach the maximal value 1 iff all $\theta_j = 0$ ($j = 1, \ldots, r$) and the minimal value -1 iff all $\theta_j = \pi$ ($j = 1, \ldots, r$). The latter can only occur if all group orders are even. Note that in this case, the whole interval $[-1, 1]$ can be covered by an appropriate choice of the θ_j's, $j = r_0 + 1, \ldots, r$. □

We cannot say anything about the (non-)existence of gaps in the case when all orders $p_1, \ldots, p_a$ are even. If e.g. $\Gamma = \mathbb{Z} \times \mathbb{Z}_2$, the bands do not overlap, but touch each other and fill the whole half line $[0, \infty)$ (cf. Remark 9.3).



FIGURE 6. The Cayley graph associated to the group $\Gamma = \mathbb{Z}\times\mathbb{Z}_2\times\mathbb{Z}_1$, where the trivial group $\mathbb{Z}_1$ leads to the attachment of a loop at each vertex. On the right, the corresponding period cell is shown. Note that Δ_{X_0} has spectral gaps in contrast to the example without loops.

9.E. **Cayley graphs with loops.** If we set one (or more) of the group orders p_j equal to 1 we formally attach a loop (or more) at each vertex (see Figure 6).

Theorem 9.5. *The Laplacian of a Cayley graph associated to an arbitrary finitely generated abelian discrete group* Γ *has spectral gaps provided we attach at each vertex a fixed number of loops.*

Proof. Formally, the assertion follows from Theorem 9.4. Note that $\hat{\mathbb{Z}}_1 = \{0\}$, i.e., the corresponding component of θ cannot be π and therefore, the minimum -1 cannot be achieved in (9.2). □

This is an analogue of gap generation by decoration of the graph as discussed by Aizenman and Schenker in [AS00].

9.F. **Cayley graphs and the borderline case.** In the borderline case, the eigenvalue problem of the limit operator is more complicated. Here, functions u_j on $I_j \cong [0,1]$ satisfy $-u_j'' = \lambda u_j$ with the boundary conditions

$$u_j(0) = u(0), \quad \mathrm{e}^{-\mathrm{i}\theta_j} u_j(1) = u(0) \quad \text{and} \quad \sum_{k=1}^{r} \big(\mathrm{e}^{-\mathrm{i}\theta_k} u_k'(1) - u_k'(0)\big) = c\lambda u(0) \tag{9.3}$$

for all $j = 1, \dots, r$ where c^2 is the volume of the (unscaled) vertex neighbourhood (cf. (7.3)). Again, we make the Ansatz

$$u_j(x) := Z\cos(\omega x) + A_j \sin(\omega x)$$

with $\lambda = \omega^2 > 0$ and arrive at the coefficient matrix $M_c(\omega)$ given by

$$\begin{pmatrix} \mathrm{e}^{-\mathrm{i}\theta_1}\sin\omega & 0 & \cdots & 0 & (\mathrm{e}^{-\mathrm{i}\theta_1}\cos\omega - 1) \\ 0 & \mathrm{e}^{-\mathrm{i}\theta_2}\sin\omega & \cdots & 0 & (\mathrm{e}^{-\mathrm{i}\theta_2}\cos\omega - 1) \\ \vdots & & & 0 & \vdots \\ 0 & 0 & \cdots & \mathrm{e}^{-\mathrm{i}\theta_r}\sin\omega & (\mathrm{e}^{-\mathrm{i}\theta_r}\cos\omega - 1) \\ (\mathrm{e}^{-\mathrm{i}\theta_1}\cos\omega - 1) & (\mathrm{e}^{-\mathrm{i}\theta_2}\cos\omega - 1) & \cdots & (\mathrm{e}^{-\mathrm{i}\theta_r}\cos\omega - 1) & -(\sin\omega \sum_{k=1}^{r} \mathrm{e}^{-\mathrm{i}\theta_k} + c\omega) \end{pmatrix}$$

for the variables $A_1, \dots, A_r, Z$. Note that formally the case $c = 0$ corresponds to the Kirchhoff boundary condition case and $M_0(\omega) = M(\omega)$. In a similar way as

before we have

$$\det M(\omega) = (\sin^{r-1}\omega)\mathrm{e}^{-\mathrm{i}\sum_k \theta_k}\big(2\sum_k(\cos\omega - \cos\theta_k) - c\omega\sin\omega\big).$$

Non-trivial solutions of the eigenvalue problem exist iff $\det M(\omega) = 0$, i.e. if $\omega = \ell\pi$, $\ell \in \mathbb{N}$, or

$$\cos\omega - \frac{1}{2rc} = \frac{1}{r}\sum_{k=1}^{r}\cos\theta_k. \tag{9.4}$$

Again, the solutions $\omega = \ell\pi$ correspond to Dirichlet eigenfunctions on each edge and produce therefore bands degenerated to a point $\{(\ell\pi)^2\}$.

The situation in the case $c > 0$ is more complicated since the relation (9.4) is no longer 2π-periodic in ω. In such a case the spectrum could be more complicated; recall the example of a lattice graph discussed in [EG96] shows where number-theoretic properties of parameters play role. This interesting question will be considered separately.

Acknowledgements

The authors appreciate P. Kuchment who made available to them the paper [KuZ03] prior to publication. O.P. is grateful for the hospitality extended to him at the Nuclear Physics Institute of Czech Academy of Sciences where a part of this work was done. The research was partially supported by the GAAS grant A1048101 and the SPECT program of the European Science Foundation.

References

[AC95] C. Anné and B. Colbois, *Spectre du Laplacien agissant sur les p-formes différentielles et écrasement d'anses*, Math. Ann. **303** (1995), no. 3, 545–573.

[AMSC01] A. N. Andriotis, M. Menon, D. Srivastava, and L. Chernozatonskii, *Rectification properties of carbon nanotube "Y–junctions"*, Phys. Rev. Lett. **87** (2001), 066802.

[AS00] M. Aizenman and J. H. Schenker, *The creation of spectral gaps by graph decoration*, Lett. Math. Phys. **53** (2000), 253–262.

[CdV86] Y. Colin de Verdière, *Sur la multiplicité de la première valeur propre non nulle du laplacien*, Comment. Math. Helv. **61** (1986), no. 2, 254–270. MR **88b:**58140

[Cha84] I. Chavel, *Eigenvalues in Riemannian geometry*, Academic Press, Orlando, 1984.

[Dav96] E. B. Davies, *Spectral theory and differential operators*, Cambridge University Press, Cambridge, 1996.

[EG96] P. Exner and R. Gawlista, *Band spectra of rectangular graph superlattices*, Phys. Rev. B **53** (1996), no. 11, 7275–7286.

[EŠ89] P. Exner and P. Šeba, *Bound states in curved quantum waveguides*, J. Math. Phys. **30** (1989), no. 11, 2574–2580.

[JM92] Sh. Jimbo and Y. Morita, *Remarks on the behavior of certain eigenvalues on a singularly perturbed domain with several thin channels*, Commun. Partial Differ. Equations **17** (1992), 523–552.

[K66] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1966.

[KoS99] V. Kostrykin and R. Schrader, *Kirchhoff's rule for quantum wires*, J. Phys. A **32** (1999), no. 4, 595–630.

[Ku02] P. Kuchment, *Graph models for waves in thin structures*, Waves Random Media **12** (2002), no. 4, R1–R24.

[KuZ01] P. Kuchment and H. Zeng, *Convergence of spectra of mesoscopic systems collapsing onto a graph*, J. Math. Anal. Appl. **258** (2001), no. 2, 671–700.

[KuZ03] ———, *Asymptotics of spectra of Neumann Laplacians in thin domains*, Advances in differential equations and mathematical physics (Birmingham, AL, 2002), Contemp. Math., vol. 327, Amer. Math. Soc., Providence, RI, 2003, pp. 199–213. MR 1 991 542

[LP03] F. Lledó and O. Post, *Non-commutative Floquet theory and spectral gaps*, Preprint (2003).

[P00] O. Post, *Periodic manifolds, spectral gaps, and eigenvalues in gaps*, Ph.D. thesis, Technische Universität Braunschweig, 2000.

[P03] ———, *Periodic manifolds with spectral gaps*, J. Diff. Equations **187** (2003), 23–45.

[PRL+00] C. Papadopoulos, A. Rakitin, J. Li, V. S. Vedeneev, and J. M. Xu, *Electronic properties in Y–junction carbon nanotubes*, Phys. Rev. Lett. **85** (2000), 3476–3479.

[Ro84] J.-P. Roth, *Le spectre du laplacien sur un graphe*, Théorie du potentiel (Orsay, 1983), Lecture Notes in Math., vol. 1096, Springer, Berlin, 1984, pp. 521–539.

[RueS53] K. Ruedenberg and C. W. Scherr, *Free–electron network model for conjugated systems, I. Theory*, J. Chem. Phys. **21** (1953), 1565–1581.

[RS-1] M. Reed and B. Simon, *Methods of modern mathematical physics I: Functional analysis*, Academic Press, New York, 1980.

[RS-4] ———, *Methods of modern mathematical physics IV: Analysis of operators*, Academic Press, New York, 1978.

[RubS01] J. Rubinstein and M. Schatzman, *Variational problems on multiply connected thin strips. I. Basic estimates and convergence of the Laplacian spectrum*, Arch. Ration. Mech. Anal. **160** (2001), no. 4, 271–308. MR 1 869 667

[Sa00] Y Saito, *The limiting equation for Neumann Laplacians on shrinking domains.*, Electron. J. Differ. Equ. **31** (2000), 25 p., `http://www.emis.de/journals/EJDE/Volumes/2000/31/abstr.html`.

[Tay96] M. E. Taylor, *Partial differential equations. Basic theory*, Springer-Verlag, New York, 1996.

[TBG+02] M. Terrones, F. Banhart, N. Grobert, J.-C. Charlier, H. Terrones, and P. M. Ajayan, *Molecular junctions by joining single-walled carbon nanotubes*, Phys. Rev. Lett. **89** (2002), 075505.

Department of Theoretical Physics, NPI, Academy of Sciences, 25068 Řež near Prague, and Doppler Institute, Czech Technical University, Břehová 7, 11519 Prague, Czechia

Institut für Reine und Angewandte Mathematik, Rheinisch-Westfälische Technische Hochschule Aachen, Templergraben 55, 52062 Aachen, Germany