Effective Dynamics of a Magnetic Vortex in a Local potential

Y. Strauss [a) c)] and I.M. Sigal [a) b)]

a) Department of Mathematics, University of Toronto,
Toronto, ON, Canada, M5S 3G3. URL: www.math.toronto.edu/sigal

b) Department of Mathematics, University of Notre Dame,
Notre Dame, IN, USA, 46556-4618.

c) Einstein Institute of Mathematics, The Hebrew University of Jerusalem,
Jerusalem, Israel 91904. E-mail: ystrauss@math.huji.ac.il

Abstract

We study simplified mean field model of superconductor dynamics in the presence of impurities or for variable superconductor depth. This model is given by the gradient-flow version of the Ginzburg-Landau equations (Grokov-Eliashberg equations) with an addition of a potential term. We find a dynamical law of motion of the vortex center, involving the potential, such that for datum close to a (static) magnetic vortex the solution is close, for all times, to a magnetic vortex whose center obeys this law.

1 Introduction

1.1 Preliminary discussion and statement of the problem

In this work we consider the gradient flow version of the Ginzburg-Landau equations in the presence of a local potential. Specifically, we study certain solutions of the system of equations

$$\begin{aligned} \partial_t \psi &= \Delta_A \psi + \frac{\lambda}{2}(1-|\psi|^2)\psi - \epsilon W \psi \\ \partial_t A &= -\nabla \times \nabla \times A + Im(\overline{\psi}\nabla_A \psi) \end{aligned} \tag{1.1}$$

where $(\psi(t), A(t)) : \mathbb{R}^2 \mapsto \mathbb{C} \times \mathbb{R}^2$ for each $t \geq 0$, $\nabla_A = \nabla - iA$ and $\Delta_A = \nabla_A^2$. The function $W : \mathbb{R}^2 \mapsto \mathbb{R}$ is a localized potential and is assumed

to be smooth and sufficiently fast decaying at infinity. The parameter ϵ measures the strength of the potential. In the study of superconductivity the function ψ is the order parameter with $|\psi|$ measuring the density of superconducting Cooper pairs and the vector field A is the vector potential for the magnetic field. The potential $W(x)$ arises with impurities, variations of superconductor geometry or other inhomogeneities in the matter.

Setting $\epsilon = 0$ in Eq. (1.1), one obtains the following set of coupled equations for ψ and A

$$\begin{aligned} \partial_t \psi &= \Delta_A \psi + \frac{\lambda}{2}(1 - |\psi|^2)\psi \\ \partial_t A &= -\nabla \times \nabla \times A + \mathrm{Im}\,(\overline{\psi}\nabla_A \psi)\,. \end{aligned} \tag{1.2}$$

This system of equations is used to describe the dynamics of superconductors in the mean field approximation and near the normal-superconducting state transition point. They are called the *Gorkov-Eliashberg* equations or *time-dependent Ginzburg-Landau* (TDGL) equations [GE, T].

The global well–posedness of the TDGL equations was shown by Demoulini and Stuart [DS]. Their proof can easily be extended to Eq. (1.1). This will be outlined elsewhere.

The stationary version of the TDGL equations is the Ginzburg-Landau (GL) equations

$$\begin{aligned} \Delta_A \psi + \frac{\lambda}{2}(1 - |\psi|^2)\psi &= 0 \\ -\nabla \times \nabla \times A + Im(\overline{\psi}\nabla_A \psi) &= 0\,. \end{aligned} \tag{1.3}$$

Let $u = (\psi, A)$. Denote by $\mathcal{E}_{GL}(u)$ the Ginzburg-Landau energy functional

$$\mathcal{E}_{GL}(u) = \frac{1}{2}\int_{\mathbb{R}^2} \left[|\nabla_A \psi|^2 + |\nabla \times A|^2 + \frac{\lambda}{4}(|\psi|^2 - 1)^2 \right]. \tag{1.4}$$

The Ginzburg-Landau equations, Eq. (1.3) are the Euler-Lagrange equations derived from $\mathcal{E}_{GL}$.

Superconductors are characterized by two length scales called *penetration depth* and *coherence depth*. In the scaling units corresponding to the form of $\mathcal{E}_{GL}$ given in Eq. (1.4) the penetration depth, measuring the scale of variations in the magnetic field, is equal to 1. The coherence length, measuring the scale of variations in the order parameter ψ is $1/m_\lambda$ where $m_\lambda = \min(\sqrt{\lambda}, 2)$. For $\lambda = 1$ the two length scales are equal and we distinguish between type I superconductors, for which $\lambda < 1$ and type II superconductors for which $\lambda > 1$. Experimentally, type I and type II materials differ

in their magnetic behavior. In type I superconductors magnetic fields are excluded from the bulk of the material except for a very thin layer near the surface. In type II superconductors magnetic fields penetrate the material in vortex structures. In general, Type II superconductors can sustain magnetic fields much higher than type I superconductors without losing their superconducting state. The existence of magnetic vortices and of type II superconductors was predicted in 1957 by Abrikosov [A](in addition Abrikosov predicted the existence of large arrays of magnetic vortices, called Abrikosov lattices, in type II materials).

Configurations (ψ, A) with finite energy are classified by the Brouwer degree of ψ, i.e., the topological degree

$$deg(\psi) := deg\left(\left.\frac{\psi}{|\psi|}\right|_{|x|=R}\right), \qquad deg(\psi) \in \mathbb{Z}$$

where R is large enough. For the magnetic field $B = \nabla \times A$ the degree, $deg(\psi)$, is related to the magnetic flux as

$$\int_{\mathbb{R}^2} B = 2\pi \deg(\psi) \in \mathbb{Z}$$

(quantization of magnetic flux).

Magnetic (Abrikosov) vortices are equivariant solutions of Eq. (1.3) of the form

$$\psi^{(n)}(x) = f_n(r)e^{in\theta}, \qquad A^{(n)}(x) = a_n(r)\nabla(n\theta) \tag{1.5}$$

where (r, θ) are the polar coordinates of $x \in \mathbb{R}^2$ and $n \in \mathbb{Z}$. Calculation of the degree of $\psi^{(n)}$ for a magnetic vortex solution given by Eq. (1.5) yields $deg(\psi^{(n)}) = n$. The pair $(\psi^{(n)}, A^{(n)})$ is called an n-vortex. The existence of n-vortices was proved by Plohr [P] and by Berger and Chen [BC] by variational methods. Stability of n-vortices was proved by Gustafson and Sigal [GS1] and Gustafson [G2]. It is shown in [JT] that $0 < f_n < 1$, $0 < a_n < 1$ and that the asymptotic behavior for $a_n(r)$ and $f_n(r)$ as $r \to \infty$ is given by

$$f_n(r) = 1 + O(e^{-m_\lambda r}), \qquad a_n(r) = 1 + O(e^{-r})\,.$$

It is also known that $f_n(r)$ vanishes like r^n and $a_n(r)$ vanishes like r^2 as r goes to the origin.

For an n-vortex $v^{(n)} = (\psi^{(n)}, A^{(n)})$ the GL energy functional $\mathcal{E}_{GL}$ is a smooth functional on the affine space given by

$$X^{(n)} \equiv \left\{ u : \mathbb{R}^2 \to \mathbb{C} \times \mathbb{R}^2 \mid u - v^{(n)} \in H^1(\mathbb{R}^2; \mathbb{C} \times \mathbb{R}^2) \right\}. \tag{1.6}$$

As mentioned above, type II superconductors can sustain very large magnetic fields (over 10^5 Gauss). However, a major obstacle in the attempt to produce large magnetic fields is the dissipation of energy due to the *creeping* or *flow* of vortices [T]. One way to overcome this problem is to *pin down* the vortices to particular locations in the material. The pinning down of vortices is achieved by the presence of point defects, impurities, inhomogeneities or by a variation in the thickness of the sample of superconducting material [DG]. Rubinstein [Rub1] considered a model of the effect of pinning in which the quartic term in $\mathcal{E}_{GL}$ is replaced by the modified term

$$\frac{\lambda}{2}(|\psi|^2 - f^2(x))^2 \,.$$

The function $f(x)$ can be thought of as measuring the quality of the superconductor (consider for example the large λ limit).

In [BBH] Bethuel, Brezis and Hélein study the large λ limit of a simplified version of the GL functional without magnetic field:

$$\mathcal{E}_\epsilon(\psi) = \int_\Omega \left[\frac{1}{2}|\nabla\psi|^2 + \frac{1}{4\epsilon^2}(|\psi|^2 - 1)^2\right] ,$$

where Ω is a simply connected bounded domain in $\mathbb{R}^2$. Since there is no magnetic field in the problem one needs to introduce a mechanism for the confinment of vortices to Ω. The minimization problem for $\mathcal{E}_\epsilon(\psi)$ is considered, therefore, over all ψ satisfying the Dirichlet boundary condition

$$\psi(x) = g(x), \qquad x \in \partial\Omega$$

where $g : \partial\Omega \to S^1$ satisfies $deg\, g = d \neq 0$. In the general context of the Bethuel-Brezis-Hélein theory Andre and Shafrir [AS] analyzed a model for the pinning of vortices in superconductors with small variable thickness which was proposed by Du and Gunzburger [DG]. This model for the pinning effect involves a weighted GL functional of the form

$$\mathcal{E}_\epsilon(\psi) = \int_\Omega p(x)\left[\frac{1}{2}|\nabla\psi|^2 + \frac{1}{4\epsilon^2}(|\psi|^2 - 1)^2\right] .$$

For this model Andre and Shafrir proved the Bethuel-Brezis-Hélein type result that the set of points $a_1, \ldots, a_d$, on which the limiting ψ_ϵ is concentrated, consists of points (possibly repeating) in which p achieves its global minimum.

More recently the effect of pinning was studied by Aftalion, Sandier and Serfaty [ASaSe], Andre, Bauman and Philips [ABP] and I.M. Sigal and F.

Ting [ST1]. In addition, Sigal and Ting investigated the stability of pinned vortices [ST2].

Our goal in the present paper is to provide a description of the dynamics of the pinning effect, i.e., we aim to derive rigorously the law governing the dynamics of a vortex in the presence of a potential. Formally speaking, this constitutes a reduction of the dynamics of the various fields in the problem to an effective dynamics of a smaller number of degrees of freedom, i.e, the vortex center and the gauge function. The methods utilized for the proof of the main theorem in the current work are close to those used by Gustafson and Sigal in their study of the dynamics of magnetic vortices [GS2]. We refer to [GS2] for a short survey of various rigorous and non-rigorous results concerning the dynamics of vortices in various time dependent versions of the GL equations. In particular, we mention here the non-rigorous results for the superconducting model of Eq. (1.2) obtained by Perez and Rubinstein [PeRu] and W. E [E] and the rigorous results obtained, for the same equations, by Gustafson and Sigal [GS2]. A simplified version of the TDGL equations, called the *nonlinear heat flow* (NLHF) equation, in which $A \equiv 0$, was studied by Neu [N], Lin [L], Jerard and Soner [JS] and Rubinstein and Sternberg [RS] (see also [Rub2] and references therein).

We conclude this subsection by noting that reviews of the Ginzburg-Landau theory of superconductivity can be found, for example, in [B, BFGLV, G1, JT, Rit, Riv, Rub2].

1.2 Symmetries of the Ginzburg-Landau equations

The Ginzburg-Landau energy functional $\mathcal{E}_{GL}$ is invariant under the following symmetries:

(i) Translation symmetry $x \to x' = x + z$

$$\psi(x) \to \psi_z(x) = \psi(x - z), \qquad A(x) \to A_z(x) = A(x - z)\,;$$

(ii) Gauge symmetry

$$\psi(x) \to \psi_\gamma(x) = e^{i\gamma(x)}\psi(x), \qquad A(x) \to A_\gamma(x) = A(x) + \nabla\gamma(x)\,;$$

(iii) Rotation Symmetry $x \to x' = Rx$, $R \in SO(2)$

$$\psi(x) \to \psi_R(x) = \psi(R^{-1}x), \qquad A(x) \to A_R(x) = RA(R^{-1}x)\,.$$

We note that for an n-vortex solution the application of the rotation symmetry group correspond to a subgroup of the gauge symmetry. Therefore, for n-vortices the relevant symmetry operations are the translation and gauge groups. Accordingly, given an n-vortex $v^{(n)} = (\psi^{(n)}, A^{(n)})$, we can apply translation symmetry with translation parameters $\mathbf{z} = (z_1, z_2) \in \mathbb{R}^2$ and gauge symmetry with a gauge function γ and find another solution $v_{\mathbf{z}\gamma}$ of Eq. (1.3) where

$$v_{\mathbf{z}\gamma} = (\psi_{\mathbf{z}\gamma},\, A_{\mathbf{z}\gamma}) = (e^{i\gamma}\psi^{(n)}(\cdot - \mathbf{z}), A^{(n)}(\cdot - \mathbf{z}) + \nabla\gamma) \tag{1.7}$$

with $\psi_{\mathbf{z}\gamma} \equiv e^{i\gamma}\psi^{(n)}(\cdot - \mathbf{z})$ and $A_{\mathbf{z}\gamma} \equiv A^{(n)}(\cdot - \mathbf{z}) + \nabla\gamma$. The latter functions can be also written as $\psi_{\mathbf{z}\gamma} = e^{i\gamma}\psi_{\mathbf{z}0}$ and $A_{\mathbf{z}\gamma} = A_{\mathbf{z}0} + \nabla\gamma$, where $\psi_{\mathbf{z}0} \equiv \psi^{(n)}(\cdot - \mathbf{z})$ and $A_{\mathbf{z}0} \equiv A^{(n)}(\cdot - \mathbf{z})$. For reasons that will become clear below we consider gauge transforms of the form

$$\gamma = \sum_{i=1,2} z_i A_i^{(n)}(\cdot - \mathbf{z}) + \tilde{\gamma} \tag{1.8}\,.$$

Then, according to Eq. (1.6) and Eq. (1.7) we require that $\tilde{\gamma} \in H^2(\mathbb{R}^2; \mathbb{R})$. In this case $v_{\mathbf{z}\gamma}$ in Eq. (1.7) will always be in $X^{(n)}$. Applying symmetry transformations in the form of Eq. (1.7) for all possible values of $\mathbf{z}$ and γ we obtain the symmetry manifold of the n-vortex

$$M_{sym}^{(n)} = \left\{ v_{\mathbf{z}\gamma} \mid \mathbf{z} \in \mathbb{R}^2, \tilde{\gamma} \in H^2(\mathbb{R}^2; \mathbb{R}) \right\} \tag{1.9}$$

with $\tilde{\gamma}$ given in Eq. (1.8). It is obvious that all points in $M_{sym}^{(n)}$ are solutions of the Ginzburg-Landau equations, Eq. (1.3).

In order to obtain Eq. (1.1) we add to $\mathcal{E}_{GL}$ an interaction term of the form

$$\mathcal{E}_{int}(u) = \int d^2x\, W(x)|\psi(x)|^2 \tag{1.10}$$

where $u \in X^{(n)}$. Define a modified energy functional

$$\mathcal{E}_\epsilon(u) = \mathcal{E}_{GL}(u) + \epsilon\, \mathcal{E}_{int}(u)\,. \tag{1.11}$$

With this definition Eq. (1.1) can be written

$$\partial_t u = -\mathcal{E}_\epsilon'(u) \tag{1.12}$$

where $\mathcal{E}_\epsilon'$ is the Fréchet derivative or, more precisely, the L^2–gradient, of $\mathcal{E}_\epsilon$. We observe that the interaction term in Eq. (1.10) does not break the gauge symmetry. However, $\mathcal{E}_{int}$ does break the translation symmetry and if

we translate ψ the interaction energy depends on the translation parameter $\mathbf{z}$. In particular, applying translations to the n-vortex solution $v^{(n)}$ we define the interaction energy

$$W_{int}^{(n)}(\mathbf{z}) \equiv \epsilon \int d^2x\, W(\mathbf{x}) |\psi^{(n)}(\mathbf{x}-\mathbf{z})|^2 \,. \tag{1.13}$$

1.3 Summary of results

The main result in this work states that the force derived from the interaction energy $W_{int}^{(n)}(\mathbf{z})$ is governing the effective dynamics of the magnetic vortex. More specifically, given an initial data $u_0 = (\psi_0, A_0)$ close to a point $v_{\mathbf{z}_0\gamma_0} \in M_{sym}^{(n)}$, we show that there exists a curve in $M_{sym}^{(n)}$, parametrized by the time t, such that for all $t \geq 0$ the solution $u(t) = (\psi(t), A(t))$ satisfies

$$u(t) - v_{\mathbf{z}(t)\gamma(t)} = O(\epsilon)$$

where the equation for the effective dynamics of the vortex center is given by

$$a_n \dot{\mathbf{z}} + \nabla_{\mathbf{z}} W_{int}^{(n)}(\mathbf{z}) = O(\epsilon^2)$$

and

$$a_n = \frac{1}{2} \|\nabla_{A^{(n)}} \psi^{(n)}\|_2^2 + \|\nabla \times A^{(n)}\|_2^2 \,. \tag{1.14}$$

1.4 Effective dynamics of vortices - main theorem

We formulate our main result giving the effective dynamics of the gradient flow Ginzburg-Landau equations with a local potential. Recall that $v^{(n)} = (\psi^{(n)}, A^{(n)})$ denotes an n-vortex solution of the static GL equations, Eq. (1.3), and $M_{sym}^{(n)}$ denotes the symmetry manifold for $v^{(n)}$ defined in Eq. (1.9). We have:

Theorem 1 (effective dynamics) *Let n be an arbitrary integer if $0 < \lambda < 1$ and $n = \pm 1$ if $\lambda > 1$. There exist constants $\epsilon_0, c, c_0 > 0$, with c, c_0 depending only on ϵ_0, such that for $0 < \epsilon < \epsilon_0$ and for a datum u_0 satisfying the condition*

$$\|u_0 - v_{\mathbf{z}_0\gamma_0}\|_{X^{(n)}} \leq c_0 \epsilon \ ,$$

for some $v_{\mathbf{z}_0\gamma_0} \in M_{sym}^{(n)}$, the solution $u(t)$ of Eq. (1.1) with datum u_0 is of the form

$$u(t) = v_{\mathbf{z}(t)\gamma(t)} + \zeta(t)$$

with $v_{\mathbf{z}(t)\gamma(t)} \in M^{(n)}_{sym}$, *the functions* $z(t)$, $\gamma(t)$ *satisfying the differential equations*

$$\begin{aligned} a_n \frac{d\mathbf{z}}{dt} &= -\nabla_{\mathbf{z}} W^{(n)}_{int}(\mathbf{z}) + O(\epsilon^2) \\ \frac{d\gamma}{dt} &= \sum_i \dot{z}_i A^{(n)}_i(\cdot - \mathbf{z}) + O_{H^{1-s}}(\epsilon^2), \qquad s > 0 \end{aligned} \tag{1.15}$$

where a_n *is given by Eq. (1.14) and the function* $\zeta(t)$ *in* $X^{(n)}$ *and bounded as*

$$\|\zeta(t)\|_{X^{(n)}} \leq c\epsilon\,. \tag{1.16}$$

□

Remark: The notation $O_X(\epsilon)$ stands for a function in X whose X-norm is bounded by $C\epsilon$.

Eqns. (1.15) will be called the *effective vortex dynamics equations*. We see that the solution to Eq. (1.1) which is initially close to an n-vortex stays close to a vortex for all times provided its center and gauge evolve according to Eqns. (1.15).

The proof of Theorem 1 is given in Section 2 with several technical statements proved in Sections 3 and 4. Finally, Appendix A contains explicit expressions for the Taylor expansions of the energy functional and its Fréchet derivative (r.h.s. of the equations of motion).

2 Proof of Theorem 1

We begin with some comments concerning notation. Throughout the discussion below the symbol H^s always means the Sobolev space $H^s(\mathbb{R}^2;\mathbb{C}\times\mathbb{R}^2)$. Let $\zeta = (\xi, F) \in L^2(\mathbb{R}^2;\mathbb{C}\times\mathbb{R}^2)$ and $\eta = (\lambda, G) \in L^2(\mathbb{R}^2;\mathbb{C}\times\mathbb{R}^2)$. The L^2 inner product of ζ and η is denoted by $\langle\zeta,\eta\rangle$ and is given by

$$\langle\zeta,\eta\rangle \equiv \int_{\mathbb{R}^2} (Re(\overline{\xi}\lambda) + F\cdot G)\,.$$

In order to make the notation less cubersome we shall denote below the parameters $\{\mathbf{z},\gamma\}$ by σ. For example, according to this convention v_σ stands for $v_{\mathbf{z}\gamma}$. We restore the full notation whenever this is necessary or if it adds to the clarity of argument. Otherwise, the shorter notation is kept throughout the course of the proof. In several steps that do not require reference to the parametrization of M_{as} we omit it altogether.

2.1 The manifold of approximate solutions

Let

$$\Sigma = \{(\mathbf{z}, \gamma) \mid \mathbf{z} \in \mathbb{R}^2, \gamma - \sum_i z_i A_i^{(n)}(\cdot - \mathbf{z}) \in H^2(\mathbb{R}^2; \mathbb{R})\}\,. \tag{2.1}$$

Denote the elements of the translation group by $T_{\mathbf{z}}$ ($\mathbf{z} \in \mathbb{R}^2$) and the elements of the gauge group by G_γ ($\gamma \in H^2(\mathbb{R}^2; \mathbb{C})$). Applying the translation and gauge transformations to a vortex solution $v^{(n)}$ of Eq. (1.3) we obtain a manifold M_{as} of solutions of this equation

$$M_{as} = \{G_\gamma T_{\mathbf{z}}\, v^{(n)} \mid (\mathbf{z}, \gamma) \in \Sigma\} = \{v_{\mathbf{z}\gamma} \mid (\mathbf{z}, \gamma) \in \Sigma\} = M_{sym}^{(n)} \subset X^{(n)} \tag{2.2}$$

The manifold M_{as} is, therefore, parametrized by $\mathbf{z}$ and γ. A point in M_{as} corresponding to a given value $\{\mathbf{z}, \gamma\}$ of the parameters (for γ this means a definite gauge function) is denoted by $v_{\mathbf{z}\gamma}$.

The manifold M_{as} is called *manifold of approximate solutions*. We argue for the introduction of M_{as} and the terminology used as follows: Inserting $v_{\mathbf{z}\gamma} \in M_{as}$ as a trial solution into Eq. (1.12) we obtain in the r.h.s

$$\mathcal{E}'_\epsilon(v_{\mathbf{z}\gamma}) = \mathcal{E}'_{GL}(v_{\mathbf{z}\gamma}) + \epsilon\, \mathcal{E}'_{int}(v_{\mathbf{z}\gamma}) = \epsilon\, \mathcal{E}'_{int}(v_{\mathbf{z}\gamma}) \tag{2.3}$$

hence the r.h.s of Eq. (1.12) (or, equivalently, of Eq. (1.1)) is of order ϵ and $v_{\mathbf{z}\gamma}$ is seen to be an approximate static solution of Eq. (1.1) for ϵ small.

Denote by $T_{\mathbf{z}\gamma}\Sigma$ the tangent space to Σ at the point $(\mathbf{z}, \gamma)$ and by $T_{v_{\mathbf{z}\gamma}} M_{as}$ the tangent space to M_{as} at the point $v_{\mathbf{z}\gamma}$. A basis for $T_{\mathbf{z}\gamma}\Sigma$ is given by

$$\{\partial_{z_i} + \langle \mathbf{z} \cdot \partial_i A, \partial_{\tilde{\gamma}} \rangle, \partial_{\tilde{\gamma}}\} \tag{2.4}$$

where $\tilde{\gamma}$ is given by Eq. (1.8) and for a function g we set $\langle g, \partial_{\tilde{\gamma}} \rangle = \int d^2x\, g(x) \partial_{\tilde{\gamma}(x)}$. Using the natural parametrization map $\beta : \Sigma \to M_{as}$ defined by $\beta(\mathbf{z}, \gamma) = v_{\mathbf{z}\gamma}$ we can push forward the basis given in Eq. (2.4) in order to obtain a basis for $T_{v_{\mathbf{z}\gamma}} M_{as}$. This later basis, is obtained via the mapping of the basis in Eq. (2.4) by the Fréchet derivative $D_{\mathbf{z}\gamma}(\beta(\mathbf{z}, \gamma)) = D_{\mathbf{z}\gamma} v_{\mathbf{z}\gamma}$. Denoting $\partial_{z_i}^A = \partial_{z_i} + \langle A_i, \partial_\gamma \rangle$, we get in $T_{v_{\mathbf{z}\gamma}} M_{as}$ the basis vectors

$$T_i^{\mathbf{z}\gamma} = \partial_{z_i}^A v_{\mathbf{z}\gamma},\ i = 1, 2 \qquad G_{\delta(x)}^{\mathbf{z}\gamma} = \partial_{\gamma(x)} v_{\mathbf{z}\gamma} \tag{2.5}$$

where the (covariant) translation vectors $T_i^{\mathbf{z}\gamma}$ are given by

$$T_i^{\mathbf{z}\gamma} = \left((\nabla_{A_{\mathbf{z}\gamma}} \psi_{\mathbf{z}\gamma})_i\ , (\nabla \times A_{\mathbf{z}\gamma}) \hat{e}_i\right)\,. \tag{2.6}$$

Here $\hat{e}_1 = (0,1)$ and $\hat{e}_2 = (-1,0)$. For a definite gauge function χ the gauge basis vectors $G^{\mathbf{z}\gamma}_{\delta(x)}$ in Eq. (2.5) are defined by the relation

$$\int d^2x\, G^{\mathbf{z}\gamma}_{\delta(x)}\chi(x) = G^{\mathbf{z}\gamma}_{\chi} \tag{2.7}$$

where

$$G^{\mathbf{z}\gamma}_{\chi} = (i\chi\psi_{\mathbf{z}\gamma}, \nabla\chi)\,. \tag{2.8}$$

The vectors in Eq. (2.5) form an orthogonal basis for $T_{v_{\mathbf{z}\gamma}}M_{as}$. We have $\langle G^{\mathbf{z}\gamma}_{\delta(x)}, T^{\mathbf{z}\gamma}_i\rangle = 0$ and $\langle T^{\mathbf{z}\gamma}_i, T^{\mathbf{z}\gamma}_j\rangle = \delta_{ij}a_n$ with

$$a_n = \|T^{\mathbf{z}\gamma}_1\|_2^2 = \|T^{\mathbf{z}\gamma}_2\|_2^2 = \frac{1}{2}\|\nabla_{A^{(n)}}\psi^{(n)}\|_2^2 + \|\nabla\times A^{(n)}\|_2^2 \tag{2.9}$$

and, in addition

$$\langle G^{\mathbf{z}\gamma}_{\delta(x)}, G^{\mathbf{z}\gamma}_{\delta(y)}\rangle = K_{\mathbf{z}\gamma}(x,y) \tag{2.10}$$

where $K_{\mathbf{z}\gamma}(x,y)$ is the integral kernel for the operator

$$K_{\mathbf{z}\gamma} = -\Delta + |\psi_{\mathbf{z}\gamma}|^2\,. \tag{2.11}$$

We note that $K_{\mathbf{z}\gamma}$ is self-adjoint and that $K_{\mathbf{z}\gamma} > 0$.

We will see below that one of the main ingredients that enter into the proof of Theorem 1 is the projection of Eq. (1.1), for each time t, on the tangent space $T_{v_{\mathbf{z}\gamma}}M_{as}$ (where $\mathbf{z}$ and γ depend on t in a suitable way). Denote $P_v : T_vX^{(n)} \to T_vM_{as}$ the projection on the tangent space to M_{as} at the point $v_{\mathbf{z}\gamma}$. We are able to use the orthogonal basis of Eq. (2.5) to obtain an explicit expression for $P_{v_{\mathbf{z}\gamma}}$. For any $\zeta \in T_{v_{\mathbf{z}\gamma}}X^{(n)}$ we have

$$P_{v_{\mathbf{z}\gamma}}\zeta = \sum_{i=1,2} a_n^{-1}T^{\mathbf{z}\gamma}_i\langle T^{\mathbf{z}\gamma}_i, \zeta\rangle + \int d^2x \int d^2y\, G^{\mathbf{z}\gamma}_{\delta(x)}K^{-1}_{\mathbf{z}\gamma}(x,y)\langle G^{\mathbf{z}\gamma}_{\delta(y)}, \zeta\rangle\,. \tag{2.12}$$

In particular, if $v(t) = v_{\mathbf{z}(t)\gamma(t)}$ is a path in M_{as}, depending on the time t, then we have

$$\dot{v}_{\mathbf{z}\gamma} = P_v\dot{v}_{\mathbf{z}\gamma} = \sum_i(-\dot{z}_i)T^{\mathbf{z}\gamma}_i + G^{\mathbf{z}\gamma}_{\dot{\tilde{\gamma}}} \tag{2.13}$$

with

$$\dot{\tilde{\gamma}} = \dot{\gamma} - \sum_i \dot{z}_i A^{(n)}_i(\cdot - \mathbf{z})\,. \tag{2.14}$$

In Eq. (2.13) and Eq. (2.14) we use the notation $\dot{v}_{\mathbf{z}\gamma} = \partial_t v_{\mathbf{z}\gamma}$, $\dot{\gamma} = \partial_t\gamma$, $\dot{\tilde{\gamma}} = \partial_t\tilde{\gamma}$ and $\dot{\mathbf{z}} = \partial_t\mathbf{z}$.

2.2 Local Sobolev spaces on the manifold of approximate solutions

In this subsection we define the notion of local Sobolev spaces on the manifold of approximate solutions M_{as}. The introduction of this concept helps to facilitate many of the estimates in this work and is based on the observation that certain estimates are not uniform on M_{as} but depend on the point $v \in M_{as}$. More specifically, at the point $v_{\mathbf{z}\gamma}$ the constants appearing in the estimates depend on the gauge function γ.

As an example for the motivation for the definition of local Sobolev spaces on M_{as} we give an estimate which is used below in the proof of Lemma 2. Consider vectors $(\xi, F) \in T_{v_{\mathbf{z}\gamma}} X^{(n)}$ with the component ξ transforming covariantly under gauge transformations and F gauge invariant. Let $\zeta = (\xi, F)$ be such a vector and let $G_\gamma \zeta$ be the action of (an appropriate unitary representation of) an element G_γ of the gauge group on ζ. We have $G_\gamma \zeta = \zeta'$ where $\zeta' = (\xi', F') = (e^{i\gamma}\xi, F)$. Construct a vector $\zeta_v = (0, Im(\overline{\xi}\nabla_A \xi))$ where ξ is the first component of ζ. Then $\zeta_v \in T_{v_{\mathbf{z}\gamma}} X^{(n)}$ has the desired transformation properties under the action of the gauge group. Suppose that we want to estimate the H^{-s} $(s > 0)$ norm of ζ_v. we have

$$\|\zeta_v\|_{H^{-s}} = \|\overline{\xi}\nabla_A \xi\|_{H^{-s}} = \|\overline{\xi}\nabla\xi - iA|\xi|^2\|_{H^{-s}} \leq \|\overline{\xi}\nabla\xi\|_{H^{-s}} + \|A|\xi|^2\|_{H^{-s}} \,. \tag{2.15}$$

For the first term on the r.h.s. of Eq. (2.15) we have the estimate

$$\begin{aligned} \|\overline{\xi}\nabla\xi\|_{H^{-s}} = \sup_{\|\eta\|_{H^s}\leq 1} |\langle \eta, \overline{\xi}\nabla\xi\rangle| &\leq \sup_{\|\eta\|_{H^s}\leq 1} \|\eta\xi\|_2 \|\nabla\xi\|_2 \\ &\leq \sup_{\|\eta\|_{H^s}\leq 1} \|\eta\|_p \|\xi\|_q \|\xi\|_{H^1} \leq C\|\xi\|^2_{H^1} \,. \end{aligned} \tag{2.16}$$

where $1/2 = 1/p + 1/q$ and q is large enough so that $H^s \subset L^p$. In order to estimate the second term on the r.h.s. of Eq. (2.15) we recall that at $v_{\mathbf{z}\gamma}$ we have $A_{\mathbf{z}\gamma} = A^{(n)}(\cdot - \mathbf{z}) + \nabla\gamma$ and so

$$\|A|\xi|^2\|_{H^{-s}} = \|(A^{(n)}(\cdot - \mathbf{z}) + \nabla\gamma)|\xi|^2\|_{H^{-s}} \leq \|A^{(n)}\|_\infty \|\xi\|_2^2 + \|\nabla\gamma|\xi|^2\|_{H^{-s}} \,. \tag{2.17}$$

For the second term on the r.h.s. of Eq. (2.17) we have the estimate

$$\begin{aligned}\|\nabla\gamma|\xi|^2\|_{H^{-s}} &= \sup_{\|\eta\|_{H^s}\leq 1} |\langle \eta, (\nabla\gamma)|\xi|^2\rangle| \leq \sup_{\|\eta\|_{H^s}\leq 1} \|\eta|\xi|^2\|_2 \|\nabla\gamma\|_2 \\ &\leq \sup_{\|\eta\|_{H^s}\leq 1} \|\eta\|_p \|\xi^2\|_q \|\gamma\|_{H^1} \leq C\|\xi\|_{2q}^2 \|\gamma\|_{H^1} \leq C\|\xi\|_{H^1}^2 \|\gamma\|_{H^1}\end{aligned} \tag{2.18}$$

where again $1/2 = 1/q + 1/p$ and q is large enough so that $H^s \subset L^p$. Using Eq. (2.16)-(2.18) we obtain

$$\|\zeta_v\|_{H^{-s}} \leq C_\gamma \|\zeta_v\|_{H^1}^2 \tag{2.19}$$

where

$$C_\gamma = C(1 + \|\gamma\|_{H^1}) \tag{2.20}$$

we see that this estimate is not uniform on M_{as} but depends on the gauge function γ.

In order to simplify estimates of the type considered here we define the notion of *local Sobolev spaces* on M_{as}. With a point $v_{\mathbf{z}\gamma} \in M_{as}$, and for any real s, we associate a local Sobolev space $H^s_{v_{\mathbf{z}\gamma}}$ such that, given $\zeta = (\xi, F) \in T_{v_{\mathbf{z}\gamma}} X^{(n)}$ with the gauge transformation properties described above, its $H^s_{v_{\mathbf{z}\gamma}}$ norm is defined to be

$$\|\zeta\|_{H^s_{v_{\mathbf{z}\gamma}}} \equiv \|G_{-\gamma}\zeta\|_{H^s} \,. \tag{2.21}$$

The definition of $H^s_{v_{\mathbf{z}\gamma}}$ preserves all of the properties of Sobolev spaces. In particular, the Sobolev embedding theorems are all valid for the local Sobolev spaces. Note also that $L^p_{v_{\mathbf{z}\gamma}} = L^p$ for all p.

Suppose we want to obtain an estimate for the vector $\zeta_v = (0, Im(\overline{\xi}\nabla_A\xi))$ in the local $H^{-s}_{v_{\mathbf{z}\gamma}}$ $(s > 0)$ norm. We have

$$\|\zeta_v\|_{H^{-s}_{v_{\mathbf{z}\gamma}}} = \|\overline{\xi}\nabla_A\xi\|_{H^{-s}} = \|(\overline{e^{-i\gamma}\xi})(e^{-i\gamma}\nabla_A\xi)\|_{H^{-s}} = \|\overline{\xi'}\nabla_{A^{(n)}}\xi'\|_{H^{-s}} \tag{2.22}$$

where $\xi' = e^{-i\gamma}\xi$. Repeating the sequence of estimates in Eq. (2.16)-(2.18) for the r.h.s. of Eq. (2.22) we find that

$$\|\overline{\xi'}\nabla_{A^{(n)}}\xi'\|_{H^{-s}} \leq C\|\xi'\|_{H^1}^2 = C\|e^{-i\gamma}\xi\|_{H^1}^2 = C\|\xi\|_{H^1_{v_{\mathbf{z}\gamma}}}^2 \tag{2.23}$$

with $C = 1 + \|A^{(n)}\|_\infty$, a constant independent of γ. Thus we arrive at the (local) estimate

$$\|\zeta_v\|_{H^{-s}_{v_{\mathbf{z}\gamma}}} \leq C\|\zeta_v\|_{H^1_{v_{\mathbf{z}\gamma}}}^2 \tag{2.24}$$

which has the same form as Eq. (2.19). The emphasis is on the fact that the constant C here does not depend on γ.

In accord with the discussion here most of the estimates involved in the proof of the main theorem of this work are performed using the local Sobolev spaces $H^s_{v_{\mathbf{z}\gamma}}$.

2.3 Proof of Theorem 1

The strategy for the proof of Theorem 1 is as follows: Given $\epsilon > 0$ small enough we show that there exists a neighborhood of M_{as} such that, if the initial data u_0 for Eq. (1.1) is in this neighborhood, then the distance of the solution $u(t)$ from M_{as} can be controlled for all times $t \geq 0$. For such a solution we define in a suitable way a projection on M_{as} so as to obtain, for each $t \geq 0$ a unique point $v(t) = v(u(t)) \in M_{as}$ corresponding to $u(t)$. We think of $u(t)$ as inducing a (uniquely defined) trajectory on M_{as}, traced by $v(t)$. We then obtain effective equations of motion for the point $v(t)$ in terms of the parametrization of M_{as} by the translation parameters $\mathbf{z}$ and the gauge function γ, i.e, we obtain equations of motion for these parameters and prove their accuracy to order ϵ^2 for all $t \geq 0$. The process is considered as a reduction of the problem of the analysis of the dynamics generated by Eq. (1.1) to a dynamical problem consisting of a small number (in a sense) of degrees of freedom, providing an effective overall behavior of the original system.

As mentioned above, the proof of Theorem 1 consists of several steps:

Step 1- Decomposition

The first step in the proof is to establish the validity of an appropriate decomposition for any $u \in X^{(n)}$ which is close enough to M_{as}:

Proposition 1 (decomposition) *Define a neighborhood $U_\delta \subset X^{(n)}$ of M_{as} in $X^{(n)}$ as follows*

$$U_\delta = \{u \mid u \in X^{(n)} \text{ and } \exists (\mathbf{z}, \gamma) \in \Sigma,\, \|u - v_{\mathbf{z}\gamma}\|_{X^{(n)}} < \delta\}\,. \tag{2.25}$$

Then there exist $\delta > 0$ and a C^1 map $M_{dec} : U_\delta \mapsto \Sigma$ such that, if we denote $v(u) = \beta(M_{dec}(u))$ for $u \in U_\delta$ (where $\beta(\mathbf{z}, \gamma) = v_{\mathbf{z}\gamma}$; the parametrization mapping for M_{as}), we have $u - v(u) \in H^1(\mathbb{R}^2; \mathbb{C} \times \mathbb{R}^2)$ and

$$P_{v(u)}(u - v(u)) = 0\,. \tag{2.26}$$

□

Proof:

Given the expression in Eq. (2.12) for the projection operator $P_{v_{\mathbf{z}\gamma}}$ we see that condition (2.26) is equivalent to the following conditions

$$\langle T_i^{\mathbf{z}\gamma}, (u - v_{\mathbf{z}\gamma})\rangle = 0,\ i = 1, 2 \qquad \langle G^{\mathbf{z}\gamma}_{\delta_{(x)}}, (u - v_{\mathbf{z},\gamma})\rangle = 0\,. \tag{2.27}$$

Define a function $g : U_\delta \times \Sigma \to \mathbb{R}^2 \times L^2(\mathbb{R}^2)$ by

$$g(u; \mathbf{z}, \gamma) = (\langle T_i^{\mathbf{z}\gamma}, (u - v_{\mathbf{z}\gamma})\rangle, \langle G^{\mathbf{z}\gamma}_{\delta_{(x)}}, (u - v_{\mathbf{z}\gamma})\rangle)\,. \tag{2.28}$$

It is obvious that $g(v_\sigma; \sigma) = 0$. Taking the Fréchet derivative we obtain a map $D_\sigma g(v_\sigma; \sigma) : T_\sigma\Sigma \mapsto \mathbb{R}^2 \times L^2(\mathbb{R}^2)$

$$D_\sigma g(v_\sigma; \sigma) = -(\langle T_i^\sigma, D_\sigma v_\sigma\rangle, \langle G^\sigma_{\delta_{(x)}}, D_\sigma v_\sigma\,\rangle)\,. \tag{2.29}$$

Using the basis of $T_\sigma\Sigma$ given in Eq. (2.4) we can express Eq. (2.28) as a transformation on the coordinate vector in that basis. We then obtain a matrix representation $[D_\sigma g(v_\sigma; \sigma)]_R$ of $D_\sigma g(v_\sigma; \sigma)$ with

$$[D_\sigma g(v_\sigma; \sigma)]_R : \mathbb{R}^2 \times H^2(\mathbb{R}^2; \mathbb{R}) \mapsto \mathbb{R}^2 \times L^2(\mathbb{R}^2)$$

and

$$[D_\sigma g(v_\sigma; \sigma)]_R = diag\{-a_n, -a_n, K_\sigma\}\,. \tag{2.30}$$

Since K_σ is invertible we find that $[D_\sigma g(v_\sigma; \sigma)]_R$ is invertible (hence also $D_\sigma g(v_\sigma; \sigma)$). The implicit function theorem then implies that for any $v_\sigma \in M_{as}$ there is a unique C^1 map M_{dec} from $B_{X^{(n)}}(v_\sigma; \delta)$, a ball in $X^{(n)}$ of size δ centered at v_σ, to Σ such that, for $\sigma = M_{dec}(u)$, Eq. (2.27) and hence Eq. (2.26) are satisfied. We note that $\|K_\sigma^{-1}\|_{L^2\to H^2}$ is uniformly bounded on M_{as} and also that $D_\sigma g(u; \sigma)$ and $D^2_\sigma g(u; \sigma)$ contain only covariant derivatives of T_i^σ and $G^\sigma_{\delta_{(x)}}$. We therefore find that the coordinate transformation representations

$$[D_\sigma g(u; \sigma)]_R : X^{(n)} \times (\mathbb{R}^2 \times H^2(\mathbb{R}^2; \mathbb{R})) \mapsto \mathbb{R}^2 \times L^2(\mathbb{R}^2)$$

and

$$[D^2_\sigma g(u; \sigma)]_R : X^{(n)} \times (\mathbb{R}^2 \times H^2(\mathbb{R}^2; \mathbb{R})) \times (\mathbb{R}^2 \times H^2(\mathbb{R}^2; \mathbb{R})) \mapsto \mathbb{R}^2 \times L^2(\mathbb{R}^2)$$

are uniformly bounded in σ for all balls $B_{X^{(n)}}(v_\sigma; \delta_0)$, where δ_0 is independent of σ. Thus, we can choose δ independent of the point v_σ and the existence of an appropriate neighborhood U_δ is established. ■

Consider a solution $u(t)$ of Eq. (1.1) which, for a time interval $t \in [0, T_\delta]$ satisfies $u(t) \in U_\delta$. Proposition 1 then implies that for such a solution it is possible to find, for each $t \in [0, T_\delta]$ a unique point $v(u(t)) \in M_{as}$ such that the condition in Eq. (2.26) holds. If we denote $\zeta_v(t) = u(t) - v(u(t))$ then $\zeta(t) \in X^{(n)}$ and we obtain a unique decomposition

$$u(t) = v(u(t)) + \zeta_v(t) \tag{2.31}$$

with

$$P_{v(u(t))}\zeta_v(t) = 0\,. \tag{2.32}$$

For the sake of bravity we omit in the sequel from our notation the time t and all indications for the procedure of obtaining the point $v(u(t))$ when given a solution $u(t)$. Thus we write, unless a temporary need arises to restore the full notation, $u(t)$, $v(u(t))$ and $\zeta_v(t)$ as u, v and ζ_v respectively. Eq. (2.31) will then be written in short as $u = v + \zeta_v$, Eq. (2.32) will be written $P_v\zeta_v = 0$, etc.

Step 2- Effective dynamics equations

Once the existence of the decomposition, Eq. (2.31)-(2.32), is established, we can project Eq. (1.1) on the tangent space to M_{as} in order to obtain the effective equations of motion. Applying the projection P_v to Eq. (1.12) we get

$$P_v\dot{u} = -P_v\mathcal{E}'_\epsilon(u)\,. \tag{2.33}$$

Making use of the decomposition in Eq. (2.31) we obtain

$$P_v\dot{v} = -P_v\mathcal{E}'_\epsilon(u) - P_v\dot{\zeta}_v\,. \tag{2.34}$$

We expand in a Taylor series the Fréchet derivative of the energy functional

$$\mathcal{E}'_\epsilon(u) = \mathcal{E}'_\epsilon(v) + L_{\epsilon;v}\zeta_v + N_v(\zeta_v) \tag{2.35}$$

where $L_{\epsilon;v} \equiv \mathcal{E}''_\epsilon(v)$ and $N_v(\zeta_v)$ is the non-linear term defined by $N_v(\zeta_v) = \mathcal{E}'_\epsilon(u) - \mathcal{E}'_\epsilon(v) - L_{\epsilon;v}\zeta_v$. Inserting Eq. (2.35) into Eq. (2.34) we get

$$P_v\dot{v} = -P_v(\mathcal{E}'_\epsilon(v) + L_{\epsilon;v}\zeta_v + N_v(\zeta_v)) - P_v\dot{\zeta}_v\,. \tag{2.36}$$

Note further that $P_v\dot{v} = \dot{v}$ (since $\dot{v} \in T_vM_{as}$) and hence

$$\dot{v} + P_v\mathcal{E}'_\epsilon(v) = -P_vL_{\epsilon;v}\zeta_v - P_vN_v(\zeta_v) - P_v\dot{\zeta}_v\,. \tag{2.37}$$

The last term on the r.h.s. of Eq. (2.37) can be written in a more convenient form by using Eq. (2.32). We have

$$0 = \partial_t(P_v\zeta_v) = \dot{P}_v\zeta_v + P_v\dot{\zeta}_v$$

and so

$$P_v\dot{\zeta}_v = -\dot{P}_v\zeta_v\,. \tag{2.38}$$

Inserting Eq. (2.38) into Eq. (2.37) we obtain

$$\dot{v} + P_v\mathcal{E}'_\epsilon(v) = -P_vL_{\epsilon;v}\zeta_v - P_vN_v(\zeta_v) + \dot{P}_v\zeta_v\,. \tag{2.39}$$

Eq. (2.39) is the starting point for the derivation of the effective dynamics on the manifold M_{as} induced by the gradient flow equations.

We begin by estimating Eq. (2.39):

Proposition 2 *Suppose that a solution u of Eq. (1.1) satisfy the condition that $u(t) \in U_\delta$ for $t \in [0, T_\delta]$, where $U_\delta \subset X^{(n)}$ is the neighborhood of M_{as} given by Proposition 1. For such u the decomposition $u = v + \zeta_v$ is valid and we have the following estimate for $t \in [0, T_\delta]$*

$$\begin{aligned}\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} \le C[&\epsilon\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}} \\ &+ \|\zeta_{v_\sigma}\|^3_{H^1_{v_\sigma}} + (|\dot{\mathbf{z}}| + \|\dot{\tilde{\gamma}}\|_2)\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}]\end{aligned} \tag{2.40}$$

for some $C > 0$. □

Proof:
A first estimate of Eq. (2.39) gives

$$\begin{aligned}\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} \le \|P_{v_\sigma}L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\|_{H^{-s}_{v_\sigma}} &+ \|P_{v_\sigma}N_{v_\sigma}(\zeta_{v_\sigma})\|_{H^{-s}_{v_\sigma}} \\ &+ \|\dot{P}_{v_\sigma}\zeta_{v_\sigma}\|_{H^{-s}_{v_\sigma}}\,.\end{aligned} \tag{2.41}$$

The proposition is a result of Lemma 1, Lemma 2 and Lemma 3 below, which provide estimates on the terms on the r.h.s. of Eq. (2.41):

Lemma 1 (approximate zero modes property) *There exists a constant $C > 0$ such that, for all $v \in M_{as}$ and any vector $\zeta \in T_vX^{(n)}$, we have*

$$\|P_vL_{v;\epsilon}\zeta\|_2 \le \epsilon C\|\zeta\|_2 \tag{2.42}$$

where ϵ is the potential strength parameter appearing in Eq. (1.1). □

Lemma 1 is proved in Section 4. This lemma provides a bound on the first term on the r.h.s. of Eq. (2.41) (since $L^2 = L^2_{v_{\mathbf{z}\gamma}}$). A straightforward technical calculation provides an estimate for the middle term on the r.h.s of Eq. (2.41). This is also done in Section 4 where we obtain the following result:

Lemma 2 *For $\zeta \in T_v X^{(n)}$ and $s > 0$ we have*

$$\|P_{v_\sigma} N_{v_\sigma}(\zeta)\|_{H^{-s}_{v_\sigma}} \leq C(\, \|\zeta\|^2_{H^1_{v_\sigma}} + \|\zeta\|^3_{H^1_{v_\sigma}}\,). \tag{2.43}$$

□

An estimate of the last term on the r.h.s. of Eq. (2.41) is given in Lemma 3, proved in Section 4:

Lemma 3 *For $\zeta_{v_\sigma} \in T_{v_\sigma} X^{(n)}$ satisfying $P_{v_\sigma}\zeta_{v_\sigma} = 0$ we have*

$$\|\dot{P}_{v_\sigma}\zeta_{v_\sigma}\|_2 \leq C(|\dot{\mathbf{z}}| + \|\dot{\tilde{\gamma}}\|_2)\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \tag{2.44}$$

where $\dot{\tilde{\gamma}}$ is given in Eq. (2.14). □

insertion of the inequalities, Eq. (2.42)-(2.44) into Eq. (2.41) completes the proof of Proposition 2.

Step 3- A bound on the error term

Given the decomposition $u = v + \zeta_v$, for a solution u satisfying the conditions of Proposition 1, we show that the remainder term ζ_v is of order ϵ for all times $t > 0$, where ϵ is the potential strength parameter (this result also justifies the assumption that the decomposition implied by Proposition 1 exists for all times if the initial data $u(0)$ satisfies certain requirments which are specified below). The following proposition is the main step in the proof of Theorem 1:

Proposition 3 *Let $u(t)$ be a solution of Eq. (1.1) with initial data $u(0) = u_0 = (\psi_0, A_0)$. There exist constants $\epsilon_0, c_0, c > 0$, with c_0 and c depending only on ϵ_0, such that, for any $0 < \epsilon < \epsilon_0$, if u_0 is chosen in such a way that there exists a point $v_{\mathbf{z}_0\gamma_0} \in M_{as}$ with $\|u_0 - v_{\mathbf{z}_0\gamma_0}\|_{X^{(n)}} \leq c_0\epsilon$, then Proposition 1 holds for $u(t)$ for all $t \geq 0$ and the decomposition $u(t) = v_\sigma(t) + \zeta_{v_\sigma}(t)$ satisfies*

$$\|\zeta_{v_\sigma}(t)\|_{X^{(n)}} \leq c\,\epsilon, \qquad t \geq 0 \tag{2.45}$$

and

$$\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} \leq c\,\epsilon^2, \qquad t \geq 0\,. \tag{2.46}$$

□

Proof: The central result underlying the proof of Theorem 1 is the *Linear Stability Theorem* for the GL equations proved by S. Gustafson and I.M. Sigal [GS1]. This linear stability property corresponds to the *coercivity of the Hessian*. We start the proof of Proposition 3 by stating this theorem in a form convenient for our purposes.

Suppose that the manifold M_{as} corresponds to a vortex with index n, i.e., $M_{as} = M^{(n)}_{sym}$. An n-vortex is called stable if there exists some constant $\nu > 0$ such that

$$L_{0;v} \mid_{(T_v M_{as})^\perp} \geq \nu \, .$$

An n-vortex is called unstable if $L_{0;v}$ has a negative eigenvalue. We have

Linear Stability Theorem (S. Gustafson and I.M. Sigal):

(i) For all $\lambda > 0$ the vortex with $n = \pm 1$ is stable.

(ii) For $\lambda < 1$ a vortex with $|n| \geq 2$ is stable.

(ii) For $\lambda > 1$ a vortex with $|n| \geq 2$ is unstable.

□

In our work here we consider only stable vortices. Then $L_{0;v} \mid_{(T_v M_{as})^\perp}$ is a positive operator. In this case, for any vector $\zeta_v \in T_v X^{(n)}$ such that $P_v \zeta_v = 0$, the linear stability theorem implies that

$$\langle \zeta_v, L_{0;v} \zeta_v \rangle \geq \nu \| \zeta_v \|^2_{H^1} \tag{2.47}$$

for some constant $\nu > 0$. Introducing the potential W along with the potential strength parameter ϵ we have the following corollary to the linear stability theorem:

Corollary A: *For a vector $\zeta_v \in T_v X^{(n)}$ with $P_v \zeta_v = 0$ and for $\epsilon > 0$ we have*

$$\langle \zeta_v, L_{\epsilon;v} \zeta_v \rangle \geq \nu \| \zeta_v \|^2_{H^1} \, . \tag{2.48}$$

□

Proof: We calculate

$$\begin{aligned} \langle \zeta_v, L_{\epsilon;v} \zeta_v \rangle = \langle \zeta_v, (L_{0;v} + \epsilon W) \zeta_v \rangle = \langle \zeta_v, L_{0;v} \zeta_v \rangle + \epsilon \langle \zeta_v, W \zeta_v \rangle \\ \geq \langle \zeta_v, L_{0;v} \zeta_v \rangle \geq \nu \| \zeta_v \|^2_{H^1} \end{aligned}$$

since the term containing the potential W is positive. ■

The following result, which we use below, is also an immediate corollary of the linear stability theorem

Corollary B: *For a vector $\zeta_{v_\sigma} \in T_{v_\sigma} X^{(n)}$ satisfying $P_{v_\sigma}\zeta_{v_\sigma} = 0$ and for $\epsilon > 0$ we have*

$$\langle \zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \geq \nu \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}} \,. \tag{2.49}$$

□

Proof: It is easy to check that the application of the gauge transformation $G_{-\gamma}$ to the projection $P_{v_{\mathbf{z}\gamma}}$ yields $G_{-\gamma}P_{v_{\mathbf{z}\gamma}}G^*_{-\gamma} = G_{-\gamma}P_{v_{\mathbf{z}\gamma}}G_\gamma = P_{v_{\mathbf{z}0}}$. This implies that

$$0 = G_{-\gamma}P_{v_\sigma}\zeta_{v_\sigma} = G_{-\gamma}P_{v_\sigma}G^*_{-\gamma}G_{-\gamma}\zeta_{v_\sigma} = P_{v_{\mathbf{z}0}}\zeta_{v_{\mathbf{z}0}} \tag{2.50}$$

where $\zeta_{v_{\mathbf{z}0}} = G_{-\gamma}\zeta_{v_\sigma}$. In addition we have $G_{-\gamma}L_{\epsilon;v_\sigma} = L_{\epsilon;v_{\mathbf{z}0}}G_{-\gamma}$. Therefore, we obtain

$$\begin{aligned}\langle \zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle &= \langle G_{-\gamma}\zeta_{v_\sigma}, G_{-\gamma}L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \\ &= \langle \zeta_{v_{\mathbf{z}0}}, L_{\epsilon;v_{\mathbf{z}0}}\zeta_{v_{\mathbf{z}0}}\rangle \geq \nu\|\zeta_{v_{\mathbf{z}0}}\|^2_{H^1} = \nu\|G_{-\gamma}\zeta_{v_\sigma}\|^2_{H^1} = \nu\|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}} \,.\end{aligned} \tag{2.51}$$

■

In the proof of Proposition 3 we make use of the dominating linear stability properties mentioned above by analyzing the time evolution and providing an upper bound on the quantity $\langle \zeta_v, L_{\epsilon;v}\zeta_v\rangle$. Corollary A of the linear stability theorem then ensures that this yields a bound on the error term ζ_v.

In Section 3 we prove the following proposition:

Proposition 4 *Let u be a solution of Eq. (1.1) satisfying the assumption of Proposition B above and let $u = v + \zeta_v$ be the decomposition implied by Proposition 1. Then, for $t \in [0, T_\delta]$, there exist constants $\nu', C_1, C_2, C_3 > 0$ such that*

$$\begin{aligned}&\frac{1}{2}\partial_t\langle \zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \leq -\frac{1}{2}\nu'\langle \zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle + \big\{ -C_1 \\ &+ C_2\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\big[(2 + \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}})(1 + \epsilon + \|\dot v_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}}) - 1\big]\big\}\|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} \\ &+ \epsilon C_3(1 + \|\dot v_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}})\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \,.\end{aligned} \tag{2.52}$$

□

Assume that there exists a maximum time T_1 such that

$$\|\ddot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} \leq \epsilon, \qquad t \in [0, T_1]\,. \tag{2.53}$$

By Eq. (2.52) and Eq. (2.53) we have in this time interval

$$\begin{aligned}\frac{1}{2}\partial_t\langle\zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \leq\ & -\frac{1}{2}\nu'\,\langle\zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \\ & + \big\{-C_1 + C_2\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\big[(2 + \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}})(1+2\epsilon) - 1\big]\big\}\|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} \\ & + \epsilon C_3(1 + \epsilon + \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}})\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\,.\end{aligned} \tag{2.54}$$

Let $\delta > 0$ be such that the decomposition $u = v_\sigma(u) + \zeta_{v_\sigma}$ implied by Proposition 1 holds for any $u \in U_\delta \subset X^{(n)}$ (see Eq. (2.25) for the definition of U_δ). The existence of such a δ is guaranteed by Proposition 1. Suppose that there exists a maximum time T_2 such that

$$\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \leq \min\left\{1, \frac{C_1}{10C_2}, \delta\right\}\,, \qquad t \in [0, T_2]\,. \tag{2.55}$$

Set $\tau = min\{T_1, T_2\}$. Then for $t \in [0, \tau]$ we have, for some $\tilde{C}, C > 0$,

$$\frac{1}{2}\partial_t\langle\zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \leq -\frac{1}{2}\nu'\langle\zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle - \frac{C_1}{2}\|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} + \epsilon\frac{C}{2}\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\,. \tag{2.56}$$

Dropping the negative definite second term on the r.h.s. of Eq. (2.56) and using Corollary B of the linear stability theorem we get

$$\frac{1}{2}\partial_t\langle\zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \leq -\frac{1}{2}\nu'\langle\zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle + \epsilon\frac{\alpha}{2}\langle\zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle^{1/2} \tag{2.57}$$

where $\alpha > 0$. Set $\langle\zeta_{v_\sigma}(t), L_{\epsilon;v_\sigma(t)}\zeta_{v_\sigma}(t)\rangle = f^2(t)$, then Eq. (2.57) reads

$$\frac{d}{dt}f \leq -\nu' f + \epsilon\alpha\,. \tag{2.58}$$

Let $g(t)$ be a solution of Eq. (2.58) with an equality sign and initial condition $g(0) = f(0) = \langle\zeta_{v_\sigma}(0), L_{\epsilon;v_\sigma(0)}\zeta_{v_\sigma}(0)\rangle^{1/2}$ i.e.,

$$\frac{d}{dt}g = -\nu' g + \epsilon\alpha\,.$$

We get

$$g(t) = f(0)e^{-\nu' t} + \frac{\epsilon\alpha}{\nu'}(1 - e^{-\nu' t}) \leq f(0) + \frac{\epsilon\alpha}{\nu'}\,.$$

With the help of Lemma 6, stated in Section 3, we observe that

$$f^2(0) = \langle \zeta_{v_\sigma}(0), L_{\epsilon;v_\sigma(0)}\zeta_{v_\sigma}(0)\rangle = \langle \zeta_{v_\sigma}(0), \overline{P}_{v_\sigma}(0)L_{\epsilon;v_\sigma(0)}\zeta_{v_\sigma}(0)\rangle$$
$$\leq \|\zeta_{v_\sigma}(0)\|_{H^1_{v_\sigma(0)}} \|\overline{P}_{v_\sigma(0)}L_{\epsilon;v_\sigma(0)}\zeta_{v_\sigma}(0)\|_{H^{-1}_{v_\sigma(0)}} \leq C\|\zeta_{v_\sigma}(0)\|^2_{H^1_{v_\sigma(0)}}$$
$$\leq C'\|\zeta_{v_\sigma}(0)\|^2_{H^1} .$$

The validity of the last inequality stems from the fact that $v_\sigma(0)$ is some fixed point on M_{as} and we can estimate the gauge function there.

Now, Proposition 1 guarantees the existence of a neighborhood U_δ of M_{as} in which the decomposition property holds. Take $u(0)$ to be an initial data for Eq. (1.1) with $u(0) \in U_\delta$. Then we have $\zeta_{v_\sigma}(0) = u(0) - v(u(0))$ with $P_{v(u(0))}\zeta_{v_\sigma}(0) = 0$. Assume furthermore that $u(0)$ is chosen to satisfy the condition that there exists a point $v_{\mathbf{z}_0\gamma_0} = v_{\sigma_0} \in M_{as}$ such that $\|u(0) - v_{\sigma_0}\|_{X^{(n)}} = \|u(0) - v_{\sigma_0}\|_{H^1} \leq c_0\epsilon$ for some constant $c_0 > 0$. Under these conditions we have

$$\|\zeta_{v_\sigma}(0)\|_{H^1} \leq \|u(0) - v_{\sigma_0}\|_{H^1} + \|v_{\sigma_0} - v(u(0))\|_{H^1} \leq c_0\epsilon + \|v_{\sigma_0} - v(u(0))\|_{H^1} .$$

Denoting $u' = v_{\sigma_0}$, the solution $v(u')$ satisfying the orthogonality condition, Eq. (2.26), is simply $v(u') = u'$. We need to show that for u'' in the vicinity of $u' = v_{\sigma_0}$ we have

$$\|v(u') - v(u'')\|_{H^1} \leq c'\|u' - u''\|_{H^1} \tag{2.59}$$

for some constant $c' > 0$ which may depend only on ϵ, since then we would have $\|v_{\sigma_0} - v(u(0))\|_{H^1} \leq c'\|v_{\sigma_0} - u(0)\|_{H^1} \leq c'c_0\epsilon$. In order to see that Eq. (2.59) holds near $u' = v_{\sigma_0}$ we first observe that by the implicit function argument in the proof of Proposition 1 we have

$$0 = D_u[g(u; M_{dec}(u))] = D_u g(u; M_{dec}(u)) + D_u M_{dec}(u)\, D_\sigma g(u; M_{dec}(u))$$

and hence

$$D_u M_{dec}(u) = -[D_\sigma g(u; M_{dec}(u))]^{-1} D_u g(u; M_{dec}(u)) .$$

It follows that $\|D_u M_{dec}(v_{\sigma_0})\|_{X^{(n)} \to \mathbb{R}^2 \times H^2} \leq C$; since Eq. (2.30) (recall that K_σ is uniformly invertible on M_{as}) and the definition of $g(u;\sigma)$ in Eq. (2.28) imply, respectively, that the two norms $\| [D_\sigma g(v_{\sigma_0}; \sigma_0)]^{-1}\|_{\mathbb{R}^2 \times L^2 \to \mathbb{R}^2 \times H^2}$ and

$\|D_u g(v_{\sigma_0};\sigma_0)\|_{X^{(n)}\to\mathbb{R}^2\times L^2}$ are uniformly bounded on M_{as}. This observation leads to the following bound

$$\begin{aligned}\|D_u v(v_{\sigma_0})\|_{X^{(n)}\to X^{(n)}} &= \|D_u[\beta(M_{dec}(v_{\sigma_0}))]\ \|_{X^{(n)}\to X^{(n)}} \\ &\le \|\ (D_\sigma v_\sigma)_{\sigma_0}\|_{\mathbb{R}^2\times H^2\to X^{(n)}}\ \|D_u M_{dec}(v_{\sigma_0}))\|_{X^{(n)}\to\mathbb{R}^2\times H^2} \le C\end{aligned}$$

and we conclude that, for $c_0 = c_0(\epsilon)$ small enough, if $\|u(0) - v_{\sigma_0}\|_{H^1} \le c_0\epsilon$ then $f(0) \le c\epsilon$ for some constant $c = c(\epsilon) > 0$. Hence, in the time interval $t \in [0,\tau]$, we have

$$\|\zeta_v\|_{H^1} \le \nu^{-1}\langle\zeta_v, L_{\epsilon;v}\zeta_v\rangle^{1/2} = \nu^{-1}f(t) \le \nu^{-1}g(t) \le C\epsilon\,. \tag{2.60}$$

In order to close the proof of Proposition 3 we need an estimate on the time derivatives $\dot{\mathbf{z}}$ and $\dot{\tilde{\gamma}}$ appearing in the last term on the r.h.s of Eq. (2.40). The following lemma is proved in Section 4:

Lemma 4 *For $s > 0$ we have the following estimate for the parametric equations of motion*

$$|a_n\dot{\mathbf{z}} + \partial^A_{\mathbf{z}}\mathcal{E}_\epsilon(v_\sigma)| + \|\dot{\tilde{\gamma}}\|_{H^{1-s}} \le C\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} \tag{2.61}$$

where $\dot{\tilde{\gamma}}$ is defined in Eq. (2.14). □

Since $\partial^A_{z_i}\mathcal{E}_\epsilon(v_\sigma) = o(\epsilon)$ (see Eq. (2.67) below) we get that for $0 < s < 1$

$$\begin{aligned}|\dot{\mathbf{z}}| + \|\dot{\tilde{\gamma}}\|_2 &\le a_n^{-1}(|a_n\dot{\mathbf{z}} + \partial^A_{\mathbf{z}}\mathcal{E}_\epsilon(v_\sigma)| + |\partial^A_{\mathbf{z}}\mathcal{E}_\epsilon(v_\sigma)|) + \|\dot{\tilde{\gamma}}\|_{H^{1-s}} \\ &\le C(\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \epsilon)\end{aligned} \tag{2.62}$$

then, under the assumption made in Eq. (2.53), we have

$$|\dot{\mathbf{z}}| + \|\dot{\tilde{\gamma}}\|_2 \le \epsilon C, \qquad t \in [0,\tau]\,. \tag{2.63}$$

Eq. (2.40), Eq. (2.60) and Eq. (2.63) then imply that for $t \in [0,\tau]$ the following inequality is valid

$$\begin{aligned}\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} \le C[&\epsilon\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}} \\ &+\|\zeta_{v_\sigma}\|^3_{H^1_{v_\sigma}} + (|\dot{\mathbf{z}}| + \|\dot{\tilde{\gamma}}\|_2)\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}] \le C\epsilon^2\,.\end{aligned} \tag{2.64}$$

The definition of τ implies that at $t = \tau$ at least one of the conditions, Eq. (2.53) or Eq. (2.55), is no longer valid. However, we can choose ϵ_0 in such a way that for $0 < \epsilon < \epsilon_0$ this conclusion will stand in contradiction with

the results, Eq. (2.60) and Eq. (2.64). Hence $\tau = \infty$ and the proof of Proposition 3 is complete. ■

Step 4 - Effective equations for the parameters

We derive the appropriate equations for the variables $\mathbf{z}$ and γ, parametrizing the manifold M_{as}. Thus, solutions of Eq. (1.1) with an initial condition close enough to M_{as} induce, up to a small, controllable error, an effective dynamics on M_{as} in terms of equations of motion for the vortex center and gauge function in the presence of the potential W.

The effective equations of motion for the parameters $\mathbf{z}$ and $\tilde{\gamma}$ are derived using Eq. (2.61) and Proposition 3. For $\tilde{\gamma}$ we have simply

$$\|\dot{\tilde{\gamma}}\|_{H^{1-s}} \leq C\epsilon^2 \tag{2.65}$$

and for $\mathbf{z}$ we get

$$|a_n \dot{\mathbf{z}} + \partial_{\mathbf{z}}^A \mathcal{E}_\epsilon(v_\sigma)| \leq C\epsilon^2 \tag{2.66}$$

Since $\mathcal{E}_\epsilon(v_\sigma) = \mathcal{E}_{int}(v_\sigma) = W_{int}^{(n)}(\mathbf{z})$, where

$$W_{int}^{(n)}(\mathbf{z}) \equiv \epsilon \int d^2x\, W(\mathbf{x}) |\psi^{(n)}(\mathbf{x} - \mathbf{z})|^2 \tag{2.67}$$

does not depend on the vector potential A, we get finally

$$|a_n \dot{\mathbf{z}} + \nabla_{\mathbf{z}} W_{int}(\mathbf{z})| \leq C\epsilon^2 \tag{2.68}$$

This derivation of the parametric equations of motion completes the proof of Theorem 1.

3 Proof of Proposition 4

We calculate

$$\begin{aligned}
\frac{1}{2}\partial_t \langle \zeta_v, L_{\epsilon,v}\zeta_v\rangle &= \partial_t(\mathcal{E}_\epsilon(u) - \mathcal{E}_\epsilon(v) - \langle \mathcal{E}'_\epsilon(v), \zeta_v\rangle - R_v(\zeta_v)) = \\
&= \langle \mathcal{E}'_\epsilon(u), \dot{u}\rangle - \langle \mathcal{E}'_\epsilon(v), \dot{v}\rangle - \langle \mathcal{E}''_\epsilon(v)\dot{v}, \zeta_v\rangle - \langle \mathcal{E}'_\epsilon(v), \dot{\zeta}_v\rangle - \partial_t[R_v(\zeta_v)] = \\
&= \langle \mathcal{E}'_\epsilon(u) - \mathcal{E}'_\epsilon(v), \dot{u}\rangle - \langle L_{\epsilon,v}\dot{v}, \zeta_v\rangle - \partial_t[R_v(\zeta_v)] = \\
&= -\langle \mathcal{E}'_\epsilon(u) - \mathcal{E}'_\epsilon(v), \mathcal{E}'_\epsilon(u)\rangle - \langle L_{\epsilon,v}\dot{v}, \zeta_v\rangle - \partial_t[R_v(\zeta_v)] \\
= -\langle \mathcal{E}'_\epsilon(u) - \mathcal{E}'_\epsilon(v), \overline{P}_v \mathcal{E}'_\epsilon(u)\rangle &+ \langle \mathcal{E}'_\epsilon(v) - \mathcal{E}'_\epsilon(u), P_v \mathcal{E}'_\epsilon(u)\rangle - \langle L_{\epsilon,v}\dot{v}, \zeta_v\rangle - \partial_t[R_v(\zeta_v)]
\end{aligned} \tag{3,1}$$

where the energy remainder term $R_v(\zeta)$ is given in Appendix A, Eq. (A.5). Next we define the quantity $\dot{R}_v(\zeta_v)$ by

$$\partial_t[R_v(\zeta_v)] = \dot{R}_v(\zeta_v) + \langle N_v(\zeta_v), \dot{\zeta}_v \rangle \,. \tag{3.2}$$

Consider the equation of motion for the error term ζ_v. This equation is obtained by projecting Eq. (1.12) on $(T_v M_{as})^{\perp}$. We have

$$\dot{\zeta}_v = -\overline{P}_v \mathcal{E}'_\epsilon(u) - \dot{P}_v \zeta_v \,. \tag{3.3}$$

From Eq. (3.2) and Eq. (3.3) we get

$$\partial_t[R_v(\zeta_v)] = \dot{R}_v(\zeta_v) - \langle N_v(\zeta_v), \overline{P}_v \mathcal{E}'_\epsilon(u) \rangle - \langle N_v(\zeta_v), \dot{P}_v \zeta_v \rangle \,. \tag{3.4}$$

Inserting Eq. (3.4) into Eq. (3.1) we obtain the following inequality

$$\begin{aligned}
\frac{1}{2}\partial_t \langle \zeta_v, L_{\epsilon,v}\zeta_v \rangle &= -\langle L_{\epsilon,v}\zeta_v, \overline{P}_v \mathcal{E}'_\epsilon(u) \rangle - \langle \mathcal{E}'_\epsilon(u) - \mathcal{E}'_\epsilon(v), P_v \mathcal{E}'_\epsilon(u) \rangle \\
&\quad - \langle L_{\epsilon,v}\dot{v}, \zeta_v \rangle - \dot{R}_v(\zeta_v) + \langle N_v(\zeta_v), \dot{P}_v \zeta_v \rangle = \\
&\leq -\langle L_{\epsilon,v}\zeta_v, \overline{P}_v \mathcal{E}'_\epsilon(u) \rangle - \langle \mathcal{E}'_\epsilon(u) - \mathcal{E}'_\epsilon(v), P_v \mathcal{E}'_\epsilon(v) \rangle \\
&\quad - \langle L_{\epsilon,v}\dot{v}, \zeta_v \rangle - \dot{R}_v(\zeta_v) + \langle N_v(\zeta_v), \dot{P}_v \zeta_v \rangle = \\
&= -\langle \overline{P}_v L_{\epsilon,v}\zeta_v, \overline{P}_v L_{\epsilon,v}\zeta_v \rangle + \langle L_{\epsilon,v}\zeta_v, \overline{P}_v(\mathcal{E}'_\epsilon(v) + N_v(\zeta_v)) \rangle \\
&\quad - \langle L_{\epsilon;v}\zeta_v, \dot{v} + P_v \mathcal{E}'_\epsilon(v) \rangle - \langle N_v(\zeta_v), P_v \mathcal{E}'_\epsilon(v) \rangle - \dot{R}_v(\zeta_v) + \langle N_v(\zeta_v), \dot{P}_v \zeta_v \rangle
\end{aligned} \tag{3.5}$$

and so

$$\begin{aligned}
&\frac{1}{2}\partial_t \langle \zeta_v, L_{\epsilon;v}\zeta_v \rangle \leq -\langle \overline{P}_v L_{\epsilon,v}\zeta_v, \overline{P}_v L_{\epsilon,v}\zeta_v \rangle + |\langle L_{\epsilon,v}\zeta_v, \overline{P}_v(\mathcal{E}'_\epsilon(v) + N_v(\zeta_v)) \rangle| \\
&+ |\langle L_{\epsilon;v}\zeta_v, \dot{v} + P_v \mathcal{E}'_\epsilon(v) \rangle| + |\langle N_v(\zeta_v), P_v \mathcal{E}'_\epsilon(v) \rangle| + |\dot{R}_v(\zeta_v)| + |\langle N_v(\zeta_v), \dot{P}_v \zeta_v \rangle|
\end{aligned} \tag{3.6}$$

In Section 4 we establish, via straightforward calculations, the validity of the following estimates

Lemma 5 *For $\zeta \in T_v X^{(n)}$ we have*

$$\|N_{v_\sigma}(\zeta)\|_2 \leq C \|\zeta\|_{H^2_{v_\sigma}} (\|\zeta\|_{H^1_{v_\sigma}} + \|\zeta\|^2_{H^1_{v_\sigma}}) \tag{3.7}$$

□

and

Lemma 6 *For $\zeta \in T_v X^{(n)}$ the following inequalities hold*

$$\|\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta\|_{H^{-1}_{v_\sigma}} \leq C\|\zeta\|_{H^1_{v_\sigma}}\,, \qquad \|\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta\|_2 \leq C\|\zeta\|_{H^2_{v_\sigma}}\,. \tag{3.8}$$

□

Eq. (3.7), Eq. (3.8) and the easily obtainable inequality

$$\|\mathcal{E}'_\epsilon(v_\sigma)\|_{H^1_{v_\sigma}} = \epsilon\|W\psi_{\mathbf{z}0}\|_{H^1} \leq \epsilon C \tag{3.9}$$

result in the following estimates:

$$|\langle \overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}, \mathcal{E}'_\epsilon(v_\sigma)\rangle| \leq \|\overline{P}_{v_\sigma} L_{\epsilon,v_\sigma}\zeta_{v_\sigma}\|_{H^{-1}_{v_\sigma}}\|\mathcal{E}'_\epsilon(v_\sigma)\|_{H^1_{v_\sigma}} \leq \epsilon\, C\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \tag{3.10}$$

and

$$\begin{aligned}|\langle \overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}, N_{v_\sigma}(\zeta_{v_\sigma})\rangle| &\leq \|\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\|_2\|N_{v_\sigma}(\zeta_{v_\sigma})\|_2 \\ &\leq C\|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}}(\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}})\,.\end{aligned} \tag{3.11}$$

Furthermore, we have the two estimates

$$\begin{aligned}&|\langle L_{\epsilon;v_\sigma}\zeta_{v_\sigma}, \dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle| = |\langle P_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}, \dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle| \\ &\leq \|P_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-1}_{v_\sigma}} \leq \epsilon C\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}}\end{aligned} \tag{3.12}$$

(see the proof of Lemma 1) and

$$\begin{aligned}|\langle N_{v_\sigma}(\zeta_{v_\sigma}), P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle| &\leq \|P_{v_\sigma} N_{v_\sigma}(\zeta_{v_\sigma})\|_{H^{-1}_{v_\sigma}}\|\mathcal{E}'_\epsilon(v_\sigma)\|_{H^1_{v_\sigma}} \\ &\leq \epsilon C(\|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^3_{H^1_{v_\sigma}})\end{aligned} \tag{3.13}$$

where in Eq. (3.13) use has been made of Eq. (3.9) and Lemma 2.

In order to obtain a bound on the term $\dot{R}_v(\zeta_v)$ we note that Eq. (3.2) implies

$$\dot{R}_v(\zeta) = \big(\partial_t R_v(\zeta)\big)_{\zeta=const.} = \langle \dot{v}, \partial_v R_v(\zeta)\rangle \tag{3.14}$$

hence, we have

$$\begin{aligned}|\dot{R}_{v_\sigma}(\zeta_{v_\sigma})| &= |\langle \dot{v}_\sigma, \partial_{v_\sigma} R_{v_\sigma}(\zeta_{v_\sigma})\rangle| \\ &\leq (\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \|P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}})\|\partial_{v_\sigma} R_{v_\sigma}(\zeta_{v_\sigma})\|_{H^s_{v_\sigma}}\,.\end{aligned} \tag{3.15}$$

The following estimate is proved in Section 4:

Lemma 7 *For any $\zeta \in T_{v_\sigma} X$ we have*

$$\|\partial_{v_\sigma} R_{v_\sigma}(\zeta)\|_{H^1_{v_\sigma}} \leq C\|\zeta\|^2_{H^2_{v_\sigma}} \|\zeta\|_{H^1_{v_\sigma}} \,. \tag{3.16}$$

□

Lemma 7 and Eq. (3.9) then imply the following bound for $0 < s < 1$

$$|\dot{R}_{v_\sigma}(\zeta_{v_\sigma})| \leq C(\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \epsilon)\|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \,. \tag{3.17}$$

With Eq. (3.10), (3.11), (3.12), (3.13) and (3.17) providing appropriate bounds on the corresponding terms in Eq. (3.6) we get

$$\begin{aligned}\frac{1}{2}\partial_t \langle \zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle &\leq -\langle \overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}, \overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \\ &+ C(1+\epsilon+\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}})\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} \\ &+ \epsilon C(1+\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}})\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \\ &+ |\langle N_{v_\sigma}(\zeta_{v_\sigma}), \dot{P}_{v_\sigma}\zeta_{v_\sigma}\rangle| \,.\end{aligned} \tag{3.18}$$

We still need to estimate the last term on the r.h.s. of Eq. (3.18). Using Lemma 3 and Eq. (2.62) we obtain

$$\|\dot{P}_{v_\sigma}\zeta_{v_\sigma}\|_2 \leq C(\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E'}_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \epsilon)\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \,. \tag{3.19}$$

With the help of Lemma 5 and Eq. (3.19) we can provide the appropriate bound in the form

$$\begin{aligned}|\langle N_{v_\sigma}(\zeta_{v_\sigma}), \dot{P}_{v_\sigma}\zeta_{v_\sigma}\rangle| &\leq \|N_{v_\sigma}(\zeta_{v_\sigma})\|_2 \|\dot{P}_{v_\sigma}\zeta_{v_\sigma}\|_2 \\ &\leq C(\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \epsilon)(1+\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}})\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} \,.\end{aligned} \tag{3.20}$$

Inserting the bound obtained in Eq. (3.20) into the r.h.s. of Eq. (3.18) we get

$$\begin{aligned}\frac{1}{2}\partial_t \langle \zeta_{v_\sigma}, L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle &\leq -\langle \overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}, \overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta_{v_\sigma}\rangle \\ + C_2\big[(2+\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}})(1+\epsilon+\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}}) - 1\big]\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}}\|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} \\ + \epsilon C_3(1+\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}})\|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \,.\end{aligned} \tag{3.21}$$

The first term on the r.h.s. of Eq. (3.21) is handled by utilizing the linear stability properties of the Ginzburg-Landau equations. The Linear Stability Theorem is used in Section 4 in the proof of the following lemma:

Lemma 8 *For $\zeta_v \in T_v X^{(n)}$ orthogonal to $T_v M_{as}$, that is for $P_v \zeta_v = 0$, we have*

$$\langle \overline{P}_{v_\sigma} L_{\epsilon;v_\sigma} \zeta_{v_\sigma}, \overline{P}_{v_\sigma} L_{\epsilon;v_\sigma} \zeta_{v_\sigma} \rangle \geq 2C_1 \|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} \tag{3.22}$$

(the choice of the form of the constant here is purely for reasons of convenience). □

We will also need an additional lemma which is also proved in Section 4:

Lemma 9 *For $\zeta \in T_v X^{(n)}$ with $\zeta_v \in (T_v M_{as})^\perp$ we have*

$$\langle \overline{P}_v L_{\epsilon;v} \zeta_v, \overline{P}_v L_{\epsilon;v} \zeta_v \rangle \geq \nu' \langle \zeta_v, L_{\epsilon;v} \zeta_v \rangle \,. \tag{3.23}$$

□

Eq. (3.21), Lemma 8 and Lemma 9 imply that

$$\begin{aligned}
&\frac{1}{2}\partial_t \langle \zeta_{v_\sigma}, L_{\epsilon;v_\sigma} \zeta_{v_\sigma} \rangle \leq -\frac{1}{2}\nu' \langle \zeta_{v_\sigma}, L_{\epsilon;v_\sigma} \zeta_{v_\sigma} \rangle + \|\zeta_{v_\sigma}\|^2_{H^2_{v_\sigma}} \times \\
&\Big\{ -C_1 + C_2 \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \left[(2 + \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}})(1 + \epsilon + \|\dot{v}_\sigma + P_{v_\sigma} \mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}}) - 1 \right] \Big\} \\
&\quad + \epsilon C_3 \left(1 + \|\dot{v}_\sigma + P_{v_\sigma} \mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}} + \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} + \|\zeta_{v_\sigma}\|^2_{H^1_{v_\sigma}} \right) \|\zeta_{v_\sigma}\|_{H^1_{v_\sigma}} \,.
\end{aligned} \tag{3.24}$$

Eq. (3.24) is the statement of proposition 4. ■

4 Proofs of Lemmas 1-9

Proof of Lemma 3:

The starting point to the proof of Lemma 3 is the explicit expression for the projection P_{v_σ} given in Eq. (2.12). Taking the time derivative of P_{v_σ} and applying the resulting operator to a fixed vector $\zeta \in T_{v_\sigma} X^{(n)}$ satisfying the condition $P_{v_\sigma} \zeta = 0$ we get

$$\dot{P}_{v_\sigma} \zeta = a_n^{-2} \sum_{i=1,2} T_i^\sigma \partial_t \langle T_i^\sigma, \zeta \rangle + \int d^2x \, G^\sigma_{\delta(x)} K_\sigma^{-1}(x,y) \partial_t \langle G^\sigma_{\delta(y)}, \zeta \rangle \,. \tag{4.1}$$

Eq. (4.1) enable us to obtain the L^2 norm of $\dot{P}_v\zeta$

$$\begin{aligned}\|\dot{P}_{v_\sigma}\zeta\|_2^2 &= a_n^{-2}\sum_{i=1,2}|\langle\partial_t T_i^\sigma,\zeta\rangle|^2 \\ &+\int d^2x\int d^2y\;\overline{\partial_t\langle G^\sigma_{\delta(x)},\zeta\rangle}K^{-1}_\sigma(x,y)\partial_t\langle G^\sigma_{\delta(y)},\zeta\rangle\,.\end{aligned} \tag{4.2}$$

Consider the first term on the r.h.s. of Eq. (4.2). The time derivative of T_i^σ is given by

$$\partial_t T_i^\sigma = i\dot{\tilde{\gamma}}\begin{pmatrix}(\nabla_{A_\sigma}\psi_\sigma)_i\\0\end{pmatrix}+\Sigma_j(-\dot{z}_j)(\nabla_{A_\sigma}T_i^\sigma)_j \tag{4.3}$$

where

$$(\nabla_{A_\sigma}T_i^\sigma)_j=\begin{pmatrix}(\nabla_{A_\sigma}(\nabla_{A_\sigma}\psi_\sigma)_i)_j\\ \partial_j(\nabla\times A_\sigma)\hat{e}_i\end{pmatrix}\,. \tag{4.4}$$

Thus we arrive at the following estimate

$$\begin{aligned}\|\partial_t T_i^\sigma\|_2 &\le \|\dot{\tilde{\gamma}}(\nabla_{A_\sigma}\psi_\sigma)_i\|_2+\Sigma_j|\dot{z}_j|\;\|(\nabla_{A_\sigma}T_i^\sigma)_j\|_2\\ &\le \|\dot{\tilde{\gamma}}\|_2\|(\nabla_{A_\sigma}\psi_\sigma)_i\|_\infty+\Sigma_j|\dot{z}_j|\;\|(\nabla_{A_\sigma}T_i^\sigma)_j\|_2\,.\end{aligned} \tag{4.5}$$

The norms of the two vectors appearing on the r.h.s. of Eq. (4.5) are gauge and translation invariant, hence they are constant at all points of M_{as}. We conclude that the following inequality is valid

$$\|\partial_t T_i^\sigma\|_2\le C(\|\dot{\tilde{\gamma}}\|_2+\Sigma_i|\dot{z}_i|) \tag{4.6}$$

and so

$$\begin{aligned}|\langle\partial_t T_i^\sigma,\zeta\rangle| &\le \|\partial_t T_i^\sigma\|_2\|\zeta\|_2\le C(\|\dot{\tilde{\gamma}}\|_2+\sum_i|\dot{z}_i|\,)\|G_{-\gamma}\zeta\|_2\\ &\le C(\|\dot{\tilde{\gamma}}\|_2+\sum_i|\dot{z}_i|\,)\|G_{-\gamma}\zeta\|_{H^1}=C(\|\dot{\tilde{\gamma}}\|_2+\sum_i|\dot{z}_i|\,)\|\zeta\|_{H^1_{v_\sigma}}\,.\end{aligned} \tag{4.7}$$

Consider now the second term on the r.h.s. of Eq. (4.2). Let $\zeta=(\xi,F)\in T_vX^{(n)}$, the time derivative $\partial_t\langle G^\sigma_\chi,\zeta\rangle$ is given by (note that ζ is a fixed vector)

$$\begin{aligned}\partial_t\langle G^\sigma_\chi,\zeta\rangle=\partial_t\langle\int d^2x\,\chi(x)G^\sigma_{\delta(x)},\zeta\rangle&=\int d^2x\,\chi(x)\partial_t\langle G^\sigma_{\delta(x)},\zeta\rangle=\\ =\partial_t\int d^2x\,(Im[\chi\overline{\psi}_\sigma\xi]-\nabla\chi F)&=\int d^2x\,\chi(x)Im[\partial_t\overline{\psi}_\sigma\xi](x)\end{aligned} \tag{4.8}$$

hence

$$\partial_t \langle G^\sigma_{\delta(x)}, \zeta\rangle = Im[\partial_t \overline{\psi}_\sigma(x)\xi(x)]\,. \tag{4.9}$$

Now, K_σ^{-1} are uniformly bounded on M_{as} as operators from H^{-1} to H^1 (see [ST1]) and we can estimate

$$\begin{aligned}
&\left| \int d^2x \int d^2y\, \overline{\partial_t \langle G^\sigma_{\delta(x)}, \zeta\rangle} K_\sigma^{-1}(x,y) \partial_t \langle G^\sigma_{\delta(y)}, \zeta\rangle \right| = \\
&\left|(\partial_t \langle G^\sigma_{\delta(x)}, \zeta\rangle, K_\sigma^{-1}\, \partial_t \langle G^\sigma_{\delta(x)}, \zeta\rangle)\right| \le \|\partial_t \langle G^\sigma_{\delta(x)}, \zeta\rangle\|_{H^{-1}} \|K_\sigma^{-1}\, \partial_t \langle G^\sigma_{\delta(x)}, \zeta\rangle)\|_{H^1} \\
&\le \|K_\sigma^{-1}\|_{H^{-1}\to H^1} \|\partial_t \langle G^\sigma_{\delta(x)}, \zeta\rangle\|^2_{H^{-1}} = C\|Im[\partial_t \overline{\psi}_\sigma \xi]\|^2_{H^{-1}} \le C\|\partial_t \psi_\sigma \xi\|^2_{H^{-1}}
\end{aligned} \tag{4.10}$$

were here again $\zeta = (\xi, F)$. We want to estimate the r.h.s. of this inequality. We have

$$\begin{aligned}
\|\partial_t \overline{\psi}_\sigma \xi\|_{H^{-1}} &= \sup_{\|\alpha\|_{H^1}\le 1} \langle \alpha, \partial_t \overline{\psi}_\sigma \xi\rangle \le \sup_{\|\alpha\|_{H^1}\le 1} \|\alpha\xi\|_2 \|\partial_t \psi_\sigma\|_2 \\
&\le \sup_{\|\alpha\|_{H^1}\le 1} \|\alpha\|_4 \|\partial_t \psi_\sigma\|_2 \|\xi\|_4 = C\|\partial_t \psi_\sigma\|_2 \|\xi\|_{H^1_{v_\sigma}}\,.
\end{aligned} \tag{4.11}$$

Furthermore, if we use Eq. (2.13) to write explicitly the time derivative

$$\partial_t \psi_\sigma = i\dot{\tilde{\gamma}}\psi_\sigma + \sum_i (-\dot{z}_i)(\nabla_{A_\sigma}\psi_\sigma)_i \tag{4.12}$$

we obtain the following bound on $\|\partial_t \psi_\sigma\|_2$

$$\|\partial_t \psi_\sigma\|_2 \le \|\dot{\tilde{\gamma}}\|_2 + \sum_i |\dot{z}_i|\, \|(\nabla_{A_\sigma}\psi_\sigma)_i\|_2 \le C(\|\dot{\tilde{\gamma}}\|_2 + \sum_i |\dot{z}_i|) \tag{4.13}$$

where again we use the gauge invariance of $\|(\nabla_{A_\sigma}\psi_\sigma)_i\|_2$.

Thus we arrive at the result

$$\|\dot{P}_{v_\sigma}\zeta\|_2 \le C(\|\dot{\tilde{\gamma}}\|_2 + \sum_i |\dot{z}_i|\,)\|\zeta\|_{H^1_{v_\sigma}}\,. \tag{4.14}$$

This completes the proof of Lemma 3. ■

Proof of Lemma 4:

Our goal is to prove the inequality

$$|a_n \dot{\mathbf{z}} + \partial^A_{\mathbf{z}} \mathcal{E}'_\epsilon(v_\sigma)| + \|\dot{\tilde{\gamma}}\|_{H^{1-s}} \le C\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}}\,. \tag{4.15}$$

Starting with $\dot{\tilde{\gamma}}$ we have

$$\begin{aligned}\langle G^{\sigma}_{\delta(x)}, \dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle &= \langle G^{\sigma}_{\delta(x)}, \dot{v}_\sigma\rangle + \langle G^{\sigma}_{\delta(x)}, P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle = \langle G^{\sigma}_{\delta(x)}, \dot{v}_\sigma\rangle \\ &= (-\Delta + |\psi_\sigma(x)|^2)\dot{\tilde{\gamma}}(x) = (K_\sigma \dot{\tilde{\gamma}})(x)\end{aligned} \tag{4.16}$$

since $\langle G^{\mathbf{z}\gamma}_{\delta(x)}, P_v\mathcal{E}'_\epsilon(v)\rangle = Im[\overline{\psi}(\epsilon W(x)\psi)] = 0$. We find the following expression for $\dot{\tilde{\gamma}}$

$$\dot{\tilde{\gamma}}(x) = \int d^2y\, K^{-1}_\sigma(x,y)\langle G^{\sigma}_{\delta(y)}, \dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle \tag{4.17}$$

where,

$$\begin{aligned}&\langle G^{\sigma}_{\delta(y)}, \dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle = \\ &\quad = \left\{Im[\overline{\psi}_\sigma(\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma))_1] - \nabla\cdot(\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma))_2\right\}(y)\,.\end{aligned} \tag{4.18}$$

Here$(\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma))_1$ is the first, complex scalar, component and $(\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma))_2$ is the second, real vector component. Since K^{-1}_σ is a bounded operator from H^{-1-s} to H^{1-s} (uniformly in σ) we have

$$\begin{aligned}\|\dot{\tilde{\gamma}}\|_{H^{1-s}} &= \|K^{-1}_\sigma\langle G^{\sigma}_{\delta(y)}, \dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle\|_{H^{1-s}} \\ &\le C\|\langle G^{\sigma}_{\delta(y)}, \dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle\|_{H^{-1-s}}\,.\end{aligned} \tag{4.19}$$

However, using Eq. (4.18) we obtain

$$\begin{aligned}&\|\langle G^{\sigma}_{\delta(y)}, \dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\rangle\|_{H^{-1-s}} = \\ &\quad = \|Im[\overline{\psi}_{\mathbf{z}0}e^{-i\gamma}(\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma))_1] - \nabla\cdot(\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma))_2\|_{H^{-1-s}} \\ &\qquad \le C\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}}\end{aligned} \tag{4.20}$$

which is the desired bound. As for $\dot{z}_i$, we have

$$|a_n\dot{z}_i + \partial^A_{z_i}\mathcal{E}_\epsilon(v_\sigma)| = |\langle T^\sigma_i, \dot{v}_\sigma + \mathcal{E}'_\epsilon(v_\sigma)\rangle| \le \|T^\sigma_i\|_{H^s_{v_\sigma}}\|\dot{v}_\sigma + P_{v_\sigma}\mathcal{E}'_\epsilon(v_\sigma)\|_{H^{-s}_{v_\sigma}}\,. \tag{4.21}$$

Obviously $\|T^\sigma_i\|_{H^s_{v_\sigma}} = \|T^{\mathbf{z}0}_i\|_{H^s} < C$. From the last estimate, together with Eq. (4.19) and Eq. (4.20), follows Eq. (4.15). ■

Proof of Lemma 1:

For any $v \in M_{as}$, the tangent space $T_v M_{as}$ is spanned by zero eigenvectors of $L_{0;v}$. Hence, since P_v is a projection on $T_v M_{as}$ we have

$$P_v L_{0;v} = L_{0;v} P_v = 0\,. \tag{4.22}$$

Furthermore, by assumption the potential W is bounded

$$\sup_{x \in R^2} |W(x)| = C\,. \tag{4.23}$$

Take any vector $\zeta = (\xi, F) \in T_v X^{(n)}$. The above observation leads to the following estimate

$$\|P_v L_{\epsilon;v} \zeta\|_2 = \|P_v (L_{0;v} + \epsilon W)\zeta\|_2 \leq \epsilon \|W\zeta\|_2 \leq \epsilon C \|\zeta\|_2\,. \tag{4.24}$$

■

Proof of Lemma 2:

We set out to prove the following inequality

$$\|P_v N_v(\zeta)\|_{H^{-s}_{v_\sigma}} \leq C[\|\zeta\|^2_{H^1_{v_\sigma}} + \|\zeta\|^3_{H^1_{v_\sigma}}]\,. \tag{4.25}$$

The expression for $N_v(\zeta)$ is given in Appendix A, Eq. (A.3), where $\zeta = (\xi, F)$. The estimates of the terms in $N_v(\zeta)$ which do not include derivatives are straightforward. For example, denoting $\tilde{N}_1 = (0, 2Re(\overline{\psi} F \xi))$ we obtain

$$\begin{aligned} \|\tilde{N}_1\|_{H^{-s}_{v_\sigma}} = 2\|Re(\overline{\psi} F \xi)\|_{H^{-s}_{v_\sigma}} \leq 4\|Re(\overline{\psi} F \xi)\|_{L^2_{v_\sigma}} = 4\|Re(\overline{\psi} F \xi)\|_2 \\ \leq C\|\zeta\|_4^2 \leq C\|\zeta\|^2_{H^1_{v_\sigma}}\,. \end{aligned} \tag{4.26}$$

The vector $\tilde{N}_2 = (0, Im(\overline{\xi} \nabla_A \xi))$ is an example for a term which is more difficult to estimate. An estimate of this term is given in Section 4. The other problematic terms in $N_v(\zeta)$ are estimated in the same manner. ■

Proof of Lemma 5:

We want to prove the inequality

$$\|N_v(\zeta)\|_2 \leq C\|\zeta\|_{H^2_{v_\sigma}} (\|\zeta\|_{H^1_{v_\sigma}} + \|\zeta\|^2_{H^1_{v_\sigma}}) \tag{4.27}$$

Again the terms not containing any derivatives do not present any problems (see for example Eq. (4.26)). For the vector $\tilde{N}_2 = (0, Im(\overline{\xi}\nabla_{A_\sigma}\xi))$ we have

$$\begin{aligned}\|\tilde{N}_2\|_2 \leq \|\overline{\xi}\nabla_{A_\sigma}\xi\|_2 \leq \|\overline{\xi'}\nabla_{A^{(n)}}\xi'\|_2 \leq \|\xi'\|_\infty \|\nabla_{A^{(n)}}\xi'\|_2 \\ \leq C\|\xi'\|_{H^2}\|\xi'\|_{H^1} = C\|\xi\|_{H^2_{v_\sigma}}\|\xi\|_{H^1_{v_\sigma}}\end{aligned} \tag{4.28}$$

where $\xi' = e^{-i\gamma}\xi$. Other terms containing derivatives are estimated in a similar way. ■

Proof of Lemma 6:

We want to prove the inequalities

$$\|\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta\|_2 \leq C\|\zeta\|_{H^2_{v_\sigma}} \tag{4.29}$$

and

$$\|\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta\|_{H^{-1}_{v_\sigma}} \leq C\|\zeta\|_{H^1_{v_\sigma}} . \tag{4.30}$$

Since $L_{\epsilon;v_\sigma}$ is a second order linear operator no difficulties arise and the two inequalities are proved by strightforward calculations. We just note that

$$\|\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta\|_2 = \|G_{-\gamma}\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta\|_2 = \|\overline{P}_{v_{\mathbf{z}0}} L_{\epsilon;v_{\mathbf{z}0}}\zeta'\|_2 \tag{4.31}$$

and

$$\|\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta\|_{H^{-1}_{v_\sigma}} = \|G_{-\gamma}\overline{P}_{v_\sigma} L_{\epsilon;v_\sigma}\zeta\|_{H^{-1}} = \|\overline{P}_{v_{\mathbf{z}0}} L_{\epsilon;v_{\mathbf{z}0}}\zeta'\|_{H^{-1}} \tag{4.32}$$

where $\zeta' = e^{-i\gamma}\zeta$. Since the gauge function γ does not appear in the operators on the r.h.s. of Eq. (4.31) and Eq. (4.32) one obtains by a straightforward calculation

$$\|\overline{P}_{v_{\mathbf{z}0}} L_{\epsilon;v_{\mathbf{z}0}}\zeta'\|_2 \leq C\|\zeta'\|_{H^2} = \|\zeta\|_{H^2_{v_\sigma}}$$

and

$$\|\overline{P}_{v_{\mathbf{z}0}} L_{\epsilon;v_{\mathbf{z}0}}\zeta'\|_{H^{-1}} \leq C\|\zeta'\|_{H^1} = C\|\zeta'\|_{H^1_{v_\sigma}} .$$

■

Proof of Lemma 7: We want to prove the inequality

$$\|\partial_{v_\sigma} R_{v_\sigma}(\zeta)\|_{H^1_{v_\sigma}} \leq \|\zeta\|^2_{H^2_{v_\sigma}}\|\zeta\|_{H^1_{v_\sigma}} .$$

By derivation of the expression for $R_v(\zeta)$ (Eq. (A.5)) we obtain, for $\zeta = (\xi, F)$

$$\partial_v R_v(\zeta) = \left((F^2 + \frac{\lambda}{2}|\xi|^2)\xi \ , F|\xi|^2 \right) . \tag{4.33}$$

We want to estimate the norm $\|\partial_{v_\sigma} R_{v_\sigma}(\zeta)\|_{H^1_{v_\sigma}}$. Note first that

$$G_{-\gamma}\partial_{v_\sigma} R_{v_\sigma}(\zeta) = \left((F^2 + \frac{\lambda}{2}|\xi'|^2)\xi' \ , F|\xi'|^2 \right) = \partial_{v_\sigma} R_{v_\sigma}(\zeta')$$

where $\xi' = e^{-i\gamma}\xi$, $\zeta' = G_{-\gamma}\zeta$. The estimate of the L^2 norm (or $L^2_{v_\sigma}$ norm which is the same) is straightforward and we get

$$\|\partial_{v_\sigma} R_{v_\sigma}(\zeta)\|_2^2 = \|(F^2 + \frac{\lambda}{2}|\xi'|^2)\xi'\|_2^2 + \|F|\xi'|^2\|_2^2 \leq C\|\zeta'\|^6_{H^1} = C\|\zeta\|^6_{H^1_{v_\sigma}} .$$

Let $\tilde{F} = F_1 + iF_2$ be the complexification of F. We have

$$\begin{aligned} \|\nabla(\partial_v R_v(\zeta'))\|_2^2 &= \|\nabla((|\tilde{F}|^2 + \frac{\lambda}{2}|\xi'|^2)\xi')\|_2^2 + \|\nabla(\tilde{F}|\xi'|^2)\|_2^2 = \\ &= \|2Re[(\nabla\tilde{F})\overline{\tilde{F}}]\xi' + |\tilde{F}|^2(\nabla\xi') + \lambda(\nabla\xi')|\xi'|^2 + \frac{\lambda}{2}(\nabla\overline{\xi'})\xi'^2\|_2^2 \\ &\quad + \|(\nabla\tilde{F})|\xi'|^2 + 2\tilde{F}\, Re[\overline{\xi'}\nabla\xi']\|_2^2 . \end{aligned} \tag{4.34}$$

All the terms on the r.h.s. of Eq. (4.34) are handled in the same manner. For example:

$$\begin{aligned} \|(\nabla\tilde{F})\overline{\tilde{F}}\,\xi')\|_2 &\leq \|\tilde{F}\|_\infty\|\xi'\|_\infty\|\nabla\tilde{F}\|_2 \\ &\leq \|\tilde{F}\|_{H^2}\|\xi'\|_{H^2}\|\tilde{F}\|_{H^1} \leq \|\zeta'\|^2_{H^2}\|\zeta'\|_{H^1} = \|\zeta\|^2_{H^2_{v_\sigma}}\|\zeta\|_{H^1_{v_\sigma}} . \end{aligned}$$

■

Proof of Lemma 8:

First we use Eq. (4.22) and the linear stability theorem in order to arrive at the result that for any $\zeta_v \in T_v X^{(n)}$ we have

$$\begin{aligned} &\|L_{0;v}\zeta_v\|_2^2 = \langle L_{0;v}\zeta_v, L_{0;v}\zeta_v\rangle = \langle L^{1/2}_{0;v}\zeta_v, L_{0;v}\, L^{1/2}_{0;v}\zeta_v\rangle = \\ &= \langle L^{1/2}_{0;v}\zeta_v, \overline{P}_v L_{0;v}\overline{P}_v\, L^{1/2}_{0;v}\zeta_v\rangle = \langle \overline{P}_v L^{1/2}_{0;v}\zeta_v, L_{0;v}\,\overline{P}_v L^{1/2}_{0;v}\zeta_v\rangle \geq \nu\|\overline{P}_v L^{1/2}_{0;v}\zeta_v\|_2^2 . \end{aligned} \tag{4.35}$$

In addition, using Eq. (4.22) again, we have

$$\|P_v L_{0;v}^{1/2}\zeta_v\|_2^2 = \langle P_v L_{0;v} L_{0;v}^{-1/2}\zeta_v, P_v L_{0;v}^{1/2}\zeta_v\rangle = 0\,. \tag{4.36}$$

From Eq. (4.35), Eq. (4.36) and the linear stability theorem we deduce that

$$\langle L_{0;v}\zeta_v, L_{0;v}\zeta_v\rangle \geq \nu\langle \zeta_v, L_{0;v}\zeta_v\rangle \geq \nu^2\|\zeta_v\|_{H^1}^2\,. \tag{4.37}$$

We note that $\|(L_{0;v}+\Delta)\zeta_v\|_2 \leq C_1\|\zeta_v\|_{H^1} + \|\nabla\nabla F_v\|_2$, where $\zeta_v = (\xi_v, F_v)$. Furthermore, since $\zeta_v \in (T_v M_{as})^\perp$, it satisfies the gauge condition $\mathrm{Im}\,(\overline{\psi}_v\xi_v) - \nabla F_v{=}0$. We conclude that $L_{0,v}+\Delta$ is a bounded operator from H^1 to L^2, i.e. $\|L_{0,v}+\Delta\|_{H^1\to L^2} \leq C_1$. we have

$$\begin{aligned}\|\Delta\zeta\|_2^2 = \|(L_{0;v}+\Delta)\zeta - L_{0;v}\zeta\|_2^2 &\leq 2(\|(L_{0;v}+\Delta)\zeta\|_2^2 + \|L_{0;v}\zeta\|_2^2)\\ &\leq 2(C_1\|\zeta\|_{H^1}^2 + \|L_{0;v}\zeta\|_2^2)\end{aligned}$$

and so

$$\frac{1}{2}\|\Delta\zeta\|_2^2 - C_1\|\zeta\|_{H^1}^2 \leq \langle L_{0;v}\zeta, L_{0;v}\zeta\rangle\,.$$

Then, for any $0<\delta<1$ we have

$$\begin{aligned}\langle L_{0,v}\zeta_v, L_{0,v}\zeta_v\rangle &= \delta\langle L_{0,v}\zeta_v, L_{0,v}\zeta_v\rangle + (1-\delta)\langle L_{0,v}\zeta_v, L_{0,v}\zeta_v\rangle\\ &\geq \frac{\delta}{2}\|\Delta\zeta_v\|_2^2 + [(1-\delta)C_2 - \delta C_1]\|\zeta_v\|_{H^1}^2\,.\end{aligned} \tag{4.38}$$

Choosing δ small enough the second term on the r.h.s. of Eq. (4.38) is positive and we get

$$\langle L_{0,v}\zeta_v, L_{0,v}\zeta_v\rangle \geq C\|\zeta_v\|_{H^2}^2\,. \tag{4.39}$$

In addition to this basic result we have

$$\begin{aligned}\|L_{0;v}\zeta_v\|_2^2 = \|\overline{P}_v L_{0;v}\zeta_v\|_2^2 &= \|\overline{P}_v(L_{\epsilon;v}\zeta_v - \epsilon W\zeta_v)\|_2^2\\ &\leq 2(\|\overline{P}_v L_{\epsilon;v}\zeta_v\|_2^2 + \epsilon^2\|W\|_\infty^2\|\zeta_v\|_2^2)\end{aligned} \tag{4.40}$$

which leads to

$$\langle \overline{P}_v L_{\epsilon;v}\zeta_v, \overline{P}_v L_{\epsilon;v}\zeta_v\rangle \geq \left(\frac{1}{2} - \epsilon^2\|W\|_\infty^2\nu^2\right)\langle L_{0;v}\zeta, L_{0;v}\zeta_v\rangle\,. \tag{4.41}$$

The proof of Lemma 8 is completed by the following simple observation

$$\begin{aligned}\langle \overline{P}_v L_{\epsilon;v}\zeta_v, \overline{P}_v L_{\epsilon;v}\zeta_v\rangle = \langle G_{-\gamma}\overline{P}_v L_{\epsilon;v}\zeta_v, G_{-\gamma}\overline{P}_v L_{\epsilon;v}\zeta_v\rangle &=\\ = \langle \overline{P}_{v_{\mathbf{z}0}} L_{\epsilon;v_{\mathbf{z}0}}\zeta'_v, \overline{P}_{v_{\mathbf{z}0}} L_{\epsilon;v_{\mathbf{z}0}}\zeta'_v\rangle &\end{aligned} \tag{4.42}$$

where $\zeta_v' = G_{-\gamma}\zeta_v$ and we have $P_{v_{\mathbf{z}0}}\zeta_v' = 0$. Eq. (4.39), Eq. (4.41) and Eq. (4.42) imply the desired result. ■

Proof of Lemma 9:

The inequality in Lemma 9, Eq. (3.23), follows from Eq. (4.37), Eq. (4.41) and the following simple estimate

$$\begin{aligned}\langle\zeta_v, L_{\epsilon;v}\zeta_v\rangle = \langle\zeta_v, L_{0;v}\zeta_v\rangle + \epsilon\langle\zeta_v, W\zeta_v\rangle &\leq \langle\zeta_v, L_{0;v}\zeta_v\rangle + \epsilon\|W\|_\infty\|\zeta_v\|_2^2 \\ &\leq (1+\epsilon\|W\|_\infty\nu)\langle\zeta_v, L_{0;v}\zeta_v\rangle\,.\end{aligned} \tag{4.43}$$

■

Appendix A - Taylor expansions of $\mathcal{E}_\epsilon(u)$ and $\mathcal{E}_\epsilon'(u)$:

In the proof of Theorem 1 above we frequently refer to the Taylor expansion of $\mathcal{E}_\epsilon'(u)$

$$\mathcal{E}_\epsilon'(u) = \mathcal{E}_\epsilon'(v) + L_{\epsilon;v}\zeta + N_{\epsilon;v}(\zeta) \tag{A.1}$$

A strightforward calculation provides us with explicit expressions for the linear and nonlinear terms in ζ. Denoting $\zeta = (\xi, F)$, we have

$$L_{\epsilon;u}\zeta = \begin{pmatrix} [-\Delta_A + \frac{\lambda}{2}(2|\psi|^2 - 1)]\xi + \frac{\lambda}{2}\psi^2\overline{\xi} + i[2\nabla_A\psi + \psi\nabla]F + \epsilon W\xi \\ \\ Im([\overline{\nabla_A}\psi - \overline{\psi}\nabla_A]\xi) + (-\Delta + \nabla\nabla + |\psi|^2)F \end{pmatrix} \tag{A.2}$$

and

$$N_{\psi,A}(\zeta) = \begin{pmatrix} i(2\nabla_A\xi + \xi\nabla)F + F^2\psi + F^2\xi + \frac{\lambda}{2}(\overline{\psi}\xi^2 + 2|\xi|^2\psi + |\xi|^2\xi) \\ \\ 2Re(\overline{\psi}F\xi) - Im(\overline{\xi}\nabla_A\xi) + F|\xi|^2 \end{pmatrix}. \tag{A.3}$$

In addition we refer at several points to the Taylor expansion of the energy functional $\mathcal{E}_\epsilon'(u)$. Letting $u = v + \zeta$ we have

$$\mathcal{E}_\epsilon(u) = \mathcal{E}_\epsilon(v) + \langle\zeta, \mathcal{E}_\epsilon'(v)\rangle + \frac{1}{2}\langle\zeta, L_{\epsilon;v}\zeta\rangle + R_v(\zeta) \tag{A.4}$$

where $L_{\epsilon;v}$ is given in Eq. (A.2) and the remainder $R_v(\zeta)$ (containing all terms which are higher then quadratic in ζ) is given by

$$R_v(\zeta) = \int\left\{(F^2 + \frac{\lambda}{2}|\xi|^2)\mathrm{Re}\,[\overline{\xi}\psi] - F\cdot\mathrm{Im}\,[\overline{\xi}\nabla_A\xi] + \frac{1}{2}(F^2|\xi|^2 + \frac{\lambda}{4}|\xi|^4)\right\}d^2x \tag{A.5}$$

Acknowledgments

The work of Y. Strauss was supported by NSERC Grant N7109, THE ISRAEL SCIENCE FOUNDATION (Grant no. 188/02) and the Edmund Landau Center for Research in Mathematical Analysis and Related areas, sponsored by the Minerva Foundation (Germany). The work of I.M. Sigal was supported by NSF Grant DMS-040026.

References

[A] Abrikosov A.A.: On the magnetic properties of superconductors of the second group. Soviet Physics JETP **5** 1174-1182 (1957).

[ABP] Andre N. and Bauman P. and Phillips D.: Vortex pinning with bounded fields for the Ginzburg-Landau equation. Ann. Ins. Henri Poincaré Anal. Non Linéaire **20** no. 4 705-729 (2003).

[AS] Andre N. and Shafrir I.: Asymptotic behavior of minimizers for the Ginzburg-Landau functional with weight. Arch. Rational Mech. Anal. **142**, no. 1, 45-73 75-98 (1998).

[ASaSe] Aftalion A. and Sandier E. and Serfaty S.: Pinning phenomena in the Ginzburg-Landau model of superconductivity. J. Math. Pures Appl. (9) **80** no. 3 339-372 (2001).

[B] Bethuel F.: Vortices in Ginzburg-Landau equations, ICM 98, Documenta Mathematica **III** 11-19.

[BBH] Bethuel F. and Brezis H. and Hélein F.: *Ginzburg-Landau Vortices*, Birkhäuser (1994).

[BC] Berger M.S. and Chen Y.Y.: Symmetric vortices for the Ginzburg-Landau equations of superconductivity, and the nonlinear desingularization phenomenon. J. Func. Anal. **82** 259-295 (1989).

[BFGLV] Blatter G. and Feigeĺman M.V. and Geshkenbein V.B. et. al.: Vortices in high temprature superconductors. Rev. Modern Phys. **66** 1125 (1994).

[DG] Du Q. and Gunzburger M.D.: A model for superconducting thin films having variable thickness. Phys. D **69** 215-231 (1993).

[DS] Demoulini S. and Stuart D.: Gradient flow of the superconducting Ginzburg-Landau functional on the plane. Comm. Anal. Geom. **5** 121-198 (1997).

[E] E W.: Dynamics of vortices in Ginzburg-Landau thories with applications to superconductivity. Physica D **77** 383-404 (1994).

[G1] Gustafson S.: Some mathematical problems in the Ginzburg-Landau theory of superconductivity. Ph.D. thesis, Univ. Toronto (1999).

[G2] Gustafson S.: Dynamical stability of magnetic vortices, Nonlinearity **15** 1717-1728 (2002).

[GE] Gork'ov L.P. and Eliashberg G.M.: Generalization of the Ginzburg-Landau equations for non-stationary problems in the case of alloys with paramagnetic impurities. Soviet Physics JETP **27** no. 2 328-334 (1968).

[GS1] Gustafson S. and Sigal I.M.: The stability of magnetic vortices. Comm. Math. Phys. **212** 257-275 (2000).

[GS2] Gustafson S. and Sigal I.M.: Effective dynamics of magnetic vortices. Preprint, arXiv:math.AP/0312438 (2003).

[JS] Jerard R. and Soner M.: Dynamics of Ginzburg-Landau vortices. Arch. Rational Mech. Anal. **142** no. 2, 99-125 (1998).

[JT] Jaffe A. and Taubes C.: *Vortices and monopoles. Structure of static gauge theories*. Progr. in Phys. Vol. 2 Birkhäuser, Boston, MA 1980.

[L] Lin F.H.: Some dynamical properties of Ginzburg-Landau vortices. Comm. Pure. Appl. Math. **49** no. 4 323-359 (1996).

[N] Neu J.: Vortices in complex scalar fields. Physica D **43** 385-406 (1990).

[P] Plohr B. Ph.D. thesis, Princeton (1978).

[PaRi] Pacard F. and Rivière T.: *Linear and Nonlinear Aspects of Vortices*, Birkhäuser, Boston (2000).

[PeRu] Peres L. and Rubinstein J.: Vortex dynamics in $U(1)$ Ginzburg-Landau models. Physica D **64** 299-309 (1993).

[Rit] Ritter W.G.: Gauge theory: instantons, monopoles and moduli spaces. Preprint, arXiv:math-ph/0304026 v1 (2003).

[Riv] Riviére T.: Ginzburg-Landau vortices: the static model. In *Séminaire Bourbaki*, Vol. 1999/2000, Astérisque No. 276 73-103 (2002).

[Rub1] Rubinstein J.: On the equlibrium position of Ginzburg-Landau vortices. Z. Angew. Math. Phys. **46** no. 5 739-751 (1995).

[Rub2] Rubinstein J.: Six lectures on superconductivity. In *boundaries, interfaces and Transitions* (Banff. AB, 1995) CRM Proc. Lecture Notes, Vol. 13, Amer. Math. Soc., Providence, RI, pp. 163-184 (1998).

[RS] Rubinstein J. and Sternberg P.: On the slow motion of vortices in the Ginzburg-Landau heat flow. SIAM J. Math. Anal. **26** no. 6 1452-1466 (1995).

[ST1] Sigal I.M. and Ting F.: Pinning of magnetic vortices by an external potential. Algebra i Analiz **16** no. 1, 239-268 (2004).

[ST2] Sigal I.M. and Ting F.: Stability of pinned vortices (to appear).

[T] Tinkham M.: *Introduction to superconductivity.* McGraw-Hill, New York (1975).