Fokker-Planck Equations as Scaling Limits of Reversible Quantum Systems

Francois Castella
CNRS et IRMAR
Université de Rennes 1
Campus de Beaulieu
35042 Rennes Cedex, France

László Erdős
School of Mathematics
Georgiatech
Atlanta GA-30332, USA

Florian Frommlet and Peter A. Markowich
Institut für Mathematik
Universität Wien
Boltzmanngasse 9
A-1090 Wien, Austria

September 30, 1999

Abstract

We consider a quantum particle moving in a harmonic exterior potential and linearly coupled to a heat bath of quantum oscillators. Caldeira and Leggett [5] have derived the Fokker-Planck equation with friction for the Wigner distribution of the particle in the large temperature limit, however their (nonrigorous) derivation was not free of criticism, especially since the limiting equation is not of Lindblad form. In this paper we recover the correct form of their result in a rigorous way. We also point out that the source of the diffusion is physically unnatural under their scaling. We investigate the model at a fixed temperature and in the large time limit, where the origin of the diffusion is a cumulative effect of many resonant collisions. We obtain a heat equation with a friction term for the radial process in phase space and we prove the Einstein relation in this case.

Keywords: Fokker-Planck equation, Wigner distribution, scaling limit, coupled harmonic oscillators.

Contents

1 Introduction **2**

2 Source of diffusion in various kinetic models **7**
2.1 Eliminating the semiclassical parameter 7
2.2 Diffusion in the original model 8
2.3 Diffusion from resonances in the scaling limit 11
2.4 Comparison of the three models 11
2.5 Derivation of Fokker-Planck equation via Boltzmann equation 13

3 The assumptions of the Caldeira-Leggett model **15**
3.1 Heat bath and its frequencies 15
3.2 High temperature 16
3.3 Linear coupling 17

4 Preliminary results **19**
4.1 The Wigner formalism 19
4.2 Solution by characteristics 21
4.3 The thermodynamic limit 22
4.4 Digression on stochastic integrals 24

5 The Fokker-Planck equation from the original Caldeira-Leggett model **28**
5.1 Evolution without friction 28
5.2 Computing the dynamics of the test-particle when the memory vanishes 29
5.3 The Caldeira-Leggett limits: obtaining the Fokker-Planck equation 31

6 Scaling limit at high temperature: the frictionless heat equation **34**
6.1 Large space/time convergence of the Wigner distribution 34

7 Heat equation with friction at finite temperature **40**
7.1 A priori bounds and continuity results 41
7.2 Transport equation before scaling limits 44
7.3 Obtaining diffusion from scaling limit 44
7.4 Derivation of the limiting equation 46

1 Introduction

In [5], Caldeira and Leggett introduced a Hamiltonian for a quantum system of a test-particle coupled to an abstract reservoir. The Schrödinger equation for the evolution of the quantum state can be equivalently written as a kinetic (phase-space) equation for the associated Wigner distribution of the test particle-reservoir system. The goal of [5] was to derive (formally) a Fokker-Planck equation for the Wigner distribution of the test-particle by taking various limits which we explain below and by "tracing out" the reservoir coordinates. The Fokker-Planck equation represents an irreversible collisional evolution with a diffusive term, while the Schrödinger equation is reversible. Hence this derivation was expected to shed some light on the origin of diffusion in the evolution of a small system coupled to an infinite reservoir. Caldeira and Leggett used a Feynman path integral approach which has no rigorous mathematical justification (despite its great successes in formal computations). More importantly, several other steps in their derivation admittedly lack precision, both from the mathematical and from the physical point of view.

Starting from this observation, the aim of the present paper is threefold. First, we discuss the origin of the diffusion and the physical meaning of the different scalings and limiting procedures introduced in [5]. In particular, we point out in Section 3 that the model introduced in [5] is not physically satisfactory in the regime where they let the diffusion appear. Second, in Sections 4 and 5 we present a mathematically rigorous derivation of the frictionless Fokker-Planck equation from the model introduced in [5]. Third, in Sections 6 and 7 we show how to recover other types of diffusive behaviour from the Caldeira-Leggett Hamiltonian, using different, more realistic scalings and limiting procedures.

We point out that [5] heavily relies on the use of ideas from Feynman, Hibbs, and Vernon [22], [23]. In particular Feynman and Vernon [23] considered a system of the form { test "particle" (A) + reservoir (R) }. The Hamiltonian is $H_A + H_R + H_I$, where H_A is the free Hamiltonian for the test-particle, H_R is the free Hamiltonian for the reservoir, and H_I is the interaction Hamiltonian. They integrated out the reservoir variables, i.e. they computed the time evolution of the wave function of the test-particle itself, given by $Tr_R\{\exp(ith^{-1}(H_A + H_R + H_I))\}$, where Tr_R is the partial trace on the Hilbert space of the reservoir and h is the Planck constant. Feynman path integral formalism was used which is particularly powerful when the total Hamiltonian, or at least $H_R + H_I$, is quadratic (in particular the interaction H_I must be linear both in the test-particle and in the reservoir variables). In this case, one is led to computing Gaussian integrals, which, in principle, are explicit. The difficulty stems from the large (infinite) number of variables.

In this context [5] takes place. Namely, [5] introduces the following Hamiltonian,

$$\begin{aligned} H_{CL} &= H_A + H_R + H_I \\ &= \left(-\frac{h^2}{2M}\Delta_x + V(x)\right) + \sum_{j=1}^{N\Omega}\left(-\frac{h^2}{2}\Delta_{R_j} + \frac{1}{2}\omega_j^2|R_j|^2\right) + \frac{1}{\sqrt{N}}\left(\sum_{j=1}^{N\Omega} C_j R_j\right)\cdot x \ . \end{aligned} \tag{1.1}$$

The first term of (1.1) represents the Hamiltonian of the test-particle with mass M where $x \in \mathbb{R}^d$ denotes the test-particle position in dimension d. The abstract reservoir here is a set of finitely many (say $N\Omega$, which is assumed to be integer) independent oscillators written in normal variables $R_j \in \mathbb{R}^d$, having frequencies $\omega_j \in [0, \Omega]$. Here Ω is the maximum frequency of the oscillators and N is the number of oscillators per unit frequency. The typical case is the uniform frequency distribution: $\omega_j = \frac{j}{N}$. The coupling is linear in x and the R_j's, with coupling coefficients given by the C_j's. The normalization factor $N^{-1/2}$ simply stems from the central limit theorem, since, roughly speaking, the variables R_j's become independent random variables with vanishing expectation in the thermodynamic limit $N \to \infty$. The operator H acts on the Hilbert space $L_x^2(\mathbb{R}^d) \otimes \left(\bigotimes_{j=1}^{N\Omega} L_{R_j}^2(\mathbb{R}^d)\right)$. The authors of [5] consider only $d = 1$ for simplicity, as we shall do as well, but the method extends to any dimension.

Caldeira-Leggett assume that the reservoir is initially in thermal equilibrium at inverse temperature β, i.e. the initial density matrix of the system $A + R$ is given by,

$$\rho^0 = \rho_A^0 \otimes \exp\left(-\beta H_R\right) , \tag{1.2}$$

where ρ_A^0 is the initial state of the test-particle. Finally, they choose the coupling coefficients,

$$C_j := \lambda\omega_j \ , \tag{1.3}$$

with some β-dependent coupling parameter λ, and specify,

$$\lambda = \lambda_0 \beta^{1/2} \ , \tag{1.4}$$

for some fixed λ_0. Note that equation (1.3) could be interpreted mathematically as a frequency dependent coupling, whereas in physical applications the coupling is typically frequency independent. This apparent difficulty is actually an artefact; the ω_j prefactor in (1.3) stems from the three-dimensionality of the underlying phonon or photon bath implicitly described by the abstract reservoir in (1.1) (see Section 3.1).

Remark. Instead of uniformly spaced oscillator frequencies $\omega_j = \frac{j}{N}$, it is sufficient to assume that the frequency distribution tends, in the thermodynamic limit ($N \to \infty$), to a uniform distribution on $[0, \Omega]$ with density, say, c, i.e.

$$\lim_{N\to\infty} \frac{1}{N}\sum_{k=1}^{N\Omega} h(\omega_k) = c\int_0^\Omega h(\omega)d\omega, \quad \forall h \in C[0,\Omega] \ . \tag{1.5}$$

Without loss of generality $c = 1$ can be assumed because changing c to 1 is equivalent to changing $\lambda \to \sqrt{c}\lambda$ (see Section 3.1).

Now the main steps of [5] are the following:

• *First*, using that $H_I + H_R$ is quadratic and relying on Feynman path integrals, Caldeira and Leggett explicitly compute the evolution of the test-particle after tracing out the reservoir variables. The evolution equation of the test-particle involves a diffusive forcing term and a memory term (friction), the latter being non-local in time (see (2.2) below, as well as (4.14)). These terms translate the effect of the evolution of the reservoir on the test-particle. It is very standard in this context that integrating out the reservoir variables gives rise to a non-Markovian evolution for the test-particle, despite that the evolution of the full system is Markovian.

• *Second*, they perform the thermodynamical limit where the number of oscillators (per unit frequency) in (1.1) becomes infinite ($N \to \infty$). More precisely, they let the oscillator ensemble tend to a continuous distribution of oscillators with uniform density in some finite range of frequency $[0, \Omega]$. This step operates with apparently ill-defined objects, but it can easily be made rigorous as we will show.

• *Third*, they consider the semiclassical limit $h \to 0$, they perform the limit $\Omega \to \infty$, i.e. the frequency range becomes infinite (removing ultraviolet cutoff), and they let the inverse temperature β go to zero. The value $\beta\Omega$ is responsible for a potential renormalization (frequency shift).

These last two limits allow them to eliminate all the non-Markovian effects. Caldeira and Leggett state the Fokker Planck equation

$$\partial_t w + v \cdot \nabla_x w - \nabla_x V(x) \cdot \nabla_v w = \gamma \nabla_v (vw) + \sigma \Delta_v w \tag{1.6}$$

for the particle's Wigner distribution $w = w(t, x, v)$, which can be interpreted as a phase space (quasi)density, as a result of their asymptotic procedures. The friction coefficient γ is given as $\gamma = \sigma\beta/M$, which is the well-known Einstein's relation between friction, diffusivity and inverse temperature. Depending on the order of limits, V may be modified to an effective potential $V_{eff}(x) = V(x) - \omega_R^2 x^2$, where ω_R is called the frequency shift.

This type of equation is also known under the name of "Quantum Brownian motion", or "Quantum Langevin equation", and received a large interest in the context of interaction between light and matter (see, e.g. [11]).

We mention that the idea of formally deriving Fokker-Planck-like equations from a reservoir of oscillators with linear coupling has been exploited by many authors, e.g. [6], [14, 15], [12], [29], [46] (see also [16] for comments on this equation and the relationship with questions of decoherence). These authors use similar scalings as [5]. In particular, in [14, 15], [46], [29], corrections to (1.6) are derived when the temperature is large but finite, and these equations involve both a diffusive term in velocity and friction terms in space and velocity. Mathematically rigorous work on these types of models is slightly less abundant. A rigorous operator-algebraic approach is given in [13], and a path-integral approach is found in [9]. A similar model has also been used in the program of Jakšić and Pillet to study thermal relaxation with spectral methods (see [30] and references therein). Recently an analogous system with an extra white-noise is studied in [24]. Under different scalings Arai derives ballistic behaviour for the test-particle [2]. In a different context and with different scaling assumptions than [5] and others, but still with the assumption of linear coupling, we also mention [11]. The key assumption in all these papers is that the test-particle is linearly coupled to the infinite bath of harmonic oscillators, which gives rise to Gaussian computations, and many quantities of interest become explicitly computable. This certainly explains at least part of the interest that these kinds of models have received.

The paper by Caldeira and Leggett raises several questions which have to be addressed. The most serious is that the limiting equation (1.6) is not of Lindblad form (see [1], [15], [37]), which is a generic condition for quantum systems to preserve the complete positivity of the density operator along the evolution. Recall that the true quantum evolution preserves this property. This shortcoming is closely related to the fact, that the equation itself contains β (as the ratio of γ and σ), while $\beta \to 0$ limit was actually used along its derivation. This is not just a mathematical inconsistency. Either the friction term should be negligible compared to the diffusion term in (1.6) if $\beta \to 0$ limit is really taken; or there should be an extra term in the equation if β is thought of as a small but nonzero number. In this latter case this extra term should restore the Lindblad form of the equation, and it is not clear why this term could be considered negligible compared to the friction. The confusion probably comes from the unspecified order of limits.

The second most important question is the physical meaning of the system modelled and the limits taken. Indeed, [5] relies on the assumptions that **1)** the coupling C_j is linear in ω with ω ranging over the full interval $[0, \infty]$, i.e. high forcing frequencies are needed ($\Omega \to \infty$); **2)** the weak coupling is proportional to $\beta^{1/2}$, with $\beta \to 0$; **3)** the test-particle is *linearly* coupled to the reservoir of *oscillators*. We remark that none of these assumptions can be taken for granted on physical grounds, in fact quite the opposite, they pose strong restrictions on the applicability

of the result.

Finally, from mathematical point of view, it is desirable to eliminate the nonrigorous steps in the original derivation; especially since the proper order of limits actually does influence the form of the limiting equation. In addition, the systematic use of the Feynman path integral should be avoided in a rigorous proof, since it is a mathematically undefined.

The present paper has five parts:

a) In Section 2 we explain that the origin of the diffusion in the original Caldeira-Leggett model is the artificial $\Omega \to \infty$ limit. Then we explain how to modify the model to obtain diffusion via a more realistic mechanism using scaling limit. We also explain how these derivations are related to other derivations of the Fokker-Planck equation via the Boltzmann equation.

b) In Section 3, we discuss the physical implications of the assumptions and scalings considered in [5].

c) In Section 5, we present a rigorous mathematical convergence result for the model introduced in [5]. Our approach is very elementary and physically transparent.

d) In Section 6, we show that one can also recover a diffusive non-kinetic behaviour (frictionless heat equation) from the Caldeira-Leggett Hamiltonian using scaling limit and without assuming infinite frequency range.

e) In Section 7, under a different scaling limit, we derive a Fokker-Planck equation with friction but without convective terms. The temperature is finite. Einstein relation is valid in a modified form which takes into account the ground state quantum fluctuations of the heat bath.

Our main results are Theorem 5.1, 6.1 and 7.1.

Remark. The equation derived in Section 5 is of Lindblad form (see [1]). Since there is no rescaling in the variables, one can reconstruct the quantum (restricted) density matrix from the evolved Wigner distribution, hence the equation must preserve the positivity of the corresponding density matrix. The Wigner distribution itself is typically not positive. On the other hand, the heat equations in Sections 6 and 7 are positivity preserving equations in pointwise sense. After rescaling the variables, the weak limit of the Wigner distribution is a nonnegative phase space density, hence the equation must preserve this property. The time dependent quantum states (density matrices) cannot be reconstructed, but the heat equation determines their rescaled weak limits at any time.

2 Source of diffusion in various kinetic models

In order to explain the origin of diffusion in [5], we have to analyze the effects of the limits introduced there. To avoid Feynman path integrals, we present the basic idea of [5] in the mathematical language we will use in our proofs.

2.1 Eliminating the semiclassical parameter

We take the Hamiltonian as in [5] (see (1.1)) with $M = 1$ and specify the choice $V(x) = \frac{1}{2}x^2$ (harmonic oscillator), in the spirit of [12], [2], [29], [46], [11]. We use the fact that, for Gaussian Hamiltonians, the evolution equation for the Wigner transform of the density matrix is a first order linear partial differential equation ([47], [38], [26]), which can be solved by the method of characteristics (see also [46] for a similar observation).

In the quadratic case, we can scale h out of the equation (1.1). Let

$$H := \frac{1}{2}\Big(-\Delta_x + x^2\Big) + \frac{1}{2}\sum_{j=1}^{N\Omega}\Big(-\Delta_{R_j} + \omega_j^2 R_j^2\Big) + \frac{1}{\sqrt{N}}\Big(\sum_{j=1}^{N\Omega} C_j R_j\Big)\cdot x, \tag{2.1}$$

then $\exp\left(-ith^{-1}H_{CL}\right)$ and $\exp\left(-itH\right)$ are unitarily equivalent under the rescaling of variables $x \to xh^{-1/2}$, $R_j \to R_j h^{-1/2}$.

Hence *mathematically* (1.1) is the same as (2.1) and we will prefer to work with (2.1).

However, the *physical* interpretations are different. The Hamiltonian (1.1) is written in macroscopic coordinates, i.e. the particle position x is measured in meters and time t is measured in seconds. The Hamiltonian (2.1) is on microscopic (atomic) scales, where x is measured in Angströms, i.e. the particle is confined to a lengthscale of a few Å. The time is also measured in atomic time units. Hence the interpretation affects not only the lengthscale of the particle confinement, which depends on the size of the actual physical device, but more importantly the timescale of the evolution.

Naturally, one prefers the macroscopic interpretation for applications. However, we shall point out that in realistic models the linear coupling assumption is questionnable on macroscopic scales and even on microscopic scales it is a serious restriction (see Section 3.3). In order to assume the linearity only on microscopic scales, which we consider more realistic assumption for applications of these models, but still to be able to follow the evolution on larger (than atomic) time scales, we will introduce *scaling limits* of these models in Sections 6 and 7. In these sections the word "macroscopic" will refer to the scaling to be introduced there and it should not be confused with the scale separation provided by the semiclassical limit.

Hence our point of view is microscopic, we use the Hamiltonian (2.1), in our units $h = 1$, and we assume linear coupling on microscopic scales. In Section 5 we prove rigorous convergence to

the Fokker-Planck equation for the Wigner distribution on atomic scales. Due to the unitarity equivalence, this gives immediately the same Fokker-Planck equation on macroscopic scales if (1.1) can be used (with $V(x) = \frac{1}{2}x^2$), i.e. if the linear coupling is considered valid on macroscopic scales. Moreover, in this case the potential $\frac{1}{2}x^2$ can be replaced by an arbitrary potential $V(x)$, as (1.1) stands. Recall that any potential $V(x)$, apart from the quadratic ones, gives rise to a genuine pseudodifferential operator in the Wigner equation. In the semiclassical limit ($h \to 0$) this converges to the differential operator term $\nabla_x V \cdot \nabla_v w$ in (see also (1.6)). This fact is well-known for general semiclassical Wigner equations [38], [39], [27], [40]. We will not prove Theorem 5.1 for a general potential because our main goal is to find the origin of diffusivity which is independent of the confining potential. We restrict ourselves to the most convenient quadratic case.

More importantly, we present two different scaling limits starting from (2.1) which allows one to follow the dynamics up to macroscopic times. However, we believe that not just our result on the original Caldeira-Leggett model (in Section 5) can be extended to include general potential, but also the resonance effect in Sections 6 and 7. Due to the lack of explicit solutions, this requires extra analysis which we leave to further works.

2.2 Diffusion in the original model

After integrating out the reservoir variables in the equations for the characteristics, it eventually reduces to the following ODE for the particle's position variable $X(t)$ (see (4.14) for the exact result),

$$X''(t) + X(t) = f(t) + \lambda^2 \int_0^t S(t-s)X(s)\, ds \ . \tag{2.2}$$

Here λ is as in (1.3)-(1.4), S is an explicit function corresponding to the memory effects, and the forcing term f is,

$$f(t) = -\frac{\lambda}{\sqrt{N}} \sum_{j=1}^{N\Omega} \omega_j \Big[R_j \cos \omega_j t + P_j \frac{\sin \omega_j t}{\omega_j} \Big] \ , \tag{2.3}$$

where R_j, P_j are the initial position and momentum variables of the oscillators. Let $R_j^* := \sqrt{2\beta}\omega_j R_j$ and $P_j^* := \sqrt{2\beta} P_j$ be their rescaled versions. In the high temperature limit these become standard Gaussian variables since the classical Gibbs distribution is given by,

$$\prod_j e^{-\beta(P_j^2 + \omega_j^2 R_j^2)} = \prod_j e^{-\frac{1}{2}[(P_j^*)^2 + (R_j^*)^2]} \ ,$$

and at high temperature the quantum Gibbs distribution converges to the classical one (for the precise formulas, see (4.15)-(4.17)). Hence the choice (1.3) for C_j, together with (1.4) gives that,

$$f(t) = -\frac{\lambda_0}{\sqrt{2}} \sum_{j=1}^{N\Omega} \Big[\frac{R_j^*}{\sqrt{N}} \cos(\omega_j t) + \frac{P_j^*}{\sqrt{N}} \sin(\omega_j t) \Big] , \tag{2.4}$$

as $\beta \to 0$ with R_j^*, P_j^* being standard Gaussians.

After integration by parts in the memory term in (2.2) we obtain (see (4.36))

$$X''(t) + X(t) = f(t) + \lambda^2 \Omega X(t) - (M \star X')(t) - xM(t) \tag{2.5}$$

where M is an approximate Dirac delta function $M(t) \sim \lambda^2 \delta_0(t)$ in the limit $\Omega \to \infty$. Here $\star$ stands for convolution. The term $\lambda^2\Omega$ is the frequency shift of the test-particle oscillator. The friction term $M \star X'$ has a main Markovian part $\lambda^2 X'$ and a non-Markovian part which is negligible as $\Omega \to \infty$.

The effect of the limits introduced in [5] are as follows

• The high temperature limit ($\beta \to 0$) plays two roles. First, it ensures that the full friction term vanishes (recall $\lambda \sim \beta^{1/2}$). Second, it makes the rescaled initial data R_j^*, P_j^* standard Gaussians.

• In the thermodynamic limit ($N \to \infty$) the sum in (2.4) becomes the sum of the real and imaginary parts of the truncated complex white noise,

$$dW^{(\Omega)}(t) := \int_0^\Omega e^{i\omega t} g(d\omega) ,$$

where $g(d\omega)$'s are independent centered Gaussian random variables with variance $\mathbf{E}\Big[g(d\omega)^2\Big] = d\omega$ (for precise definition see Section 4.4).

• Removing the ultraviolet cutoff ($\Omega \to \infty$) gives the (complex) white noise,

$$dW(t) = \int_0^\infty e^{i\omega t} g(d\omega) \tag{2.6}$$

for the forcing term. Moreover, the simultaneous limit $\beta \to 0$, $\Omega \to \infty$ may lead to a constant phase shift $\lambda^2\Omega \sim \beta\Omega$.

In summary, the solution $X(t)$ to (2.2) converges to the solution of a pure harmonic oscillator with a white noise forcing, i.e. $\theta X(t) + \sigma X'(t) \sim (\eta \star dW)(t)$, where $\eta(s) = \theta \sin s + \sigma \cos s$ is the harmonic oscillator trajectory (with initial condition $\eta(0) = \sigma$, $\eta'(0) = \theta$). In particular the mean square displacement (both in space and velocity)

$$\mathbf{E}\Big|\theta X(t) + \sigma X'(t)\Big|^2 \sim \mathbf{E}\Big|\Big(\eta \star dW^{(\Omega)}\Big)(t)\Big|^2 = \int_0^\Omega \Big| \int_0^t \eta(t-s) e^{-i\omega s} ds \Big|^2 d\omega \tag{2.7}$$

behaves *quadratically* in t for small t for every finite Ω, hence it is *not* diffusive. The diffusive behavior (linear mean square displacement) is regained only *after* the $\Omega \to \infty$ limit or after long times.

We emphasize that, from this point of view, the irreversibility (the v-Laplacian) in the CL model immediately stems from the particular asymptotic distribution of the frequencies (uniform from zero to infinity) in the forcing term. In other terms this model demonstrates diffusion in a setup where a plain diffusive forcing mechanism was essentially put in by hand. Diffusion appears already in the microscopic (atomic) time scale as a result of high frequency oscillators. This means that there is a shorter, unexplored time scale on which most of the oscillators live, hence the initial Hamiltonian with the Caldeira-Leggett limits should not be considered microscopic, rather mesoscopic. This problem is especially transparent if the heat bath is provided by phonons (crystal lattice vibrations) which have an ultraviolet cutoff. We emphasize that introducing the Planck constant does not eliminate this problem, since large frequency *and* large wavelength (which is necessary for linear approximations) mean large propagation speed and the sound speed in metals ($10^3 - 10^4 \frac{m}{s}$) cannot be considered a very huge number in macroscopic units. In case of photons (electro-magnetic field), huge Ω is more realistic. We return to this point in see Section 3.3.

In contrast to this diffusive mechanism, the source of the diffusion in more realistic models dealing with a moving test-particle interacting with many degrees of freedom is the *scaling limit*. This means that in these models the full frequency spectrum of the diffusion is collected over a long time from the cumulative effects of interactions with bounded frequency, and the diffusive behaviour is visible only on a much larger time (and sometimes space) scale than that of the microscopic interaction (collision) mechanism. This makes a key difference between the present model and other works dealing, for instance, with collisional models as scaling limits of microscopic dynamics, i.e. macroscopic long time behaviour of Schrödinger equations (see e.g. [44], [45], [34], [28], [18], [19], [20], [21], [40], [41], [7], [8], [32] or also [4]).

We remedy this drawback of the CL scaling in Sections 6 and 7, as we indicate now.

2.3 Diffusion from resonances in the scaling limit

In Section 6, we show that one can also recover a diffusive non-kinetic behaviour from the Caldeira-Leggett Hamiltonian under a more realistic space-time scaling limit. Namely, for a *fixed* cutoff in frequency Ω, and after the high-temperature limit, we consider the resulting dynamics for the test-particle for large time $t \sim \alpha^{-2}$ and large space and velocity variables x, $v \sim \alpha^{-1}$. Here $\alpha \to 0$ is a scaling parameter. We prove that the phase space density is subject to a heat equation both in velocity and position variables. In particular, the energy of the

test-particle increases up to α^{-2} due to the resonances with bath particles of high energy (but bounded frequency). Recall that the temperature of the heat bath is $\beta^{-1} \to \infty$, hence bath particles can have large energy even with bounded frequency.

In this case the diffusion indeed comes from the cumulative effect of bounded frequency interactions via a change of scale (microscopic to macroscopic behaviour). This is in fact a high energy diffusion in phase space; the test-particle is heated up. The forcing frequency distribution can be quite arbitrary, the only condition is that it has to carry energy at the resonant frequency. The diffusion comes from a pure resonance effect, and this seems to be a more universal physical feature in this context (see [11]). However, the high temperature limit is still essential in this derivation.

In Section 7, we keep the temperature fixed and we rescale only time, $t = T\delta^{-1}$ (where $\delta \to 0$ plays to role of α^2 above), space and velocity remain unscaled. The reason is that the bath temperature is finite, hence the typical energy ("temperature") of the test-particle remains finite as well. Since the particle Hamiltonian is confining (energy level sets are compact in phase space), the particle remains effectively localized. As a result we get a small scale diffusion in phase space with friction, after integrating out the fast circular motion. Again the diffusion comes from resonance and is developed over a long time period, and the contributing bath frequencies are bounded.

2.4 Comparison of the three models

The main goal in all these Caldeira-Leggett type models is to derive diffusion. The time dependence of the mean square displacement of the characteristics (2.7) is quadratic for small time (unless $\Omega \to \infty$) and is linear for large time. To see diffusion on *all* times considered, there are two alternatives: either we take $\Omega \to \infty$ or we rescale time.

I.) If $\Omega \to \infty$, then λ must go to zero to keep the frequency shift $\lambda^2\Omega$ finite. Up to a positive time t, the total effect of the friction term is of order $\lambda^2 t$, while the diffusive (forcing) term is roughly of order $\lambda^2 t/\beta$ for larger times, see (5.14), however for short times it is only quadratic in t. Hence for finite times $\lambda^2 t \to 0$, the friction term vanishes. Moreover, the diffusive term vanishes as well, unless $\beta \to 0$ is chosen such that $\lambda^2 \sim \beta$. The frequency shift is of order $\beta\Omega$ and its actual size depends on the simultaneous limits $\beta \to 0$, $\Omega \to \infty$. If $\beta \to 0$ is taken first, then $\Omega \to \infty$, then there is no frequency shift. If $\beta\Omega$ is kept at a positive constant along the limits, then we see a frequency shift. These two cases are described in Theorem 5.1, where frictionless Fokker-Planck equations are derived on the microscopic time scale.

II.) If we consider long times, i.e. $t \sim \alpha^{-2}T$, $\alpha \to 0$ and T is fixed, then the size of the diffusive term is roughly $\lambda^2\alpha^{-2}T/\beta$ for *all* T. To compensate for the blowup α^{-2}, we can either rescale space and velocity ($x = \alpha^{-1}X$, $v = \alpha^{-1}V$) or we set $\lambda^2 \sim \alpha^2$.

II/a. If we rescale space and velocity as well, then the friction term has a size $\lambda^2 T$ and the diffusion term is of order $\lambda^2 T/\beta$ (in the new variables). One would like to keep λ and β fixed to see both friction and diffusion. But since the phase shift, $\lambda^2\Omega$, has to be kept finite, it forces keeping Ω finite as well. This is the most realistic physical situation. However, the friction has a non-Markovian part, whose size is $\lambda^2 T$ if Ω is fixed (and it goes to zero only if $\Omega \to \infty$). Hence the limiting equation must have a term which is nonlocal in time. This is the extra term which is missing in (1.6), but its inclusion would lead to an integro-differential equation and not to Fokker-Planck.

To derive a differential equation, the non-Markovian friction part has to be killed. With finite Ω it is possible only if $\lambda \to 0$, and then the full friction is eliminated. In order not to eliminate the diffusive term as well, $\beta \sim \lambda^2$ is necessary. This again leads to the high temperature limit, but now Ω is fixed and the diffusion comes from long-time cumulative resonance effects. The fast oscillator motion on the microscopic time scale has to be integrated out; either in time or by a radial averaging. This is the model in Section 6.

II/b. If we set $\lambda^2 \sim \alpha^2$ and keep β finite, then we see a finite diffusion on a microscopic space and velocity scale. The friction term $\lambda^2 t$ remains positive and the ratio of the friction to the diffusion is β, which gives Einstein relation. Hence Ω could be kept fixed to see the diffusion mechanism.

However, the non-Markovian part of the memory does not vanish unless $\Omega \to \infty$. The qualitative analysis of Section 7 shows that Ω can grow very slowly (like $|\log\alpha|^7$), i.e. the non-Markovian part of the friction is weak for large times and moderately large Ω. This was probably the heuristic idea of Caldeira and Leggett to neglect this term. However, this effect shows up only after time rescaling; for finite microscopic times t this term is not negligible.

Hence we let $\Omega \to \infty$, and assume that $\lambda^2\Omega$ converges to a fixed number (possibly zero). This number gives the frequency shift. Again, we see that the size of the frequency shift delicately depends on the simultaneous limiting procedure. This is the model of Section 7 (where $\delta := \alpha^2$ is introduced for brevity).

We point out that in models II/a and II/b the origin of the diffusion is the time rescaling. Since the forcing frequencies are kept finite, there is no diffusion on the microscopic scale; it becomes visible only after the large time rescaling. Hence the physically questionnable limits, $\beta \to 0$, $\Omega \to \infty$ (see Section 3) have nothing to do with the emergence of the diffusion in these models.

However, at least one of these limits is necessary to arrive at a differential equation instead of an integro-differential equation with time delayed memory term. In model II/a. (Section 6) we used $\beta \to 0$ and kept Ω fixed, while in II/b. (Section 7) we let $\Omega \to \infty$ and kept β finite.

We always consider nonnegative times $t \geq 0$. However, most of our computations are valid for *any* time, except those which are directly responsible for the emergence of the diffusion (Laplacian, or linear mean square displacement). We shall point out these steps. If time were evolved backward, $t < 0$, then the same argument would yield an opposite sign of the Laplacian (so that along the evolution it is regularizing) in the final limiting equations. This is the usual phenomenon of irreversibility of the parabolic equations.

2.5 Derivation of Fokker-Planck equation via Boltzmann equation

In the Caldeira-Leggett type models we assumed that the test-particle is localized and is subject to a harmonic heat bath with linear interaction. This usually describes particles trapped in a microscopic cavity.

For transport phenomena it is more natural to consider a free test-particle subject to a collision mechanism. In these models the collisions are provided by impurities (Lorenz gas) or by a system of many noninteracting particles (Rayleigh gas or phonon models) and one focuses only on the dynamics of the test-particle. The goal is to derive an equation for the reduced phase space distribution from the Hamiltonian dynamics with many degrees of freedom. A scaling limit is necessary to eliminate the details of the single collisions and to keep only their cumulative long-time effects. The effect of a single collision is weakened. One can introduce a weak coupling parameter $\lambda \to 0$; one can consider a gas at low density $\varrho \to 0$ or, in the Rayleigh gas case, one can let the mass ratio of the gas particle and test-particle m/M go to zero. In all cases the time is rescaled as $t = T\delta^{-1}$. The first scale on which collision effects are visible is $\delta \sim \lambda^2$ (weak coupling or van-Hove limit) or $\delta \sim \varrho$ (low density or Grad limit) and $\delta \sim m/M$ (heavy test-particle limit).

In classical mechanics, the limiting equation is the linear Landau equation (or: diffusion on the energy surface) for the van-Hove limit [33]; the linear Boltzmann equation for the low density case ([25], [43], [3]); and the Fokker-Planck equation for the heavy test-particle case ([17]). The Fokker-Planck equation can be obtained in a two step limit as well: first one obtains a linear Boltzmann equation via a low density limit, then a Fokker-Planck equation from a mass rescaling (for an excellent review see [44]).

In quantum mechanics the limiting equation is the linear Boltzmann equation both in the case of the Lorenz gas (see [18] for the low density case and [19] for the weak coupling case) and in the case of the weakly coupled phonons [20]. In the model of [20] a more realistic nonlinear

phonon coupling is considered.

In all cases when the first nontrivial limiting equation is Boltzmann, one needs an extra limiting procedure to derive a diffusive equation. For example if the momentum change in the collisions is small (e.g. the mass ratio m/M is small), then a Taylor expansion in the Boltzmann collision operator gives the Fokker-Planck equation in the first nontrivial order (see [36], for rigorous proof [31]). The smallness of the collisions has to be compensated by an extra time rescaling. However, the two step time rescaling cannot be considered as a fully satisfactory derivation since in the first (Boltzmann) limit correlations are neglected which could become relevant on a larger time scale. The proper (but much harder) procedure is to follow the Hamiltonian dynamics up to the desired (larger) time scale.

We remark that a considerably more difficult collision mechanism is when all particles interact, they are identical, and we are interested in the evolution of the one particle marginal distribution (or density matrix). In this case, the limiting equation is expected to be a nonlinear Boltzmann equation and in classical mechanics it was proven by Lanford [35]. In quantum mechanics the correlation structure is complicated and even the first nontrivial (Boltzmann) time scale is not understood rigorously.

Finally, we compare our model II/b to these free kinetic models with collisions. The closest related model is a free electron subject to a weakly coupled phonon interaction considered in [20], where a (linear) Boltzmann equation was derived. In both models the time scale is the van Hove scale $t \sim \lambda^{-2}$, where λ is the coupling constant. In case of the realistic (nonlinear) electron-phonon coupling in [20], each phonon mode contributes equally to the collision mechanism. However, in the model II/b the source of the diffusion is resonance which originates merely in the test-particle confinement, however for the rigorous proof we need to use the special form of the linear coupling and test-particle Hamiltonian. Phonons with frequencies away from the base frequency of the test-particle Hamiltonian do not contribute, while phonons near the resonance frequency have a strong long time effect. In particular, it is easy to see that the Duhamel expansion used in [20] diverges for the model II/b, which is also an indication that there is no Boltzmann equation behind the Fokker-Planck equation derived in Section 7.

3 The assumptions of the Caldeira-Leggett model

We discuss the three key assumptions of [5] mentioned in the Introduction. As we have explained in Section 2.2, all these assumptions play an important *mathematical* role in the Caldeira-Leggett argument, but their physical content has not been explained. We do not wish to enter long physical speculations in this section, just we would like to point out that the type of

assumptions needed for the mathematics of the problem cannot be taken for granted physically.

3.1 Heat bath and its frequencies

Caldeira and Leggett make two assumptions on the frequencies: the frequency distribution is uniform on $[0, \infty]$ and the coupling coefficients are proportional to ω (see (1.3)). In order to see their validity, the first physical question is the source of the harmonic heat bath. We note that it is not so easy to realize a harmonic heat bath physically; the standard gases or fluids, which often serve as an environment, are not harmonic oscillators, but phonons and photons are. However phonons and photons have a quite specific frequency distribution. We refer, e.g. to [42], or also [10] for an account on phonon- or photon- models.

The infinite ultraviolet cutoff $\Omega \to \infty$ is particularly questionnable in the phonon model, where the lattice spacing is a natural shortest lengthscale and typically the Schrödinger equation for electrons lives on the same lengthscale. In photon models removing the ultraviolet cutoff is more acceptable since it is not related to other physical quantities on atomic scales. However in Section 3.3 we will see that the linear coupling assumption sets an upper limit on the frequency cutoff. We mention that in a photon picture one has to consider a velocity coupling instead of a position coupling in (2.1) ($R_j \frac{\partial}{\partial x}$ instead of $R_j x$), however the present method goes through for velocity coupling as well.

Setting frequency dependent coupling coefficient (1.3) is equivalent to having a *frequency independent* coupling and a quadratic distribution of oscillator frequencies. Indeed, we can construct a Hamiltonian, similar to (2.1) which gives the same dynamics for the test-particle, and the coupling is frequency independent. To do that, we consider ω_j^2 copies of independent harmonic oscillators with the same frequency ω_j and variable $R_j^{(i)}$, $i = 1, 2, \ldots, \omega_j^2$ (assuming for the moment that ω_j^2 is an integer). Then for all j the j^{th} mode of the reservoir Hamiltonian $-\frac{1}{2}\Delta_{R_j} + \frac{1}{2}\omega_j^2 |R_j|^2$ is replaced with,

$$\sum_{i=1}^{\omega_j^2} \Big(-\frac{1}{2}\Delta_{R_j^{(i)}} + \frac{1}{2}\omega_j^2 |R_j^{(i)}|^2 \Big) \ ,$$

and $C_j R_j = \lambda \omega_j R_j$ is replaced with $\lambda \sum_{i=1}^{\omega_j^2} R_j^{(i)}$ in the interaction term. This modification does not change the marginal evolution of the test-particle as it is seen by changing the variable,

$$R_j \to \frac{1}{\omega_j} \sum_{i=1}^{\omega_j^2} R_j^{(i)} \ ,$$

and integrating out the irrelevant relative differences of the $R_j^{(i)}$'s. This explains how to change the problem with a coupling constant that depends linearly on the frequency into an equivalent

problem of a test-particle coupled to a reservoir in a frequency independent way, at the expense of considering quadratically distributed oscillators, i.e. the number of oscillators with frequency ω_j is proportional to ω_j^2 . If the ω_j^2's are not integers, then this equivalence becomes exact only after the thermodynamic limit, yet to be described. We will not prove this fact rigorously here. The proof is not complicated, but it is unnecessary for the present discussion; we start our rigorous work from (2.1); other possible representations are used only for physical explanations.

Using this observation, the assumption (1.3) suggests that the abstract reservoir of oscillators in (2.1) comes from phonons or photons in the three-dimensional space. In general, in the d-dimensional physical space, the number of photons/phonons having a given frequency ω is proportional to ω^{d-1} if the dispersion relation is massless. Hence the factor ω_j in (1.3) should not enter the coupling parameter, but rather should be considered as the square root (by the standard normalization) of the number of photons/phonons with given frequency ω_j in $\mathbb{R}^3$.

3.2 High temperature

Choosing temperature dependent coupling and letting the temperature go to infinity is quite convenient in the Caldeira-Leggett argument (Section 2.2), however both assumptions are slightly suprising and should not be needed to detect diffusion. Naturally, the Einstein relation (if valid) in a frictionless diffusive equation forces $\beta \to 0$. But as we mentioned in Section 2.5, Fokker-Planck equation can be obtained in the $m/M \to 0$ limit of the classical Rayleigh gas for *any* value of the temperature of the external bath and one obtains the Einstein relation as well. One would like to prove a similar result starting from quantum mechanics with (massless) phonons at finite temperature; this was the main motivation for introducing a new scaling limit in Section 7 (model II/b in Section 2.4).

In conjunction of the $\beta \to 0$ limit we also mention that if we think of the oscillators as phonons, then large temperature ($> 10^3 K$) destroys the lattice structure of the metal by melting. Large temperatures could be more realistic in a photon picture.

3.3 Linear coupling

The linear coupling between the heat bath and the test-particle is the key assumption in [5], as well as in many other works in this field, and also in this article. By linear coupling we mean an interaction that is linear in *both* the reservoir and the particle variables. Linearity in the reservoir variables is widely used and accepted since in typical physical situations these variables are small (like ion displacement in the lattice), hence nonlinear interactions can be approximated by linearization. Linearity in the particle variable is a serious assumption, since it implicitly assumes microscopically localized particles. It is especially questionnable in models

for electron transport. However, we readily admit that our work heavily relies on this assumption, because this makes the model almost explicitly computable. Without this assumption only the Boltzmann equation has been proved in the case of a free test-particle coupled to a phonon field with different, more involved methods [20].

We briefly discuss the linearity assumption below in the phonon and photon case.

The case of phonons.

Let Λ be a crystal lattice in $\mathbb{R}^d$, and Λ^* be its dual lattice. Written in normal variables (see e.g. [42]) and after linearization in the phonon variables the interaction of an electron with the crystal lattice is,

$$H_I = \sum_{k \in \Lambda^*} C_\omega \cdot R_\omega \exp(ik \cdot x) \ , \tag{3.1}$$

where $\omega = \omega(k) = |k|$ from the usual dispersion relation (more precisely, we have $H_I = \sum_{k \in \Lambda^*} D(k) \cdot \tilde{R}_k \exp(ik \cdot x)$ where the $\tilde{R}_k$ is the normal mode of the lattice vibration with wave vector k, and $D(k)$ is the k-th Fourier component of the electron-photon interaction, but we can assume (3.1) at least for radial coupling). The essential point in (3.1) is that this interaction is a priori highly non-linear in x. One can reach linear coupling by assuming that the quantity $k \cdot x$ in (3.1) remains small during the full evolution of the system, and linearize the exponential accordingly. This means that the wavelength ($= O(|\text{wavevector}|^{-1}) = O(|k|^{-1})$) of the crystal oscillation should be bigger than the displacement of the particle (x) during its full evolution. Furthermore, in the original Caldeira-Leggett model (as well as in Section 5.3) the ultraviolet cutoff was removed ($\Omega \to \infty$) in order to obtain diffusion (see Section 2.2). Therefore, we are led to assume huge frequencies together with big wavelengths, whereas the typical sound speed in metals (= frequency $\times$ wavelength) is bounded. One can take the idealized "infinite sound speed", or, equivalently, "infinitely stiff lattice" limit, but given the actual size of the physical parameters this is a physically questionable procedure.

On the level of the Hamiltonian, notice that $\sum \omega_j R_j \sim \sum_j \sum_i R_j^{(i)} = \sum_{k \in \Lambda^*} R_k$, to which the particle coordinate is coupled (2.1), is just the displacement of the ion at the origin as the normal modes are the Fourier transforms of the displacement vectors. In other words, the test-particle is assumed to remain in the vicinity of the origin (on atomic scales), hence it interacts with one single ion of the crystal lattice for all its dynamics (see e.g. [13]). On the other hand, we wish to derive a diffusive equation for the electron, i.e. for large values of time it is expected to move away from the origin. Even if the diffusion appears only in the velocity (see (1.5)), the large velocity implies large fluctuation in the configuration variable as well.

In summary, the linear model effectively involves an implicit mean-field assumption by requiring that the test-particle is coupled to the same mode for all its evolution, which seems

incompatible with the finite sound speed of the metals along with the removed UV cutoff. This leaves a serious doubt on the applicability of the linear coupling assumption for diffusion models for electron propagation in an ionic lattice (see also [2] for a brief criticism of this assumption).

The case of photons.

As mentioned before, one can interpret (2.1) as the interaction of an electron with an external electro-magnetic field (up to coupling with velocities - see above). In this case, the electron interacts with the field through (see, e.g., [11]),

$$A(x) = \int_k \sum_e (\hbar/\omega)^{1/2} \Big[e a_e(k) \exp(ikx) + e a_e^+(k) \exp(-ikx) \Big] , \tag{3.2}$$

where k is the momentum of the photon, e its polarization, $\omega = c|k|$, c is the speed of light, and $a_e(k)$, $a_e^+(k)$ are the annihilation and creation operators. To be more precise, the interaction potential in this case is $V(x) = -(q/M)A(x) \cdot p$, where q is the electric charge of the electron, M its mass, and $p = -i\frac{\partial}{\partial x}$ its momentum.

In this case, one can argue that the speed of light is large, so that, contrary to the phonon picture, one can reach simultaneously high frequencies and large wavelengths. But again, the linearization needed for our purposes means that we couple the electron to the homogeneous field $A(0)$ instead of $A(x)$ during the full evolution of the electron, and this assumption is questionnable for large values of time in the kinetic regime, when the electron evolves far away from the origin. It is more realistic in models to study energy dissipation of a localized electron coupled to a radiation field with bounded frequency. This is the main reason why we wanted to eliminate the $\Omega \to \infty$ limit from the derivation (Section 7). Note however that the replacement of $A(x)$ by $A(0)$ is fairly standard (at least for small displacements of the electron): it is found in all the above mentioned works, it is also the basic approximation in the so-called Pauli-Fierz-Kramers transform (see [11]).

4 Preliminary results

4.1 The Wigner formalism

The density matrix,

$$\rho^{N,\varepsilon} := \rho^{N,\varepsilon}(t, x, y, R, Q) , \tag{4.1}$$

which is the solution of,

$$i\partial_t \rho^{N,\varepsilon} = [H, \rho^{N,\varepsilon}] , \tag{4.2}$$

represents the state of the system "particle + reservoir" at time t with the reservoir variables $R = (R_1, \dots, R_{N\Omega})$, $Q = (Q_1, \dots, Q_{N\Omega})$. We index the density matrix by N and the superscript $\varepsilon = (\beta, \Omega, \lambda)$ stands for all the other scaling parameters; recall that β is the inverse temperature, Ω is the frequency range and λ is the coupling strength in the Hamiltonian (2.1).

We take the initial data (independent of ε for simplicity),

$$\rho_A^0 \otimes e^{-\beta H_R} \ , \tag{4.3}$$

with $\rho_A^0 := \rho_A^{N,\varepsilon}(t=0)$. Here $H_R := \frac{1}{2}\sum_{k=1}^{N\Omega}\Big(-\Delta_{R_k} + \omega_k^2 R_k^2\Big)$ is the reservoir Hamiltonian and $\rho_A^{N,\varepsilon}(t,x,y)$ is the density matrix at time t of the test-particle. It is defined by

$$\rho_A^{N,\varepsilon}(t,x,y) := \int_{\mathbb{R}^{N\Omega}} \rho^{N,\varepsilon}(t,x,y,R,R)\, dR \ , \tag{4.4}$$

with the obvious notation $dR = dR_1 \dots dR_{N\Omega}$. As usual, we do not distinguish between operators and their kernels in the notation. Following [5], we have to compute,

$$Tr_R\Big(e^{-itH}(\rho_A^0 \otimes e^{-\beta H_R})e^{itH}\Big) \ , \tag{4.5}$$

where Tr_R is the partial trace over the reservoir variables. We observe that the Hamiltonian (2.1) is quadratic, so that equation (4.2) can actually be transformed into a first order transport partial differential equation by using the Wigner transform. Indeed, let us define the Wigner transform $w^{N,\varepsilon}(t)$ of $\rho^{N,\varepsilon}(t)$ by,

$$w^{N,\varepsilon}(t,x,v,R,P) := \tag{4.6}$$

$$:= \int_{\mathbb{R}^{N\Omega+1}} \rho^{N,\varepsilon}\Big(t, x+\frac{y}{2}, x-\frac{y}{2}, R+\frac{Q}{2}, R-\frac{Q}{2}\Big) \times \ \exp\Big(-i[yv + \sum_{k=1}^{N\Omega} Q_k P_k]\Big)\, dy\, dQ \ .$$

Also, let us define the Wigner transform of $\rho_A^{N,\varepsilon}$ by,

$$w_A^{N,\varepsilon}(t,x,v) := \int_{\mathbb{R}} \rho_A^{N,\varepsilon}\Big(t, x+\frac{y}{2}, x-\frac{y}{2}\Big) \ \exp(-iyv)\, dy \ . \tag{4.7}$$

We have the well-known property,

$$w_A^{N,\varepsilon}(t,x,v) := \int_{\mathbb{R}^{2N\Omega}} w^{N,\varepsilon}(t,x,v,R,P)\, dR\, dP \ , \tag{4.8}$$

and the initial datum for $w^{N,\varepsilon}$ is easily computed from (4.3) and the Mehler kernel,

$$w^{N,\varepsilon}(t=0,x,v,R,P) = w_0(x,v) W_0^{N,\varepsilon}(R,P) \tag{4.9}$$

with

$$W_0^{N,\varepsilon}(R,P) : = \prod_{k=1}^{N\Omega}\Big[\, 4\pi\Big(\frac{\cosh(\beta\omega_k)-1}{\cosh(\beta\omega_k)+1}\Big)^{1/2} \times \exp\Big(-\{\frac{\omega_k(\cosh(\beta\omega_k)-1)}{\sinh(\beta\omega_k)}R_k^2\}\Big)\ \exp\Big(-\{\frac{\sinh(\beta\omega_k)}{\omega_k(\cosh(\beta\omega_k)+1)}P_k^2\}\Big)\Big]\ .$$

Here, $w_0(x,v)$ is the initial datum for the test-particle, i.e. it is the Wigner transform of $\rho_A^0(x,y)$. Here and in the sequel, we shall assume the following regularity for w_0,

$$\widehat{w}_0(\xi,\eta) := \int_{\mathbb{R}^2} w_0(x,v)\exp(-i[x\xi+v\eta])\,dx\,dv \quad \in L^1(\mathbb{R}_\xi\times\mathbb{R}_\eta)\ . \tag{4.10}$$

It is well known that, if $\rho^{N,\varepsilon}$ satisfies the Von-Neumann equation (4.2) with Hamiltonian given by (2.1), then its Wigner transform (4.6) satisfies the following partial differential equation,

$$\partial_t w^{N,\varepsilon} + v\,\partial_x w^{N,\varepsilon} - x\,\partial_v w^{N,\varepsilon} + \sum_{k=1}^{N\Omega}\Big(P_k\,\partial_{R_k} w^{N,\varepsilon} - \omega_k^2 R_k\,\partial_{P_k} w^{N,\varepsilon}\Big) \tag{4.11}$$
$$-\frac{\lambda}{\sqrt{N}}\Big(\sum_{k=1}^{N\Omega}\omega_k R_k\Big)\,\partial_v w^{N,\varepsilon} - \frac{\lambda}{\sqrt{N}}\Big(\sum_{k=1}^{N\Omega}\omega_k x\,\partial_{P_k} w^{N,\varepsilon}\Big) = 0\ .$$

As a conclusion we can now rephrase our original problem in the Wigner formalism: following [5], we want to derive a diffusive behaviour for $w_A^{N,\varepsilon}(t)$, the trace of $w^{N,\varepsilon}(t)$, in the thermodynamic limit ($N\to\infty$) and in certain limiting regimes of ε. Here, $w^{N,\varepsilon}$ satisfies (4.11) with initial datum (4.9).

4.2 Solution by characteristics

Equation (4.11) can easily be solved by the method of characteristics. In fact, for all values of time t, and for all smooth, compactly supported test functions $\phi(x,v)$, we have,

$$\int_{\mathbb{R}^2} w_A^{N,\varepsilon}(t,x,v)\overline{\phi}(x,v)\,dx\,dv = \int_{\mathbb{R}^{2N\Omega+2}} w(t=0,x,v,R,P)\,\overline{\phi}(X(t),V(t))\,dx\,dv\,dR\,dP \tag{4.12}$$
$$= \int_{\mathbb{R}^{2N\Omega+6}} \hat{w}_0(\xi,\eta)\overline{\hat{\phi}(\theta,\sigma)}e^{i(x\xi+v\eta)}e^{-i(X(t)\theta+V(t)\sigma)}W_0^{N,\varepsilon}(R,P)\,dx\,dv\,dR\,dP\,d\xi\,d\eta\,d\theta\,d\sigma,$$

where we have introduced the (forward) characteristics,

$$X'(t) = V(t)\ , \quad V'(t) = -X(t) - \frac{\lambda}{\sqrt{N}}\sum_{k=1}^{N\Omega}\omega_k R_k(t) \tag{4.13}$$
$$R_k'(t) = P_k(t)\ , \quad P_k'(t) = -\omega_k^2 R_k(t) - \frac{\lambda}{\sqrt{N}}\omega_k X(t)\ ,$$

with initial data $X(0) = x$, $V(0) = v$, $R_k(0) = R_k$ and $P_k(0) = P_k$. Here we used that the flow (4.13) preserves the Lebesgue measure over $\mathbb{R}^{2(N\Omega+1)}$. For simplicity, we did not index the characteristics by N, ε, but $X(t), V(t)$ in (4.12) depend on N, ε. However, sometimes we will use $X_N(t)$ for special emphasis.

Integrating with respect to $R_k(t)$ in (4.13) and inserting the result in the equation for $X(t)$ gives,

$$\begin{aligned} X''(t) + X(t) &= -\frac{\lambda}{\sqrt{N}} \sum_{k=1}^{N\Omega} \omega_k \Big[R_k \cos \omega_k t + P_k \frac{\sin \omega_k t}{\omega_k} \Big] \\ &\quad + \frac{\lambda^2}{N} \sum_{k=1}^{N\Omega} \int_0^t \omega_k \sin \omega_k (t-s) X(s) ds \ . \end{aligned} \tag{4.14}$$

The right-hand-side of (4.14) is of the form 'forcing term + memory term'.

In view of (4.9) and (4.12), the partial trace over the oscillators is an integral with respect to a Gaussian distribution in R_k, P_k with (unnormalized) density,

$$\exp \Big[- \frac{\omega_k (\cosh \beta\omega_k - 1)}{\sinh \beta\omega_k} R_k^2 - \frac{\sinh \beta\omega_k}{\omega_k (\cosh \beta\omega_k + 1)} P_k^2 \Big] \ . \tag{4.15}$$

Changing variables such that,

$$r_k = \sqrt{\frac{2\omega_k (\cosh \beta\omega_k - 1)}{\sinh \beta\omega_k}} R_k \ , \quad p_k = \sqrt{\frac{2 \sinh \beta\omega_k}{\omega_k (\cosh \beta\omega_k + 1)}} P_k \ , \tag{4.16}$$

we obtain (after normalization) the standard Gauss measure,

$$d\mu_N = \prod_{k=1}^{N\Omega} \frac{1}{2\pi} e^{-\frac{1}{2}(r_k^2 + p_k^2)} dr_k dp_k \ , \tag{4.17}$$

i.e. r_k, p_k are independent standard Gaussian variables. The integration with respect to this probability measure will be denoted by $\mathbf{E}_N$.

Using these new variables and integration by parts with respect to s, the equation (4.14) for $X_N(t) = X(t)$ becomes,

$$X_N''(t) + X_N(t) = f_N(t) + \lambda^2 \Omega X_N(t) - (M_N \star X_N')(t) - x M_N(t) \ , \tag{4.18}$$

with,

$$f_N(t) := -\frac{\lambda}{\sqrt{N}} \sum_{k=1}^{N\Omega} A_\beta(\omega) \Big[r_k \cos \omega_k t + p_k \sin \omega_k t \Big] \ , \tag{4.19}$$

and,

$$M_N(t) := \frac{\lambda^2}{N} \sum_{k=1}^{N\Omega} \cos \omega_k t \ . \tag{4.20}$$

Here we defined,

$$A(\omega) = A_\beta(\omega) := \sqrt{\frac{\omega(\cosh \beta\omega + 1)}{2 \sinh \beta\omega}} \ . \tag{4.21}$$

We see that the memory term is split into three parts. The term $\lambda^2 \Omega X_N$ induces a frequency shift of the test-particle oscillator, $M_N \star X_N'$ is the friction term and the last inhomogeneous term will be irrelevant. We define

$$a^2 = a_\varepsilon^2 := 1 - \lambda^2 \Omega$$

(recall that ε stands for the triple (β, Ω, λ)), and we always assume that a_ε is uniformly separated from zero, e.g. $\frac{1}{2} \leq a_\varepsilon \leq 1$. We can rewrite (4.18) as

$$X_N''(t) + a^2 X_N(t) = f_N(t) - (M_N \star X_N')(t) - x M_N(t) \ . \tag{4.22}$$

4.3 The thermodynamic limit

We now perform the limit $N \to \infty$. A possible way is to solve (4.14) (iteratively), and compute the limit in the corresponding formulae (see (4.42) later). This rigorously gives the thermodynamic limit but we present an alternative approach which is more illuminating to explain the asymptotic diffusion that we shall recover in Section 5.3. We first need an a priori bound.

Lemma 4.1 *Let $X_N(t)$ solve (4.22) with initial conditions $X(0) = x$, $X'(0) = v$, and let*

$$F_N(t) := \sup_{s \leq t} \mathbf{E}_N |X_N(t)| + \sup_{s \leq t} \mathbf{E}_N |X_N'(t)| \ . \tag{4.23}$$

Then there is a constant $C > 0$ such that

$$F_N(t) \leq C e^{Kt} \left(|x| + |v| + K|x| + \sup_{s \leq t} \left\{ s e^{-Ks} \right\} \left[\lambda^2 \Omega \left(\beta^{-1} + \Omega \right) \right]^{1/2} \right) \ . \tag{4.24}$$

uniformly in N, where

$$K = K(\lambda, \Omega) := C\lambda^2 \left(1 + \frac{1}{|\Omega - a|} \right) \ . \tag{4.25}$$

and $a^2 = 1 - \lambda^2 \Omega \in [\frac{1}{4}, 1]$.

Proof. From the fundamental solution of (4.22), one has

$$\begin{aligned} X_N(t) &= x\cos at + va^{-1}\sin at \\ &\quad + \int_0^t a^{-1}\sin a(t-s)\Big[f_N(s) - (M_N \star X_N')(s) - xM_N(s)\Big]ds\ , \\ X_N'(t) &= -xa\sin at + v\cos at \\ &\quad + \int_0^t \cos a(t-s)\Big[f_N(s) - (M_N \star X_N')(s) - xM_N(s)\Big]ds\ . \end{aligned} \tag{4.26}$$

First step. To estimate the memory term in (4.26), we write,

$$\begin{aligned} \int_0^t \sin[a(t-s)](M_N \star X_N')(s)ds &= \Big(\sin(a\,\cdot\,) \star M_N \star X_N'\Big)(t) \\ &= \int_0^t \Big(\int_0^s \sin[a(s-u)]M_N(u)du\Big)X_N'(t-s)ds\ , \end{aligned} \tag{4.27}$$

An easy calculation shows that the inner integral is bounded by

$$\Big|\int_0^s \sin[a(s-u)]M_N(u)du\Big| = \Big|\Big(M_N \star \sin(a\,\cdot\,)\Big)(s)\Big| \le k\lambda^2\Big(1 + \frac{1}{|a-\Omega|}\Big)\ , \tag{4.28}$$

with a universal constant k uniformly in N. Indeed, notice that,

$$\lim_{N\to\infty} M_N(s) = \lambda^2\frac{\sin\Omega s}{s} =: M(s)\ , \tag{4.29}$$

uniformly for $s \in [0,t]$. Moreover $\int_0^s \sin[a(s-u)]M(u)du$ can be estimated by splitting the integration into two regimes $u \le 1$ and $u \ge 1$ (or $u \le s$ regime only if $s \le 1$) and both regimes can be estimated by elementary integration by parts to obtain (4.28).

Hence the expected value of the integral of the memory terms in (4.26) is estimated by,

$$\begin{aligned} &\mathbf{E}_N\Big|\int_0^t a^{-1}\sin a(t-s)\Big[-(M_N \star X_N')(s) - xM_N(s)\Big]ds\Big| \\ &\qquad \le a^{-1}k\lambda^2\Big(1 + \frac{1}{|a-\Omega|}\Big)\Big[|x| + \int_0^t F_N(s)\,ds\Big]\ , \end{aligned} \tag{4.30}$$

and similarly for the cosine term in (4.26).

Second step. For the forcing term one computes,

$$\mathbf{E}_N\Big|\int_0^t \sin[a(t-s)]f_N(s)ds\Big| \le t\sup_{s\le t}\Big(\mathbf{E}_N|f_N(s)|^2\Big)^{1/2}\ . \tag{4.31}$$

We have,

$$\mathbf{E}_N|f_N(s)|^2 = \frac{\lambda^2}{N}\sum_{k=1}^{N\Omega} A_\beta^2(\omega) \le \hat{k}\lambda^2\Omega\Big(\beta^{-1} + \Omega\Big)\ , \tag{4.32}$$

where $\hat{k}$ is again some positive constant, independent of N. Indeed, this sum is an approximating Riemann sum for the integral,

$$\lambda^2 \int_0^\Omega A_\beta^2(\omega) d\omega = \lambda^2 \int_0^\Omega \frac{\omega(\cosh \beta\omega + 1)}{2 \sinh \beta\omega} \, d\omega \ ,$$

which satisfies the estimate (4.32). Hence we obtain,

$$\begin{aligned} \mathbf{E}_N \Big[|X_N(t)| + |X_N'(t)| \Big] \ &\leq \ |x| + |v| + k\lambda^2 \Big(1 + \frac{1}{|a - \Omega|} \Big) \Big[|x| + \int_0^t F_N(s) \, ds \Big] \\ &\quad + t \Big[\hat{k} \lambda^2 \Omega \Big(\beta^{-1} + \Omega \Big) \Big]^{1/2} . \end{aligned} \tag{4.33}$$

By a standard Gronwall-type argument we conclude (4.24). ■

4.4 Digression on stochastic integrals

Stochastic integration is integration with respect to a random measure. Once the measure is specified, the integrals are defined as limits of integrals of stepfunctions. We do not develop this notion here, just indicate how it is related to the present problem.

Definition 4.1 *The ensemble of random variables $g(A)$, A running over the Borel sets of $\mathbb{R}$, is called standard Gaussian random measure if $g(A)$ is a centered real Gaussian random variable for all A and $\mathbf{E} g(A) g(B) = |A \cap B|$ where $|\cdot|$ is the Lebesgue measure.*

In the thermodynamic limit $N \to \infty$, the forcing term (4.19) converges in an $L^2(d\mu_N)$ sense towards the stochastic integral,

$$f(t) := -\lambda \int_0^\Omega A_\beta(\omega) \Big[r(d\omega) \cos \omega t + p(d\omega) \sin \omega t \Big] \ , \tag{4.34}$$

where $r(d\omega), p(d\omega)$ are independent standard Gaussian random measures. The expectation with respect to their joint measure is denoted by $\mathbf{E}$. Clearly $f_N(t)$ is a Riemann sum approximation of $f(t)$ by choosing $r_k := N^{1/2} r\Big(\Big[\frac{k-1}{N}, \frac{k}{N} \Big] \Big)$ and $p_k := N^{1/2} p\Big(\Big[\frac{k-1}{N}, \frac{k}{N} \Big] \Big)$, since their distribution is $d\mu_N$ (see (4.17)). In particular we can realize all f_N's and f on a common probability space. Note that $f(t)$ is formally a white noise (see (2.6)) when the 'hyperbolic factor' $A_\beta(\omega)$ is replaced by one and $\Omega = \infty$.

Lemma 4.2 *For $1 < \Omega < \infty$ there exist a random function $X(t)$ such that,*

$$\lim_{N \to \infty} \Big(\sup_{s \leq t} \mathbf{E} |X_N(s) - X(s)| + \sup_{s \leq t} \mathbf{E} |X_N'(s) - X'(s)| \Big) = 0 \ , \tag{4.35}$$

and $X(t)$ almost surely satisfies the equation,

$$X''(t) + a^2 X(t) = f(t) - (M \star X')(t) - xM(t) \ , \tag{4.36}$$

with initial conditions $X(0) = x$, $X'(0) = v$. Moreover,

$$F(t) := \sup_{s \le t} \mathbf{E}|X(s)| + \sup_{s \le t} \mathbf{E}|X'(s)| \ ,$$

satisfies the same estimate as $F_N(t)$ (see (4.24)),

$$F(t) \le Ce^{Kt}\Big(|x| + |v| + K|x| + \sup_{s \le t} \Big\{ se^{-Ks} \Big\} \Big[\lambda^2 \Omega \Big(\beta^{-1} + \Omega \Big) \Big]^{1/2} \Big) \ . \tag{4.37}$$

Proof. Let us define $X(t)$ by the integral equation,

$$\begin{aligned} X(t) &= x \cos at + va^{-1} \sin at \\ &\quad + \int_0^t a^{-1} \sin[a(t-s)] \Big[f(s) - (M \star X')(s) - xM(s) \Big] ds \ , \end{aligned} \tag{4.38}$$

Since,

$$\int_0^t \mathbf{E}|f(s)|^2 ds = \lambda^2 \int_0^\Omega \frac{\omega(\cosh \beta\omega + 1)}{2 \sinh \beta\omega} d\omega < \infty \ ,$$

$X(t)$ is well defined almost surely and satisfies (4.36). Moreover, the uniformity of (4.24) in N, and (4.35) shows that $F(t)$ satisfies (4.37). So we are left with proving (4.35).

Let $Z_N(s) := X_N(s) - X(s)$, then it satisfies (from (4.26) and (4.38)),

$$\begin{aligned} Z_N(t) &= \int_0^t a^{-1} \sin[a(t-s)] \Big[f_N(s) - f(s) - (M \star Z_N')(s) \\ &\quad - (M_N - M) \star X_N'(s) - x(M_N - M)(s) \Big] ds \ , \end{aligned}$$

and a similar formula holds $Z_N'(t)$. Clearly $Z_N(0) = Z_N'(0) = 0$. Hence, similarly to (4.33),

$$\begin{aligned} \mathbf{E}\Big(|Z_N(s)| + |Z_N'(s)| \Big) &\le K \int_0^t \widetilde{F}_N(s) ds \\ &\quad + a^{-1} t \sup_{s \le t} \left(\Big\{ |x| + t \sup_{u \le t} \mathbf{E}|X_N'(u)| \Big\} |M_N(s) - M(s)| + \mathbf{E}|f_N(s) - f(s)| \right) \ , \end{aligned}$$

with $\widetilde{F}_N(t) = \sup_{s \le t} \mathbf{E}|Z_N(s)| + \sup_{s \le t} \mathbf{E}|Z_N'(s)|$. We use again a Gronwall argument to obtain (4.35), based upon the control of $\sup_{u \le t} \mathbf{E}|X_N'(u)|$ from Lemma 4.1 and the facts that $|M_N(s) - M(s)| \to 0$ (see (4.29)) and $\mathbf{E}|f_N(s) - f(s)| \to 0$ uniformly for $s \le t$ as $N \to \infty$.

In order to check $\mathbf{E}|f_N(s) - f(s)| \to 0$, we observe that,

$$r_k = N^{1/2} r\Big(\Big[\frac{k-1}{N}, \frac{k}{N} \Big] \Big) = N^{1/2} \int \mathbf{1}\Big(\omega \in [\frac{k-1}{N}, \frac{k}{N}] \Big) r(d\omega) \ ,$$

to obtain,

$$\mathbf{E}|f(s) - f_N(s)|^2 = \lambda^2 \int_0^\Omega \left[A_\beta(\omega) - \sum_{k=1}^{N\Omega} A_\beta(\omega_k) \cdot \mathbf{1}\Big(\omega \in [\frac{k-1}{N}, \frac{k}{N}]\Big)\right]^2 d\omega \ , \tag{4.39}$$

which goes to zero as $N \to \infty$, uniformly in $s \leq t$. For uniformly spaced frequencies, $\omega_k = \frac{k}{N}$, (4.39) is straightforward. For frequencies satisfying only the uniform density condition (1.5) with $c = 1$, first one has to verify that

$$\lim_{N\to\infty} \frac{1}{N} \#\Big\{k \ : \ |\omega_k - \frac{k}{N}| \geq \eta\Big\} = 0$$

for any $\eta > 0$, and then using the continuity of the function $A_\beta(\omega)$ to conclude the result. ■

Let us remark that for the present paper there is no need to use stochastic integrals. A reader who is unfamiliar with this concept, can keep the finite sums $\sum_{k=1}^{N\Omega}$ instead of $\int_0^\Omega d\omega$, $f_N(t)$ instead of $f(t)$, and keep on thinking of $\mathbf{E}$ as expectation $\mathbf{E}_N$ with respect to the finite dimensional measure $d\mu_N$. We shall compute various expectations involving $f(t)$. The results are given as an ordinary $\int_0^\Omega(\ldots)d\omega$ integral. However, one can keep the finite dimensional approximations $f_N(t)$, and perform the expectations with respect to $d\mu_N$. In this case the expectations involve a finite sum over the frequencies, like $\sum_{k=1}^{N\Omega}(\ldots)$. It is sufficient to take the $N \to \infty$ limit only in this sum, which is a Riemann sum for the integral $\int_0^\Omega(\ldots)d\omega$ using (1.5) with $c = 1$. However, for notational simplicity we will use the continuous formalism. Note that the thermodynamic limit $N \to \infty$ is always taken before any other limits.

The conclusion of Section 4 is the,

Lemma 4.3 *Assume (1.5) with $c = 1$ and assume (4.10). Let $w_A^{N,\varepsilon}(t)$ be defined as (4.8), while $w^{N,\varepsilon}(t)$ is the solution of (4.11) with initial datum (4.9). Then, in the thermodynamic limit, we have for all $\phi(x,v) \in C_c^\infty(\mathbb{R}^2)$ locally uniformly for $t \in \mathbb{R}$,*

$$\lim_{N\to\infty} \int_{\mathbb{R}^2} w_A^{N,\varepsilon}(t,x,v)\overline{\phi}(x,v)dx\ dv = \int_{\mathbb{R}^2} w_A^{\varepsilon}(t,x,v)\overline{\phi}(x,v)dx\ dv \ , \tag{4.40}$$

where w_A^ε is defined by,

$$\int_{\mathbb{R}^2} w_A^{\varepsilon}(t,x,v)\overline{\phi}(x,v)dxdv = \tag{4.41}$$

$$= \mathbf{E}\int_{\mathbb{R}^6} \hat{w}_0(\xi,\eta)\overline{\hat{\phi}(\theta,\sigma)}e^{i(x\xi+v\eta)}e^{-i(X(t)\theta+X'(t)\sigma)}d\xi\ d\eta\ dx\ dv\ d\theta\ d\sigma \ ,$$

and X satisfies (4.36) .

For the proof one only has to observe that the dominated convergence theorem applies and use Lemma 4.2 and (4.12) (recalling that X is actually X_N in that formula). ■

Remark. As an alternative proof which avoids any reference to probabilistic concepts, we can easily compute the right-hand-side of (4.12) directly by performing a finite dimensional Gaussian integration with respect to $d\mu_N$ (again, $X(t)$ is actually $X_N(t)$ in (4.12)). In this case all the integrals $\int_0^{N\Omega}(\ldots)d\omega$ are finite sums and the $N \to \infty$ limit is taken only after having performed the $d\mu_N$ integration. We easily find that the right-hand-side of (4.12) is equal to,

$$\int_{\mathbb{R}^2} \hat{w_0}\Big(A(t)\theta + A'(t)\sigma\ ,\ B(t)\theta + B'(t)\sigma\Big)\ \overline{\hat{\phi}(\theta,\sigma)} \tag{4.42}$$
$$\times \exp\Big[-\int_0^\Omega \frac{[A_\omega(t)\theta + A'_\omega(t)\sigma]^2}{2\lambda_\omega}\,d\omega - \int_0^\Omega \frac{[B_\omega(t)\theta + B'_\omega(t)\sigma]^2}{2\mu_\omega}d\omega\Big]\ d\theta\ d\sigma\ ,$$

where $\lambda_\omega = [2\omega(\cosh(\beta\omega)-1)]/[\sinh(\beta\omega)]$, $\mu_\omega = [2\sinh(\beta\omega)]/[\omega(\cosh(\beta\omega)+1)]$, and,

$$\begin{aligned}
\Psi(t) &= \lambda^2 \int_0^\Omega \int_0^t \omega \sin(\omega[t-s])\sin(s)\ ds\ d\omega\ ,\\
A(t) &= \cos(t) + (\Psi \star A)(t)\ ,\\
B(t) &= \sin(t) + (\Psi \star B)(t)\ ,\\
A_\omega(t) &= -\int_0^t \lambda\omega\cos(\omega s)\sin(t-s)\ ds + (\Psi \star A_\omega)(t)\ ,\\
B_\omega(t) &= -\int_0^t \lambda\sin(\omega s)\sin(t-s)\ ds + (\Psi \star B_\omega)(t)\ .
\end{aligned}$$

5 The Fokker-Planck equation from the original Caldeira-Leggett model

5.1 Evolution without friction

In the spirit of [5], we would like to exhibit a scaling where the solution of (4.36) is close to the solution $\widetilde{X}(t)$ of the equation without friction term below. The scaling parameters are $\varepsilon = (\beta, \Omega, \lambda)$. The frictionless equation (compare with (4.36)) is,

$$\widetilde{X}''(t) + a^2\widetilde{X}(t) = f(t)\ , \qquad \text{with,} \qquad X(0) = x\ ,\quad X'(0) = v\ , \tag{5.1}$$

recalling that $a^2 = a_\varepsilon^2 = 1 - \lambda^2\Omega \in [\frac{1}{4}, 1]$.

We need a continuity result ensuring that $X(t)$ and $\widetilde{X}(t)$ are close. If $Y(t) = X(t) - \widetilde{X}(t)$, then,

$$Y''(t) + a^2 Y(t) = -(M \star X')(t) - xM(t)\ , \tag{5.2}$$

with initial conditions $Y(0) = Y'(0) = 0$. Given the bound (4.37) on $X(t)$ and (4.28) it is trivial to see that,

$$\mathbf{E}\Big(|Y(t)| + |Y'(t)|\Big) \leq Kte^{Kt}\left(|x| + |v| + K|x| + \sup_{s\leq t}\Big\{se^{-Ks}\Big\}\Big[\lambda^2\Omega\Big(\beta^{-1} + \Omega\Big)\Big]^{1/2}\right) , \quad (5.3)$$

where $K = C\lambda^2(1 + \frac{1}{|\Omega - a|})$ (see (4.25)). So in particular the solution of (4.36) tends to the solution of (5.1) in a very strong norm if the right-hand-side of (5.3) goes to zero. This happens for example for such limiting regimes of $\varepsilon = (\beta, \Omega, \lambda)$ that $\lambda \to 0$ and $\Omega \to \infty$ in such a way that $a^2 = 1 - \lambda^2\Omega \in [\frac{1}{4}, 1]$ and $\lambda^2\beta^{-1/2} \to 0$.

Hence, as soon as one can ensure a small right-hand-side in (5.3), we can replace X by $\widetilde{X}$ in (4.40)-(4.41) by the Lebesgue theorem, since the x, v, θ, σ integrations range over a bounded domain (ϕ is compactly supported) and we assumed $\widehat{w}_0(\xi, \eta) \in L^1$ (see (4.10)). This proves

Lemma 5.1 *Let $\widetilde{w}^\varepsilon_A$ be defined as,*

$$\int_{\mathbb{R}^2} \widetilde{w}^\varepsilon_A(t, x, v)\overline{\phi}(x, v)dxdv = \mathbf{E}\int_{\mathbb{R}^6} \hat{w}_0(\xi, \eta)\overline{\hat{\phi}(\theta, \sigma)}e^{i(x\xi + v\eta)}e^{-i(\widetilde{X}(t)\theta + \widetilde{X}'(t)\sigma)}d\xi\, d\eta\, dx\, dv\, d\theta\, d\sigma \ , \quad (5.4)$$

analogously to (4.41). Then,

$$\lim_\varepsilon \int_{\mathbb{R}^2} \widetilde{w}^\varepsilon_A(t, x, v)\overline{\phi}(x, v)dxdv = \lim_\varepsilon \int_{\mathbb{R}^2} w^\varepsilon_A(t, x, v)\overline{\phi}(x, v)dxdv \ , \quad (5.5)$$

for any limit of the parameters $\varepsilon = (\beta, \Omega, \lambda)$ for which the right hand side of (5.3) goes to zero. ■

5.2 Computing the dynamics of the test-particle when the memory vanishes

In this section we compute $w^\varepsilon(t, x, v)$ when X is actually replaced by $\widetilde{X}$, the solution of (5.1), in (4.41). We have,

$$\begin{aligned} \widetilde{X}(t) &= x\cos at + va^{-1}\sin at + \int_0^t a^{-1}\sin a(t-s)f(s)ds \ , \quad (5.6) \\ \widetilde{X}'(t) &= -xa\sin at + v\cos at + \int_0^t \cos[a(t-s)]f(s)ds \ . \end{aligned}$$

Hence

$$\begin{aligned} \int_{\mathbb{R}^2} \widetilde{w}^\varepsilon_A(t, x, v)\overline{\phi(x, v)}dx\, dv &= \mathbf{E}\int_{\mathbb{R}^6} \widehat{w}_0(\xi, \eta)\overline{\widehat{\phi}(\theta, \sigma)}e^{i(x\xi + v\eta)}e^{-i(\widetilde{X}(t)\theta + \widetilde{X}'(t)\sigma)}\, d\xi\, d\eta\, dx\, dv\, d\theta\, d\sigma \\ &= \mathbf{E}\int_{\mathbb{R}^2} \widehat{w}_0\Big(\xi_{\theta,\sigma}(t), \eta_{\theta,\sigma}(t)\Big)\overline{\widehat{\phi}(\theta, \sigma)}e^{-i\int_0^t \eta_{\theta,\sigma}(t-s)f(s)ds}\, d\theta\, d\sigma \ , \quad (5.7) \end{aligned}$$

with,

$$\eta_{\theta,\sigma}(t) := \theta a^{-1} \sin at + \sigma \cos at \ , \qquad \xi_{\theta,\sigma}(t) := \theta \cos at - \sigma a \sin at \ , \tag{5.8}$$

which are, by the way, harmonic oscillator trajectories,

$$\frac{d}{dt}\eta_{\theta,\sigma}(t) = \xi_{\theta,\sigma}(t) \ , \quad \frac{d}{dt}\xi_{\theta,\sigma}(t) = -a^2 \eta_{\theta,\sigma}(t) \ . \tag{5.9}$$

After performing the expectation in (5.7), we arrive at

Lemma 5.2 *With the notations above, we have for any $t \geq 0$,*

$$\int_{\mathbb{R}^2} \widetilde{w}_A^\varepsilon(t,x,v)\overline{\phi(x,v)}\, dx\, dv = \int_{\mathbb{R}^2} \widehat{w}_0\Big(\xi_{\theta,\sigma}(t), \eta_{\theta,\sigma}(t)\Big)\overline{\widehat{\phi}(\theta,\sigma)} e^{-\frac{1}{2}Q(t)}\, d\theta\, d\sigma \ , \tag{5.10}$$

with

$$Q(t) := Q(t;\theta,\sigma;\beta,a) = \lambda^2 \int_0^\Omega A_\beta^2(\omega) H(t,\omega) d\omega \ , \tag{5.11}$$

$$H(t,\omega) := H(t,\omega;\theta,\sigma;a) = \Big| \int_0^t \eta_{\theta,\sigma}(s) e^{-i\omega s} ds \Big|^2 \ . \tag{5.12}$$

The functions $\xi_{\theta,\sigma}$, $\eta_{\theta,\sigma}$ are defined by (5.8). The function $H(t,\omega)$ satisfies the following estimate

$$H(t,\omega) \leq 2\gamma^2 \left\{ \Big| \frac{e^{it(a-\omega)} - 1}{a-\omega} \Big|^2 + \frac{4}{(a+\omega)^2} \right\} \tag{5.13}$$

with $\gamma^2 := \theta^2 + a^2\sigma^2$. Assuming $\Omega > 1$ we also have

$$Q(t) = I\lambda^2 t \gamma^2 \frac{\cosh \beta a + 1}{2a \sinh \beta a} + \lambda^2 \gamma^2 B(t) \tag{5.14}$$

with $I := \frac{\pi}{2}$ and with a function B satisfying $B(0) = 0$ and

$$|B(t)| \leq C[1 + \beta^{-1}][1 + (\log t)_+][1 + \log \Omega] \tag{5.15}$$

with a universal constant C. Also, we have the estimate:

$$Q(t) = \mathbf{E}\Big(f \star \eta_{\theta,\sigma} \Big)^2(t) = \mathbf{E}\Big(\theta \widetilde{X}(t) + \sigma \widetilde{X}'(t) \Big)^2 + \mathcal{O}\Big[(|x| + |v|)(|\theta| + |\sigma|) \Big] \ . \tag{5.16}$$

Remark. Notice that $Q(t)$ grows quadratically in t for small t (since H does so). This means that the test-particle as described by the Wigner distribution w_A^ε has a ballistic behaviour when the memory effects disappear (quadratic growth of the mean squared displacement $\mathbf{E}\widetilde{X}^2(t)$). In

the rest of this paper we show that, under several specific scaling limits, one can indeed replace w_A^ε with $\widetilde{w}_A^\varepsilon$ (see Lemma 5.1) *and* recover a linear growth for $Q(t)$, i.e. a diffusive behaviour for the test-particle. In particular, this is where the time asymmetric condition $t \geq 0$ is used.

Proof. We only have to show the estimates (5.13) and (5.15). These are straightforward calculations. We use the following notation,

$$a\sigma + i\theta = \gamma e^{i\phi} \; . \tag{5.17}$$

(i.e. $\theta = \gamma \sin\phi$, $a\sigma = \gamma\cos\phi$ and $\gamma^2 = \theta^2 + a^2\sigma^2$). Hence, from (5.8),

$$\eta_{\theta,\sigma}(t) = \frac{\gamma}{2a}\Big(e^{i(\phi - at)} + e^{-i(\phi - at)}\Big) \; , \tag{5.18}$$

and

$$H(t,\omega) \;\; = \;\; \frac{\gamma^2}{4a^2}\Big|e^{2i\phi}\frac{e^{-it(a+\omega)}-1}{a+\omega} - \frac{e^{it(a-\omega)}-1}{a-\omega}\Big|^2 \; , \tag{5.19}$$

which proves (5.13).

To prove (5.14)-(5.15), for any $\Omega > 1$ we obtain, by extracting the worst singularity

$$\begin{aligned} Q(t) \;\; &= \;\; \lambda^2 \int_0^\Omega \frac{\omega(\cosh\beta\omega + 1)}{2\sinh\beta\omega} H(t,\omega) d\omega \\ &= \;\; \lambda^2 \frac{\gamma^2}{4a^2}\widetilde{B}(t) + \lambda^2 \frac{\gamma^2}{4a^2}\int_0^\Omega \frac{\omega(\cosh\beta\omega+1)}{2\sinh\beta\omega}\Big|\frac{e^{it(a-\omega)}-1}{a-\omega}\Big|^2 d\omega \; , \end{aligned} \tag{5.20}$$

with,

$$\widetilde{B}(t) := \int_0^\Omega \frac{\omega(\cosh\beta\omega+1)}{2\sinh\beta\omega}\Big\{\Big|\frac{e^{-it(a+\omega)}-1}{a+\omega}\Big|^2 - 2\mathrm{Re}\Big(e^{2i\phi}\frac{e^{-it(a+\omega)}-1}{a+\omega}\frac{e^{it(a-\omega)}-1}{a-\omega}\Big)\Big\}d\omega \; , \tag{5.21}$$

and $\widetilde{B}(0) = 0$. With the substitution $\omega' = t(a-\omega)$ in (5.21), one easily computes

$$|\widetilde{B}(t)| \leq C[1+\beta^{-1}][1+(\log t)_+][1+\log\Omega] \; . \tag{5.22}$$

The second integral in (5.20) is proportional to t for large t since $\Omega > 1$. Obviously it becomes uniformly bounded if $\Omega < a \leq 1$ (a trivial behaviour), and this is the very reason why we assumed $\Omega > 1$ in this section. Then the main contribution comes from $\omega \sim a$, and by the same change of variables as above, the result is,

$$Q(t) = \lambda^2\gamma^2 B(t) + I\lambda^2 t\gamma^2 \frac{\cosh a\beta + 1}{2a\sinh a\beta} \tag{5.23}$$

with $I := \frac{\pi}{2}$, and $\widetilde{B}(t)$ is replaced by some $B(t)$ which also satisfies (5.22) and $B(0) = 0$. ■

5.3 The Caldeira-Leggett limits: obtaining the Fokker-Planck equation

In this section we rigorously perform the scaling limit introduced in [5]. We prove the following,

Theorem 5.1 *Let w_A^ε be the Wigner distribution of the test-particle after the thermodynamic limit, as given by Lemma 4.3. We recall that ε stands for (β, Ω, λ). Let $\lambda = \lambda_0 \beta^{1/2}$ with some fixed λ_0, as in [5].*

a) [Nonzero frequency shift.] *Assume that $a^2 = 1 - \lambda^2 \Omega = 1 - \lambda_0^2 \beta \Omega \in [\frac{1}{4}, 1]$ is fixed. Then for any $t \geq 0$ the following weak limit exists*

$$W(t,x,v) = \lim_{\substack{\Omega \to \infty, \beta \to 0 \\ \beta\Omega = (1-a^2)\lambda_0^{-2}}} w_A^\varepsilon(t,x,v) \; . \tag{5.24}$$

The limit holds in the topology of $C^0([0,\infty)_t; \mathcal{D}'_{x,v})$. Moreover, W satisfies the Fokker-Planck equation,

$$\partial_t W + v\partial_x W - a^2 x \partial_v W - \frac{\lambda_0^2 \pi}{2} \Delta_v W = 0 \; , \tag{5.25}$$

with initial datum $W(t=0) = w_0$ satisfying (4.10)

b) [No frequency shift.] *For any $t \geq 0$ the following weak limit exists,*

$$W(t,x,v) = \lim_{\Omega \to \infty} \lim_{\beta \to 0} w_A^\varepsilon(t,x,v) \; . \tag{5.26}$$

[the order of limits cannot be interchanged], and W satisfies the Fokker-Planck equation,

$$\partial_t W + v\partial_x W - x \partial_v W - \frac{\lambda_0^2 \pi}{2} \Delta_v W = 0 \; , \tag{5.27}$$

with initial datum $W(t=0) = w_0$ satisfying (4.10)

Proof. For the proof of part a) first notice that Lemma 5.1 applies since the right hand side of (5.3) goes to zero under the prescribed limits. Hence X can be replaced by $\widetilde{X}$ and we can therefore rely on Lemma 5.2 above. On the other hand, since we assumed $\lambda = \lambda_0 \beta^{1/2}$, we readily observe,

$$\lim{}^* \, Q(t) = \lambda_0^2 \lim{}^* \int_0^\Omega \beta A_\beta^2(\omega) H(t,\omega) d\omega = \lambda_0^2 \int_0^\infty \Big| \int_0^t \eta_{\theta,\sigma}^2(s) e^{-i\omega s} ds \Big|^2 d\omega \; , \tag{5.28}$$

where $\lim^*$ stands for the simultaneous limit $\beta \to 0$, $\Omega \to \infty$ such that $a^2 = 1 - \lambda_0^2 \beta \Omega \in [\frac{1}{4}, 1]$ is fixed. Here we used that $\beta A_\beta(\omega)^2 \to 1$ in our limit if $\omega \leq \Omega^{1/2}$ and that $H(t,\omega) \in L^1(d\omega)$, see

(5.13). The contribution $\omega \geq \Omega^{1/2}$ to the integral vanishes in the limit by the estimate (5.13) and the trivial bound $\frac{z\cosh z+1}{\sinh z} \leq 2(1+z)$. Hence from the unitarity of the Fourier transform

$$\int_0^\infty \Big| \int_0^t g(s) e^{-i\omega s} ds \Big|^2 d\omega = \pi \int_0^t |g(s)|^2 ds \ , \tag{5.29}$$

which is valid for any real function g, we obtain

$$\lim{}^* \ Q(t) \ = \ \lambda_0^2 \pi \int_0^t \eta_{\theta,\sigma}^2(s) ds \ . \tag{5.30}$$

Here $t \geq 0$ is used, and this step is the origin of irreversibility. The end of the calculation is trivial. From Lemma 5.2 together with (5.30) we have,

$$\begin{aligned} \lim{}^* \int_{\mathbb{R}^2} w_A^\varepsilon(t,x,v) \overline{\phi(x,v)} \, dx \, dv \ &= \ \int_{\mathbb{R}^2} \widehat{w}_0 \Big(\xi_{\theta,\sigma}(t), \eta_{\theta,\sigma}(t) \Big) \\ &\quad \times \overline{\widehat{\phi}(\theta,\sigma)} e^{-I\lambda_0^2 \int_0^t \eta_{\theta,\sigma}^2(s) ds} \, d\theta \, d\sigma \ , \end{aligned} \tag{5.31}$$

where η and ξ are defined in (5.8) and $I = \frac{\pi}{2}$. We can define,

$$W(t,x,v) := \lim{}^* \ w_A^\varepsilon(t,x,v) \ , \tag{5.32}$$

as a weak limit given by (5.31). Then differentiating (5.31) gives (using (5.8)),

$$\begin{aligned} \int_{\mathbb{R}^2} \partial_t W(t,x,v) \overline{\phi(x,v)} dx \, dv \ &= \ \int_{\mathbb{R}^2} \partial_t \widehat{W}(t,\theta,\sigma) \overline{\widehat{\phi}(\theta,\sigma)} d\theta \, d\sigma \\ = \int_{\mathbb{R}^2} \Big[-a^2 \eta_{\theta,\sigma}(t) \partial_\xi + \xi_{\theta,\sigma}(t) \partial_\eta - I\lambda_0^2 \eta_{\theta,\sigma}^2(t) \Big] \widehat{w}_0 \Big(\xi_{\theta,\sigma}(t), \eta_{\theta,\sigma}(t) \Big) \overline{\widehat{\phi}(\theta,\sigma)} e^{-I\lambda_0^2 \int_0^t \eta_{\theta,\sigma}^2(s) ds} d\theta \, d\sigma \ . \end{aligned} \tag{5.33}$$

Letting $t = 0$, we have,

$$\partial_t \Big|_{t=0} \widehat{W}(t,\theta,\sigma) = \Big[-a^2 \sigma \partial_\theta + \theta \partial_\sigma - I\lambda_0^2 \sigma^2 \Big] \widehat{W}(t,\theta,\sigma) \Big|_{t=0} \ , \tag{5.34}$$

which is exactly the Fokker-Planck equation (5.27) after Fourier transforming,

$$\partial_t \Big|_{t=0} W(t,x,v) = \Big[a^2 x \partial_v - v \partial_x + I\lambda_0^2 \Delta_v \Big] W(t,x,v) \Big|_{t=0} \ . \tag{5.35}$$

Considering $t = 0$ is not a restriction, since the proof works for any L^1 initial condition.

The proof of part b) is completely analogous. We again notice that under the prescribed limits the right hand side of (5.3) goes to zero, hence Lemma 5.1 applies. Here $\eta_{\theta,\sigma}$ and $\xi_{\theta,\sigma}$ depend on the limiting parameters, since $a^2 = 1 - \lambda^2\Omega = 1 - \lambda_0^2 \beta \Omega$. But $\lim_{\beta \to 0} a = 1$, hence

$$\lim_{\beta \to 0} \eta_{\theta,\sigma}(s) = \theta \sin s + \sigma \cos s \ , \qquad \lim_{\beta \to 0} \xi_{\theta,\sigma}(s) = \theta \cos s - \sigma \sin s \tag{5.36}$$

uniformly for $s \in [0,t]$. Therefore

$$\begin{aligned}\lim_{\Omega\to\infty}\lim_{\beta\to 0} Q(t) &= \lambda_0^2 \lim_{\Omega\to\infty} \int_0^\Omega \Big| \int_0^t [\theta \sin s + \sigma \cos s] e^{-i\omega s} ds \Big|^2 d\omega \qquad (5.37)\\ &= \lambda_0^2 \int_0^\infty \Big| \int_0^t [\theta \sin s + \sigma \cos s] e^{-i\omega s} ds \Big|^2 d\omega \\ &= \pi \lambda_0^2 \int_0^t [\theta \sin s + \sigma \cos s]^2 ds \ .\end{aligned}$$

Again, the last step is robust in a sense that it does not use the particular form of the function $[\theta \sin s + \sigma \cos s]$, instead it uses (5.29). But it is rigid in a sense that $\Omega = \infty$ is essential to get diffusive (linear) behaviour for the mean square displacement (5.16).

To conclude, we follow the calculation (5.31)-(5.35). In addition to the limit (5.37), we have to replace $\xi_{\theta,\sigma}(s), \eta_{\theta,\sigma}(s)$ by their limiting values (5.36) in the argument of $\widehat{w}_0$ to arrive at the analogue of (5.31). Dominated convergence theorem applies if we assume, additionally, that $\widehat{w}_0$ is continuous and bounded. However $\widehat{w}_0 \in L^1$, hence it can be approximated by such functions in L^1-norm. Using that the flow $(\theta,\sigma) \mapsto \Big(\xi_{\theta,\sigma}(s), \eta_{\theta,\sigma}(s)\Big)$ is measure preserving and that $\widehat{\phi}$ is bounded, one can easily see that the approximation error can be made arbitrarily small.

The rest of the calculation is identical to the proof of part a) and we obtain (5.27). ■

6 Scaling limit at high temperature: the frictionless heat equation

We propose a different way to get diffusion from the Hamiltonian (2.1). As we mentioned, obtaining diffusion for the test-particle means that we have to extract linear dependence in time for $Q(t)$. In this section, linear growth is obtained from time rescaling and from the special form of linear combinations of $\sin s$ and $\cos s$ in Lemma 5.2. It relies on a resonance effect which comes from a singularity near $\omega \sim a$. The system $\widetilde{X}''(t) + a^2 \widetilde{X}(t)$ (see (5.1)) picks up those modes from the forcing term $f(t)$ in (4.34) for which the frequency ω is close to its eigenfrequency. So, in this section we assume $\Omega > 1$ but finite, contrary to the previous section.

This effect is more robust (see the remark after (5.37)) in the sense that one *could* leave the hyperbolic functions βA_β^2 in (5.28) without ensuring a limit where it goes to 1. In other terms, we do not need the high temperature limit $\beta \to 0$ to obtain diffusion, unlike in Section 5.3, where this limit made the $d\omega$ measure uniform and we recovered a white noise forcing term.

Nevertheless, Lemma 5.2 needs the right-hand-side of (5.3) to go to zero in order to be applicable (one needs the friction to vanish), and this cannot be achieved keeping β fixed (Section 2.4), hence we again set $\lambda = \lambda_0 \beta^{1/2}$, $\beta \to 0$.

6.1 Large space/time convergence of the Wigner distribution

Let α be a small parameter. We describe the behaviour of the test-particle, as given by its Wigner distribution w_A^ε on time scales of order $1/\alpha^2$. We consider the diffusive scaling, i.e. the space coordinate scales as $1/\alpha$. Since the test-particle is a fast harmonic oscillator, and energies are transferred back and forth between space and velocity, we also have to consider velocities of order $1/\alpha$. Hence we introduce the following scaling,

$$t = T\alpha^{-2}, \qquad x = X\alpha^{-1}, \qquad v = V\alpha^{-1} \; , \tag{6.1}$$

where the capital letters are unscaled quantities (macroscopic variables). The rescaled reduced Wigner transform is defined as,

$$W_T^{\varepsilon,\alpha}(X,V) := w_A^\varepsilon(T\alpha^{-2}, X\alpha^{-1}, V\alpha^{-1}) \; , \tag{6.2}$$

where w_A^ε is defined in Lemma 4.3 (after the thermodynamic limit). Its Fourier transform is,

$$\widehat{W}_T^{\varepsilon,\alpha}(\Theta,\Sigma) = \alpha^2 \widehat{w}_A^\varepsilon(T\alpha^{-2}, \Theta\alpha, \Sigma\alpha) \; , \tag{6.3}$$

where we use $\Theta = \theta\alpha^{-1}$ and $\Sigma = \sigma\alpha^{-1}$ rescaled dual variables. The initial condition is,

$$W_{T=0}^{\varepsilon,\alpha}(X,V) = W_0(X,V) \; , \qquad \widehat{W}_{T=0}^{\varepsilon,\alpha}(\Theta,\Sigma) = \widehat{W}_0(\Theta,\Sigma) \; , \tag{6.4}$$

and we assume that,

$$\widehat{W}_0(\Theta,\Sigma) \in L^1(\mathbb{R}_\Theta \times \mathbb{R}_\Sigma) \; . \tag{6.5}$$

The macroscopic testfunction $\Phi(X,V)$ is a smooth function with compact support, the microscopic testfunction is defined as,

$$\phi(x,v) = \Phi(x\alpha, v\alpha) = \Phi(X,V) \; , \tag{6.6}$$

and in Fourier variables, $\widehat{\phi}(\theta,\sigma) = \alpha^{-2}\widehat{\Phi}(\theta\alpha^{-1}, \sigma\alpha^{-1}) = \alpha^{-2}\widehat{\Phi}(\Theta,\Sigma)$.

We are now in position to state the theorem of this section,

Theorem 6.1 *Define the large time/space scale Wigner distribution $W_T^{\varepsilon,\alpha}(X,V)$ as in (6.2). Assume (6.5) for the initial data. Assume that $\lambda = \lambda_0\beta^{1/2}$ with a fixed $\lambda_0 > 0$ and fix the frequency cutoff $\Omega > 1$. Hence the limits of the parameters $\varepsilon = (\beta, \Omega, \lambda)$ are reduced to $\beta \to 0$. Then:*

a) *The following high-temperature limit exists in the weak sense for any $T \geq 0$:*

$$W_T^\alpha(X,V) := \lim_{\beta\to 0} W_T^{\varepsilon,\alpha}(X,V) \; . \tag{6.7}$$

b) *Define the following time average of W^α over one cycle of the harmonic oscillator (5.8),*

$$W_T^{\#,\alpha}(X,V) := \frac{1}{2\pi\alpha^2}\int_T^{T+2\pi\alpha^2} W_S^\alpha(X,V)dS\ . \tag{6.8}$$

Then the weak limit,

$$W_T^+(X,V) := \lim_{\alpha\to 0} W_T^{\#,\alpha}(X,V)\ , \tag{6.9}$$

exists for each $T \geq 0$ and it satisfies the heat equation in phase space,

$$\partial_T W_T^+ = \frac{\pi\lambda_0^2}{4}(\Delta_X + \Delta_V)W_T^+\ , \tag{6.10}$$

with initial condition $W_{T=0}^+(X,V)$ given by

$$\widehat{W}_0^+(X,V) = \frac{1}{2\pi}\int_0^{2\pi} \widehat{W}_0\Big(X\sin s + V\cos s,\ X\cos s - V\sin s\Big)ds\ . \tag{6.11}$$

c) *Define the radial average,*

$$W_T^{*,\alpha}(X,V) := \frac{1}{2\pi}\int_0^{2\pi} W_T^\alpha(R\cos s, R\sin s)ds \tag{6.12}$$

with $R := \sqrt{X^2+V^2}$, and clearly $W_T^{*,\alpha}$ depends on R only. Again, the weak limit,

$$W_T^\dagger(X,V) := \lim_{\alpha\to 0} W_T^{*,\alpha}(X,V)\ , \tag{6.13}$$

exists and the radially symmetric function $W_T^\dagger$ satisfies the heat equation (6.10) with initial condition,

$$W_{T=0}^\dagger(X,V) := \frac{1}{2\pi}\int_0^{2\pi} W_0(R\cos s, R\sin s)ds\ .$$

Remark. Here we identified the equation in a weak sense in the space and velocity variables, but in a strong sense in the time variable and some averaging ((6.8) or (6.12)) was needed to ensure the existence of the limit. If we want to consider the limit in a weak sense in time as well, then there is no need for averaging. Based upon part b), one can easily prove that $W_T^+(X,V)$ can also be identified as the weak limit in space, velocity and time, i.e. we have the following

Corollary 6.1 *Under the above conditions the weak limit*

$$W_T^+(X,V) := \lim_{\alpha\to 0}\lim_{\beta\to 0} W_T^{\varepsilon,\alpha}(X,V)$$

exists in the topology of $\mathcal{D}'(\ [0,\infty)_T \times \mathbb{R}_X \times \mathbb{R}_V\ )$, it coincides with (6.9) and satisfies (6.10). ■

Proof of Theorem 6.1. Using the rescaling and the definition of w_A^ε (4.41), we have,

$$\begin{aligned}\langle W_T^{\varepsilon,\alpha},\Phi\rangle &= \int_{\mathbb{R}^2} W_T^{\varepsilon,\alpha}(X,V)\overline{\Phi(X,V)}dX\,dV = \alpha^2\int_{\mathbb{R}^2} w_A^\varepsilon(T\alpha^{-2},x,v)\overline{\phi(x,v)}dx\,dv \\ &= \alpha^2\,\mathbf{E}\int_{\mathbb{R}^6}\widehat{w}_0(\xi,\eta)\overline{\widehat{\phi}(\theta,\sigma)}e^{i(x\xi+v\eta)}e^{-i(\theta X(t)+\sigma X'(t))}d\xi\,d\eta\,dx\,dv\,d\theta\,d\sigma \qquad (6.14)\\ &= \mathbf{E}\int_{\mathbb{R}^6}\widehat{W}_0(\xi\alpha^{-1},\eta\alpha^{-1})\overline{\widehat{\Phi}(\Theta,\Sigma)}e^{i(x\xi+v\eta)}e^{-i\alpha(\Theta X(t)+\Sigma X'(t))}d\xi\,d\eta\,dx\,dv\,d\Theta\,d\Sigma\,,\end{aligned}$$

where $t = T\alpha^{-2}$.

First Step: the limit $\beta \to 0$.

Due to the choice $\lambda = \lambda_0\beta^{1/2}$, we can replace $X(t)$ by $\widetilde{X}(t)$ in the $\beta\to 0$ limit. For, the right hand side of (5.3) goes to zero as $\beta\to 0$, hence Lemma 5.1 applies. Hence,

$$\begin{aligned}&\lim_{\beta\to 0}\langle W_T^{\varepsilon,\alpha},\Phi\rangle = \\ &= \lim_{\beta\to 0}\mathbf{E}\int_{\mathbb{R}^6}\widehat{W}_0(\xi\alpha^{-1},\eta\alpha^{-1})\overline{\widehat{\Phi}(\Theta,\Sigma)}e^{i(x\xi+v\eta)}e^{-i\alpha(\Theta\widetilde{X}(t)+\Sigma\widetilde{X}'(t))}d\xi\,d\eta\,dx\,dv\,d\Theta\,d\Sigma \\ &= \lim_{\beta\to 0}\mathbf{E}\int_{\mathbb{R}^2}\widehat{W}_0\Big(\xi_{\Theta,\Sigma}(T\alpha^{-2}),\eta_{\Theta,\Sigma}(T\alpha^{-2})\Big)\overline{\widehat{\Phi}(\Theta,\Sigma)}e^{-\frac{1}{2}Q(T\alpha^{-2})}\,d\Theta\,d\Sigma\,, \qquad (6.15)\end{aligned}$$

where in the second step we also used Lemma 5.2 and the fact that $\alpha^{-1}\xi_{\alpha\Theta,\alpha\Sigma} = \xi_{\Theta,\Sigma}$ and $\alpha^{-1}\eta_{\alpha\Theta,\alpha\Sigma} = \eta_{\Theta,\Sigma}$ (see (5.8)).

Recall that both $Q(t)$ and the trajectories $\xi_{\Theta,\Sigma},\eta_{\Theta,\Sigma}$ depend on β, since $a^2 = 1-\lambda^2\Omega = 1-\lambda_0^2\beta\Omega$ appears in their definition (see (5.8)). Similarly to the argument at the end of the proof of part b) of Theorem 5.1, using that $\widehat{W}_0 \in L^1(d\Theta\,d\Sigma)$, $\widehat{\Phi}\in L^\infty\cap C^0$, $Q\geq 0$, we see that the limit can be taken inside the integral and the trajectories $\xi_{\Theta,\Sigma},\eta_{\Theta,\Sigma}$ can be replaced by their limiting values (as $a\to 1$)

$$\eta^*_{\Theta,\Sigma}(s) := \theta\sin t + \sigma\cos t \qquad \xi^*_{\Theta,\Sigma}(s) := \theta\cos t - \sigma\sin t\,. \qquad (6.16)$$

We also use (see (5.14)) that

$$\lim_{\beta\to 0} Q(t) = I\lambda_0^2 t\gamma^2 + \lambda_0^2\gamma^2 B_0(t)\,. \qquad (6.17)$$

with $B_0(t)$ satisfying $B_0(0)=0$ and

$$|B_0(t)| \leq C[1+(\log t)_+][1+\log\Omega] \qquad (6.18)$$

(see (5.15)). We also recall that $\gamma^2 = \theta^2 + \sigma^2 = \alpha^2(\Theta^2 + \Sigma^2) =: \alpha^2\Gamma^2$. Hence,

$$\begin{aligned}\lim_{\beta\to 0}\langle W_T^{\varepsilon,\alpha}, \Phi\rangle &= \int_{\mathbb{R}^2} \widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(T\alpha^{-2}), \eta^*_{\Theta,\Sigma}(T\alpha^{-2})\Big) \times \\ &\times\overline{\widehat{\Phi}(\Theta,\Sigma)}\exp\Big\{-\frac{1}{2}\Big[I\lambda_0^2 T\alpha^{-2} + \lambda_0^2 B_0(T\alpha^{-2})\Big]\alpha^2(\Theta^2+\Sigma^2)\Big\}d\Theta\, d\Sigma\ .\end{aligned} \tag{6.19}$$

This relation defines the Fourier transform,

$$\widehat{W}_T^{\alpha}(\Theta,\Sigma) := \lim_{\beta\to 0}\widehat{W}_T^{\varepsilon,\alpha}(\Theta,\Sigma)\ , \tag{6.20}$$

as a weak limit, and its inverse Fourier transform,

$$W_T^{\alpha}(X,V) := \lim_{\beta\to 0} W_T^{\varepsilon,\alpha}(X,V)\ .$$

We can compute its time derivative in Fourier space,

$$\begin{aligned}\langle\partial_T\widehat{W}_T^{\alpha}, \widehat{\Phi}\rangle &= \int \alpha^{-2}\Big[-\eta^*_{\Theta,\Sigma}(T\alpha^{-2})\partial_\xi + \xi^*_{\Theta,\Sigma}(T\alpha^{-2})\partial_\eta - \\ &-\frac{\alpha^2}{2}\Big[I\lambda_0^2 + \lambda_0^2 B_0'(T\alpha^{-2})\Big](\Theta^2+\Sigma^2)\Big]\widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(T\alpha^{-2}), \eta^*_{\Theta,\Sigma}(T\alpha^{-2})\Big) \\ &\times\overline{\widehat{\Phi}(\Theta,\Sigma)}\exp\Big\{-\frac{1}{2}\Big[I\lambda_0^2 T\alpha^{-2} + \lambda_0^2 B_0(T\alpha^{-2})\Big]\alpha^2(\Theta^2+\Sigma^2)\Big\}d\Theta\, d\Sigma\ .\end{aligned} \tag{6.21}$$

As usual, we can let $T = 0$ to obtain,

$$\partial_T\Big|_{T=0}\widehat{W}_T^{\alpha}(\Theta,\Sigma) = \alpha^{-2}\Big[-\Sigma\partial_\Theta + \Theta\partial_\Sigma - \frac{\alpha^2}{2}\Big[I\lambda_0^2 + \lambda_0^2 B_0'(0)\Big](\Theta^2+\Sigma^2)\Big]\widehat{W}_0(\Theta,\Sigma)\ . \tag{6.22}$$

Second Step: the macroscopic limit $\alpha \to 0$.

Now the difficulty in (6.22) is that the convective term is too big compared to the last diffusive term since the motion takes place on two different time scales. There is the fast (microscopic) time scale of the harmonic oscillator described by $\alpha^{-2}[-\Sigma\partial_\Theta + \Theta\partial_\Sigma]$. Then there is a slow, macroscopic diffusive scale. We present two ways to average out the fast motion.

Part b) of Theorem 6.1: Averaging over a cycle.

Here we define $W^{\#,\alpha}$ according to (6.8). Now for any T fixed the formula,

$$\lim_{\alpha\to 0}\langle\widehat{W}_T^{\#,\alpha}, \widehat{\Phi}\rangle = \lim_{\alpha\to 0}\int \widehat{W}_T^{\#,\alpha}(\Theta,\Sigma)\overline{\widehat{\Phi}(\Theta,\Sigma)}d\Theta d\Sigma \tag{6.23}$$

$$= \lim_{\alpha\to 0}\int\Big[\frac{1}{2\pi\alpha^2}\int_T^{T+2\pi\alpha^2}\widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(S\alpha^{-2}), \eta^*_{\Theta,\Sigma}(S\alpha^{-2})\Big)e^{-I_1\lambda_0^2 S(\Theta^2+\Sigma^2)}dS\Big]\overline{\widehat{\Phi}(\Theta,\Sigma)}d\Theta d\Sigma\ ,$$

defines a function,

$$\widehat{W}_T^+(\Theta,\Sigma) := \lim_{\alpha\to 0} \widehat{W}_T^{\#,\alpha}(\Theta,\Sigma)\ , \tag{6.24}$$

weakly, as we show below. Here $I_1 := \frac{I}{2} = \frac{\pi}{4}$ for brevity. Note that in (6.23) we neglected the term involving B_0 in the exponential (see (6.19)) since the estimate (6.18) readily implies $\alpha^2 B_0(T\alpha^{-2}) \to 0$. The exponential factor in (6.19) converges to that in (6.23) uniformly for all $S \le T$. Using $\widehat{\Phi} \in L^1$, we can apply the dominated convergence theorem along with approximating $\widehat{W}_0$ by bounded functions, similarly to the argument at the end of the proof of Theorem 5.1.

We have to show that the limit on the right-hand-side of (6.23) exists,

$$\begin{aligned}\langle \widehat{W}_T^{\#,\alpha}, \widehat{\Phi}\rangle \;=\; & \int_{\mathbb{R}^2}\Big[\frac{1}{2\pi\alpha^2}\int_T^{T+2\pi\alpha^2} \widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(S\alpha^{-2}),\eta^*_{\Theta,\Sigma}(S\alpha^{-2})\Big)e^{-I_1\lambda_0^2 T(\Theta^2+\Sigma^2)}dS \\ & + \frac{1}{2\pi\alpha^2}\int_T^{T+2\pi\alpha^2} \widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(S\alpha^{-2}),\eta^*_{\Theta,\Sigma}(S\alpha^{-2})\Big) \\ & \times\Big[e^{-I_1\lambda_0^2 S(\Theta^2+\Sigma^2)} - e^{-I_1\lambda_0^2 T(\Theta^2+\Sigma^2)}\Big]dS\Big]\overline{\widehat{\Phi}(\Theta,\Sigma)}d\Theta d\Sigma\ .\end{aligned} \tag{6.25}$$

The first term in (6.25) is independent of α, as it is just the integral of $\widehat{W}_0(\xi^*(s),\eta^*(s))$ over one full cycle of the harmonic oscillator (6.16),

$$\frac{1}{2\pi\alpha^2}\int_T^{T+2\pi\alpha^2} \widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(S\alpha^{-2}),\eta^*_{\Theta,\Sigma}(S\alpha^{-2})\Big)dS = \frac{1}{2\pi}\int_0^{2\pi} \widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(s),\eta^*_{\Theta,\Sigma}(s)\Big)ds\ . \tag{6.26}$$

The second term in (6.25) vanishes in the limit $\alpha \to 0$ since,

$$\Big|e^{-I_1\lambda_0^2 S(\Theta^2+\Sigma^2)} - e^{-I_1\lambda_0^2 T(\Theta^2+\Sigma^2)}\Big| \le 2\pi I_1\lambda_0\alpha^2(\Theta^2+\Sigma^2)e^{-I_1\lambda_0^2 T(\Theta^2+\Sigma^2)} \tag{6.27}$$

(use that $|S-T| \le 2\pi\alpha^2$), which kills the factor α^{-2} in (6.25) and then the length of the integration interval goes to zero. Dominated convergence theorem again has to be applied after an approximation. This shows that the limit in (6.24) makes sense and,

$$\begin{aligned}\langle W_T^+,\Phi\rangle \;&=\; \langle \widehat{W}_T^+,\widehat{\Phi}\rangle \\ &= \int_{\mathbb{R}^2}\Big[\frac{1}{2\pi}\int_0^{2\pi} \widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(s),\eta^*_{\Theta,\Sigma}(s)\Big)ds\Big]e^{-I_1\lambda_0^2 T(\Theta^2+\Sigma^2)}\overline{\widehat{\Phi}(\Theta,\Sigma)}d\Theta d\Sigma\ .\end{aligned} \tag{6.28}$$

The time derivative is,

$$\begin{aligned}&\langle \partial_T W_T^+,\Phi\rangle = \\ =\;& -I_1\lambda_0^2\int_{\mathbb{R}^2}(\Theta^2+\Sigma^2)\Big[\frac{1}{2\pi}\int_0^{2\pi} \widehat{W}_0\Big(\xi^*_{\Theta,\Sigma}(s),\eta^*_{\Theta,\Sigma}(s)\Big)ds\Big]e^{-I_1\lambda_0^2 T(\Theta^2+\Sigma^2)}\overline{\widehat{\Phi}(\Theta,\Sigma)}d\Theta d\Sigma \\ =\;& -I_1\lambda_0^2\Big\langle \widehat{W}_T^+,(\Theta^2+\Sigma^2)\widehat{\Phi}\Big\rangle = -I_1\lambda_0^2\Big\langle W_T^+,-(\Delta_X+\Delta_V)\Phi\Big\rangle\ ,\end{aligned} \tag{6.29}$$

which completes the proof of (6.10). The initial condition (6.11) is easily obtained from (6.28) by setting $T = 0$ and taking inverse Fourier transform.

Part c) of Theorem 6.1: Radial average

The other possibility to eliminate the fast modes is to use the radial function $W_T^{*,\alpha}$ defined in (6.12). Now the formula,

$$\lim_{\alpha\to 0}\langle \widehat{W}_T^{*,\alpha}, \widehat{\Phi}\rangle = \lim_{\alpha\to 0}\int \widehat{W}_T^{*,\alpha}(\Theta,\Sigma)\overline{\widehat{\Phi}(\Theta,\Sigma)}\, d\Theta\, d\Sigma \tag{6.30}$$

$$= \lim_{\alpha\to 0}\int \left[\frac{1}{2\pi}\int_0^{2\pi} \widehat{W}_0\Big(\xi^*_{\Gamma\cos s,\Gamma\sin s}(T\alpha^{-2}), \eta^*_{\Gamma\cos s,\Gamma\sin s}(T\alpha^{-2})\Big) ds\right] e^{-I_1\lambda_0^2 T(\Theta^2+\Sigma^2)}\overline{\widehat{\Phi}(\Theta,\Sigma)}\, d\Theta\, d\Sigma\ ,$$

(with $\Gamma := \sqrt{\Theta^2+\Sigma^2}$) defines a radial function,

$$\widehat{W}_T^{\dagger}(\Theta,\Sigma) := \lim_{\alpha\to 0}\widehat{W}_T^{*,\alpha}(\Theta,\Sigma)\ , \tag{6.31}$$

(depending only on $\Theta^2+\Sigma^2$) as a weak limit, as we show below. Note that in (6.30) we again neglected the term involving B_0 in the exponential for the same reason as in (6.23).

We have to show that the limit on the right-hand-side of (6.30) exists. But,

$$\xi^*_{\Gamma\cos s,\Gamma\sin s}(T\alpha^{-2}) = \Gamma\cos(s+T\alpha^{-2})\ , \qquad \eta^*_{\Gamma\cos s,\Gamma\sin s}(T\alpha^{-2}) = \Gamma\sin(s+T\alpha^{-2})\ ,$$

hence,

$$\frac{1}{2\pi}\int_0^{2\pi} \widehat{W}_0\Big(\xi^*_{\Gamma\cos s,\Gamma\sin s}(T\alpha^{-2}), \eta^*_{\Gamma\cos s,\Gamma\sin s}(T\alpha^{-2})\Big) ds$$
$$= \frac{1}{2\pi}\int_0^{2\pi} \widehat{W}_0(\Gamma\cos s,\Gamma\sin s) ds =: \widehat{W}_0^{\dagger}(\Theta,\Sigma)\ ,$$

independently of α, which is the "radialized" initial condition in Fourier space.

So it is clear that the limit on the right-hand-side of (6.30) exists,

$$\lim_{\alpha\to 0}\langle \widehat{W}_T^{*,\alpha}, \widehat{\Phi}\rangle = \int \widehat{W}_0^{\dagger}(\Theta,\Sigma) e^{-I_1\lambda_0^2 T(\Theta^2+\Sigma^2)}\overline{\widehat{\Phi}(\Theta,\Sigma)}\, d\Theta\, d\Sigma =: \langle \widehat{W}_T^{\dagger}, \widehat{\Phi}\rangle\ ,$$

and clearly $W_T^{\dagger}$ also satisfies the heat equation (6.10). This ends the proof of Theorem 6.1. ■

7 Heat equation with friction at finite temperature

Here we choose a scaling where the Markovian part of the friction term does not vanish, i.e. we can keep β fixed and still get finite diffusion. Again we look at large time $t = T\delta^{-1}$ but now we do not scale the space variable. To eliminate the fast mode, we again integrate the angle.

The result is a radial Fokker-Planck equation with friction. While the test-particle performs many cycles, it slowly diffuses out, and this diffusion is slowed down by a friction. The diffusion comes from resonance.

In this scaling limit the solution of (4.36) is close to the solution $\widetilde{X}(t)$ of an equation without a time delayed (non-Markovian) friction term, but a Markovian friction term will be present. Let us choose,

$$\lambda := \lambda_0 \delta^{1/2} \ , \tag{7.1}$$

with some $\lambda_0 \leq \sqrt{3}/2$ fixed. We compare the solution of (4.36) to that of

$$\widetilde{X}''(t) + I\lambda^2 \widetilde{X}'(t) + a^2 \widetilde{X}(t) = f(t) \ ; \qquad \widetilde{X}(0) = x \ , \quad \widetilde{X}'(0) = v \ , \tag{7.2}$$

with $a^2 := 1 - \lambda^2 \Omega = 1 - \lambda_0^2 \delta^{-1} \Omega$, and,

$$I = \int_0^\infty \frac{\sin \Omega s}{s} ds = \frac{\pi}{2} \ . \tag{7.3}$$

We choose the scaling such that $a \in [\frac{1}{2}, 1]$, hence we always assume that $\Omega \leq \delta^{-1}$, but to exploit resonance, we also assume $\Omega > 2$. The new term $\lambda^2 I \widetilde{X}'(t)$ for the approximate characteristic is due to the fact that $M(t) \sim \lambda^2 I \delta_0(t)$ as $\Omega \to 0$, where δ_0 denotes the Dirac delta measure. This term is the main part of the full friction $(M \star X')$ in (4.36). Notice that it is small compared with the pure harmonic oscillator terms, $\widetilde{X}'' + a^2 \widetilde{X}$, but it is not negligible, since we will consider long times $t \sim \lambda^{-2}$.

7.1 A priori bounds and continuity results

As in Section 5.1 we need a priori estimates for X, i.e. for,

$$F(t) := \sup_{s \leq t} \mathbf{E}|X(s)| + \sup_{s \leq t} \mathbf{E}|X'(s)| \ ,$$

and estimates on the difference between $\widetilde{X}(t)$ and $X(t)$. The estimate (4.37) in Lemma 4.2 (which originates in (4.24) in Lemma 4.1), however, is not precise enough for large times. The following estimate is a more precise version of Lemma 4.2.

Lemma 7.1 *Let $t = T\delta^{-1}$, $\lambda = \lambda_0 \delta^{1/2}$ with fixed $\lambda_0 \leq \sqrt{3}/2$ and $T \geq 0$ and we assume that $2 \leq |\log \delta|^7 \leq \Omega \leq \delta^{-1}$ We also fix $\beta > 0$, hence the limit of scaling parameters $\varepsilon = (\beta, \Omega, \lambda)$ is reduced to $\delta \to 0$, $\Omega \to \infty$ with the side condition that $\Omega \in [|\log \delta|^7, \delta^{-1}]$.*

Let X be the solution to (4.36), then,

$$F(T\delta^{-1}) \leq C(\beta, \lambda_0, T)\Big(1 + |x| + |v|\Big) \ , \tag{7.4}$$

where C is monotone increasing in T. Moreover, if $\widetilde{X}$ is the solution to (7.2), then the difference $Y(t) =: X(t) - \widetilde{X}(t)$ satisfies,

$$\lim_{\delta\to 0}\Big(\sup_{s\le T\delta^{-1}}\mathbf{E}|Y(s)| + \sup_{s\le T\delta^{-1}}\mathbf{E}|Y'(s)|\Big) = 0\ . \tag{7.5}$$

In particular,

$$\lim_{\delta\to 0}\int_{\mathbb{R}^2}\widetilde{w}_A^\varepsilon(s,x,v)\overline{\phi}(x,v)dxdv = \lim_{\delta\to 0}\int_{\mathbb{R}^2} w_A^\varepsilon(s,x,v)\overline{\phi}(x,v)dxdv\ , \tag{7.6}$$

uniformly for all $s \le T\delta^{-1}$, where $\widetilde{w}_A^\varepsilon(t,x,v)$ is the Wigner transform corresponding to $\widetilde{X}$, defined exactly as (5.4), but $\widetilde{X}(t)$ now being the solution to (7.2).

Proof. We follow essentially the proof of Lemma 4.1. The characteristics (4.36) fulfill

$$\begin{aligned} X(t) &= x\cos at + va^{-1}\sin at + \int_0^t a^{-1}\sin a(t-s)\Big[f(s) - (M\star X')(s) - xM(s)\Big]ds \\ X'(t) &= -xa\sin at + v\cos at + \int_0^t \cos a(t-s)\Big[f(s) - (M\star X')(s) - xM(s)\Big]ds\ . \end{aligned} \tag{7.7}$$

Similarly to the proof of (4.30) one obtains

$$\mathbf{E}\Big|\int_0^t a^{-1}\sin a(t-s)\Big[(M\star X')(s) + xM(s)\Big]ds\Big| \le K\Big[\int_0^t F(s)ds + |x|\Big]\ , \tag{7.8}$$

recalling the value of K (4.25), and the cosine term in $X'(t)$ is similar.

Now we estimate the random forcing term. First we use

$$\mathbf{E}\Big|\int_0^t f(s)\ a^{-1}\sin a(t-s)\ ds\Big| \le \left(\mathbf{E}\Big|\int_0^t f(s)\ a^{-1}\sin a(t-s)\ ds\Big|^2\right)^{1/2}\ , \tag{7.9}$$

then notice that $a^{-1}\sin a(t-s) = \eta_{\theta,\sigma}(t-s)$ with $\theta = 1$, $\sigma = 0$ (see (5.8)). Hence (cf. (5.12))

$$\mathbf{E}\Big|\int_0^t f(s)a^{-1}\sin a(t-s)\ ds\Big|^2 \le \lambda^2\int_0^\Omega A_\beta^2(\omega)H(t,\omega;1,0;a) \tag{7.10}$$

which is just $Q(t) = Q(t;1,0;\beta,a)$, see (5.11). Hence from (5.14), (5.15) we get

$$\mathbf{E}\Big|\int_0^t f(s)\ a^{-1}\sin a(t-s)\ ds\Big|^2 \le C_1^2(\beta,\lambda_0,T) \tag{7.11}$$

using the relations among the parameters; $t = T\delta^{-1}$, $\lambda = \lambda_0\delta^{1/2}$ and $\Omega \le \delta^{-1}$. Similar estimate is valid for the cosine term.

The estimates (7.8), (7.9) and (7.11) lead to the a priori bound,

$$F(t) \le |x| + |v| + K\Big[\int_0^t F(s)ds + |x|\Big] + C_1(\beta, \lambda_0, T) \ , \tag{7.12}$$

and by the standard Gronwall argument we obtain,

$$F(t) \ \ \le \ \ C_2(\beta, \lambda_0, T)\Big(1 + |x| + |v|\Big) \ . \tag{7.13}$$

By monotonicity of C_2 in T, we get the a priori bound (7.4) on $X(t)$ and $X'(t)$.

From the equation (4.36) we also get a similar bound for $X''(t)$. We estimate

$$\mathbf{E}|X''(t)| \ \ \le \ \ a^2\mathbf{E}|X(t)| + \Big(\mathbf{E}|f(t)|^2\Big)^{1/2} + |x||M(t)| + \int_0^t |M(s)| \ \mathbf{E}|X'(t-s)|ds \ .$$

For the forcing term we use

$$\mathbf{E}|f(t)|^2 = \lambda^2 \int_0^\Omega \frac{\omega(\cosh\beta\omega + 1)}{2\sinh\beta\omega} \, d\omega \le C_3(\beta)\lambda^2\Omega^2$$

(see (4.32)) and that

$$|M(s)| = \lambda^2\Big|\frac{\sin\Omega s}{s}\Big| \le \frac{2\Omega\lambda^2}{1+\Omega s} \ . \tag{7.14}$$

These estimates, along with $t = T\delta^{-1}$, $\lambda = \lambda_0\delta^{1/2}$ and $\Omega \le \delta^{-1}$, give that

$$\sup_{s \le T\delta^{-1}} \mathbf{E}|X''(s)| \le C_4(\beta, \lambda_0, T)\Big(|x| + |v| + \Omega^{1/2}\Big) \ , \tag{7.15}$$

using the a priori bounds (4.37), and C_4 is monotone in T.

For the continuity result, notice that $Y(t) := X(t) - \widetilde{X}(t)$ satisfies the equation,

$$Y''(t) + I\lambda^2 Y'(t) + a^2 Y(t) = I\lambda^2 X'(t) - (M \star X')(t) - xM(t) \ , \tag{7.16}$$

with initial conditions $Y(0) = Y'(0) = 0$. Using (7.3) we obtain,

$$\begin{aligned} \Big|I\lambda^2 X'(s) - (M \star X')(s)\Big| \ \ &\le \ \ \lambda^2\Big|\int_0^s \frac{\sin\Omega u}{u}\Big(X'(s) - X'(s-u)\Big) \, du\Big| \\ &\quad + \lambda^2 \ |X'(s)| \ \Big|\int_s^\infty \frac{\sin\Omega u}{u} \, du\Big| \ . \end{aligned} \tag{7.17}$$

The second term is estimated by $(const)\lambda^2|X'(s)|$ with a universal constant if $s \le 1$ and by $(const)\lambda^2(\Omega s)^{-1}|X'(s)| \le (const)\lambda^2\Omega^{-1}|X'(s)|$ if $s \ge 1$.

In the first term we split the integration domain. For $u \geq \Omega^{-2/3}$ we use integration by parts, (4.37) and (7.15)

$$\lambda^2 \, \mathbf{E} \, \Big| \int_{\Omega^{-2/3}}^{s} \frac{d}{du}\Big(\frac{\cos \Omega u}{\Omega}\Big) u^{-1} \Big(X'(s) - X'(s-u) \Big) \, du \Big| \leq C_5(\beta, \lambda_0, T) \delta |\log \delta| \Omega^{-1/3} \Big(1 + |x| + |v| \Big)$$

for all $s \leq T\delta^{-1}$. For the domain $0 \leq u \leq \Omega^{-2/3}$, we use Taylor expansion: $|X'(s) - X'(s-u)| \leq |u| \sup_{\sigma \leq s} |X''(\sigma)|$ and the bound (7.15). We obtain finally, using (4.37),

$$\mathbf{E} \, \Big| I\lambda^2 X'(s) - (M \star X')(s) \Big| \;\; \leq \;\; C_6(\beta, \lambda_0, T, x, v) \delta |\log \delta| \Omega^{-1/6} \; , \tag{7.18}$$

if $1 \leq s \leq T\delta^{-1}$ and

$$\mathbf{E} \, \Big| I\lambda^2 X'(s) - (M \star X')(s) \Big| \leq \pi \lambda_0^2 \delta F(t) \leq C_7(\beta, \lambda_0, x, v) \delta \Big(1 + |\log \delta| \Omega^{-1/6} \Big) \; , \tag{7.19}$$

if $s < 1$.

We now introduce the two fundamental solutions φ and ψ of $Y'' + I\lambda^2 Y' + a^2 Y = 0$ with $\varphi(0) = 0$, $\varphi'(0) = 1$ and $\psi(0) = 1, \psi'(0) = 0$. They are explicitly given as,

$$\varphi(t) = b^{-1} e^{-I\lambda^2 t/2} \sin bt \; , \qquad\qquad \psi(t) = e^{-I\lambda^2 t/2} \cos bt + \frac{I\lambda^2}{2} \varphi(t) \; , \tag{7.20}$$

with $b := (a^2 - I^2\lambda^4/4)^{1/2}$. Note that they are bounded functions for small enough δ. Hence, by (7.14), (7.18) and (7.19),

$$\begin{aligned} \mathbf{E} \, |Y(t)| \;\; &= \;\; \mathbf{E} \, \Big| \int_0^t \varphi(t-s) \Big(I\lambda^2 X'(s) - (M \star X')(s) - xM(s) \Big) \, ds \Big| \qquad (7.21) \\ &\leq \;\; \Big(C_8(\beta, \lambda_0, T, x, v) |\log \delta| \Omega^{-1/6} + C_7(\beta, \lambda_0, x, v) \delta + 2\lambda^2 |x| \Big[1 + (\log \Omega t)_+ \Big] \Big) \|\phi\|_\infty \\ &\leq \;\; C_9(\beta, \lambda_0, T, x, v) \Omega^{-1/6} |\log \delta| \; . \end{aligned}$$

The constants C_8 and C_9 can be chosen monotone in T, so the same estimate is valid for $\sup_{s \leq T\delta^{-1}} \mathbf{E} \, |Y(s)|$. The argument for Y' is similar, which proves (7.5). ■

7.2 Transport equation before scaling limits

Armed with (7.6), it is enough to compute $\widetilde{w}_A^\varepsilon(t, x, v)$. The calculation is the same as in Section 5.2 except for the different fundamental solutions φ and ψ given in (7.20). We redefine,

$$\begin{aligned} \eta_{\theta,\sigma} \;\; &:= \;\; \theta\varphi(t) + \sigma\varphi'(t) \; , \qquad (7.22) \\ \xi_{\theta,\sigma} \;\; &:= \;\; \theta\psi(t) + \sigma\psi'(t) \; , \end{aligned}$$

and in complete analogy to Lemma 5.2 we state the,

Lemma 7.2 *We have for $t \geq 0$,*

$$\int_{\mathbb{R}^2} \widetilde{w}_A^\varepsilon(t,x,v)\overline{\phi(x,v)}\,dx\,dv = \int_{\mathbb{R}^2} \widehat{w}_0\Big(\xi_{\theta,\sigma}(t),\eta_{\theta,\sigma}(t)\Big)\overline{\widehat{\phi}(\theta,\sigma)}e^{-\frac{1}{2}Q(t)}\,d\theta\,d\sigma\,, \tag{7.23}$$

with

$$Q(t) := \lambda^2 \int_0^\Omega A_\beta^2(\omega)H(t,\omega)d\omega\,, \tag{7.24}$$

and H is given again as $H(t,\omega) = \left|\int_0^t \eta_{\theta,\sigma}(s)e^{-is\omega}ds\right|^2$, but with the new $\eta_{\theta,\sigma}$ defined in (7.22). We also have exactly the same estimate as (5.16), but with the redefined quantities. ■

7.3 Obtaining diffusion from scaling limit

In this section, and with similar arguments as in Section 6, we again obtain linear dependence in time of $Q(t)$ for large t. Indeed, we first write,

$$\varphi(t) = \frac{1}{2ib}\Big(e^{tu} - e^{t\bar{u}}\Big)\,, \qquad \text{with,} \quad u := -\frac{I\lambda^2}{2} + ib\,. \tag{7.25}$$

With these notations, we have,

$$\eta_{\theta,\sigma}(t) = \frac{1}{2ib}\bigg(\theta\Big(e^{tu} - e^{t\bar{u}}\Big) + \sigma\Big(ue^{tu} - \bar{u}e^{t\bar{u}}\Big)\bigg)\,, \tag{7.26}$$

hence,

$$H(t,\omega) \;=\; \frac{1}{4b^2}\left|(\theta+\sigma u)\frac{e^{t(u-i\omega)}-1}{u-i\omega} - (\theta+\sigma\bar{u})\frac{e^{t(\bar{u}-i\omega)}-1}{\bar{u}-i\omega}\right|^2\,. \tag{7.27}$$

We now take the scaling $t = T\delta^{-1}$ for a fixed T and $\delta \to 0$. The terms with denominator $\bar{u} - i\omega = -I\lambda^2/2 - i(\sqrt{a^2 - I^2\lambda^4/4} + \omega)$ have no singularity (they are bounded) so the first term of H is the main term. Extracting the main term, we can write (cf. (5.20)),

$$H(t,\omega) = (\theta^2 + a^2\sigma^2)\left[\frac{1}{4a^2}\left|\frac{e^{t(u-i\omega)}-1}{u-i\omega}\right|^2 + U(t,\omega)\right]\,.$$

Using $u = ai + O(\delta)$, $3/4 \leq a^2 \leq 1$, $b^2 = a^2 + O(\delta^2)$ we obtain for small enough δ that,

$$\int_0^\infty |U(T\delta^{-1},\omega)|d\omega \leq C_{10}(\beta,\lambda_0,T)|\log\delta|\,.$$

With some elementary calculations this implies,

$$\begin{aligned} Q(T\delta^{-1}) \;&=\; \lambda^2(\theta^2+a^2\sigma^2)\left[\frac{1}{4a^2}\int_0^\Omega A_\beta^2(\omega)\left|\frac{e^{T\delta^{-1}(u-i\omega)}-1}{u-i\omega}\right|^2 d\omega + B_1(T\delta^{-1})\right] \\ &=\; \lambda^2(\theta^2+a^2\sigma^2)\left[\frac{A_\beta^2(a)}{4a^2}\int_{a+\sqrt{\delta}}^{a-\sqrt{\delta}}\left|\frac{e^{T\delta^{-1}(u-i\omega)}-1}{u-i\omega}\right|^2 d\omega + B_3(T\delta^{-1})\right]\,, \end{aligned} \tag{7.28}$$

where the functions B_j $(j = 1, 2, 3)$ satisfy $|B_j(T\delta^{-1})| \leq C_{11}(\beta, \lambda_0, T)\delta^{-1/2}$. We used that the function $\omega \mapsto A_\beta^2(\omega)$ is bounded with a bounded derivative around $\omega \sim a$, and that the function $z \mapsto (e^{tz} - 1)/z$ is uniformly bounded by t in the vicinity of the imaginary axis.

Since the derivative of $z \mapsto |(e^{tz} - 1)/z|^2$ is bounded by t^2, one can replace u by ai in the last integral at the expense of an error $2\sqrt{\delta}|u - ia|t^2 = O(\delta^{-1/2})$. Finally one can evaluate,

$$\int_{a+\sqrt{\delta}}^{a-\sqrt{\delta}} \left| \frac{e^{T\delta^{-1}(a-\omega)i} - 1}{a - \omega} \right|^2 d\omega = 2\pi T\delta^{-1} + O(\delta^{-1/2})$$

At this step $T \geq 0$ is used. In summary, we obtained,

$$Q(T\delta^{-1}) = (\theta^2 + a^2\sigma^2)\Big(\lambda_0^2 T \, \frac{\pi(\cosh(\beta a) + 1)}{4a \sinh \beta a} + B_4(T\delta^{-1})\Big) \, . \tag{7.29}$$

The error satisfies $|B_4(T\delta^{-1})| \leq C_{12}(\beta, \lambda_0, T)\delta^{1/2}$, hence,

$$\lim_{\delta \to 0} Q(T\delta^{-1}) = c_\beta \lambda_0^2 \gamma^2 T \, , \tag{7.30}$$

with $\gamma := \theta^2 + \check{a}^2\sigma^2$ and

$$c_\beta := \frac{\pi(\cosh(\beta\check{a}) + 1)}{4\check{a} \sinh \beta\check{a}} \, , \tag{7.31}$$

assuming that

$$\check{a} := \lim_{\delta \to 0, \Omega \to \infty} a \; = \lim_{\delta \to 0, \Omega \to \infty} (1 - \lambda_0 \Omega \delta^{-1}) \tag{7.32}$$

exists, and $\check{a} \in [\frac{1}{4}, 1]$.

Since we will keep β fixed and choose $\lambda = \lambda_0 \delta^{1/2}$ with a fixed λ_0, δ and Ω are left as a scaling parameters from the triple $\varepsilon = (\beta, \Omega, \lambda)$. Like in Section 6 (cf.(6.2)) we introduce,

$$W_T^\varepsilon(x, v) := w_A^\varepsilon(T\delta^{-1}, x, v) \, , \tag{7.33}$$

and notice that only the time is rescaled. We will assume that $\Omega \to \infty$ along with $\delta \to 0$ in such a way that the limit (7.32) exists and $\Omega \in [|\log \delta|^7, \delta^{-1}]$. Clearly either $\Omega \sim \delta^{-1}$, in which case $\check{a} < 1$, or $\Omega \ll \delta^{-1}$, when $\check{a} = 1$. In the latter case, however, we need $\Omega \geq |\log \delta|^7$.

7.4 Derivation of the limiting equation

We need the notion of "radial" function with respect to the elliptical phase space trajectories of the oscillator $Y'' + \check{a}^2 Y$. As usual, the dual variables to the phase space coordinates (x, v) are (θ, σ). With $\check{a} > 0$ fixed, let

$$\gamma = \gamma(\theta, \sigma) := \sqrt{\theta^2 + \check{a}^2\sigma^2} \, , \qquad r = r(x, v) := \sqrt{x^2 + \check{a}^{-2}v^2} \, ,$$

which will be considered either variables or functions, depending on the context. If a function $u(x,v)$ depends only on $x^2 + \check{a}^{-2}v^2$, then it can be written as $u(x,v) = u^*(r)$ with some function u^* defined on $\mathbb{R}_+$. Then the *two dimensional* Fourier transform $\widehat{u}(\theta,\sigma) = \int \exp[-i(\theta x + \sigma v)]u(x,v)dxdv$ is a function of $\theta^2 + \check{a}^2\sigma^2$ only, hence it can be written as $\widehat{u}(\theta,\sigma) = \widetilde{u}^*(\gamma)$. Here $\widetilde{u}^*$ can be thought of as the "elliptical-radial" Fourier transform of u^*, but in order to avoid confusion, we will always perform Fourier transforms on $\mathbb{R}^2$, i.e. between $u(x,v) \leftrightarrow \widehat{u}(\theta,\sigma)$, even if these functions are "radial".

For any function $u(x,v)$ we can form the "radial" average of its Fourier transform $\widehat{u}(\theta,\sigma)$ by defining

$$\widehat{u}^{\#}(\theta,\sigma) := \frac{1}{2\pi}\int_0^{2\pi} \widehat{u}(\gamma\cos s, \check{a}^{-1}\gamma\sin s)ds \qquad \left(= \frac{1}{2\pi\gamma}\int_{\tilde{\theta}^2+\check{a}^2\tilde{\sigma}^2=\gamma^2} \widehat{u}(\tilde{\theta},\tilde{\sigma})d\tilde{\theta}d\tilde{\sigma} \right),$$

which is a function of γ, hence it can be written as

$$\widehat{u}^{\#}(\theta,\sigma) = \widetilde{u}^{\#,*}(\gamma)\ .$$

In this notation # refers to "radial" averaging, and $*$ indicates that we consider the radial part of the function. Tilde indicates that it comes from the two dimensional Fourier transform $\widehat{u}$ of the original function u.

Theorem 7.1 *Define the large time scale Wigner function $W_T^\varepsilon(x,v)$ as in (7.33). Assume that $\lambda = \lambda_0\delta^{1/2}$, $\lambda_0 \le \sqrt{3}/2$ and fix $\beta > 0$, $\check{a} \in [\frac{1}{2}, 1]$. The initial condition $W_0^\varepsilon(x,v) = w_0(x,v)$ satisfies $\widehat{w}_0(\theta,\sigma) \in L^1(\mathbb{R}_\theta \times \mathbb{R}_\sigma)$. Consider the "radial" average of $\widehat{W}_T^\varepsilon$,*

$$\widetilde{W}_T^{\#,\varepsilon}(\gamma) := \frac{1}{2\pi}\int_0^{2\pi} \widehat{W}_T^\varepsilon(\gamma\cos s, \check{a}^{-1}\gamma\sin s)ds\ . \tag{7.34}$$

Then for any $T \ge 0$ the limit,

$$\widehat{W}_T^+(\theta,\sigma) := \lim_{\substack{\delta\to 0,\Omega\to\infty \\ 1-\lambda_0^2\Omega\delta\to\check{a} \\ \Omega\ge|\log\delta|^7}} \widehat{W}_T^{\#,\varepsilon}(\theta,\sigma)\ , \tag{7.35}$$

exists in a weak sense and it is a function of $\gamma = (\theta^2 + \check{a}^2\sigma^2)^{1/2}$ only. Hence, its inverse Fourier transform $W_T^+(x,v)$ is a function of $r = (x^2 + \check{a}^{-2}v^2)^{1/2}$ only and it can be written as $W_T^{+,*}(r) := W_T^+(x,v)$. This function satisfies the "radial" Fokker-Planck equation,

$$\partial_T W_T^{+,*} = \frac{\pi\lambda_0^2}{4}\,\partial_r(rW_T^{+,*}) + \frac{c_\beta\lambda_0^2}{2}\,\Delta_r W_T^{+,*}\ , \tag{7.36}$$

(c_β is given in (7.31)) with initial condition $W_0^{+,*}(r) := W_{T=0}^+(x,v)$ whose Fourier transform $\widehat{W}_0^+(\theta,\sigma)$ is given by,

$$\widehat{W}_0^+(\theta,\sigma) := \widehat{w}_0^{\#}(\theta,\sigma) = \frac{1}{2\pi}\int_0^{2\pi} \widehat{w}_0(\gamma\cos s, \check{a}^{-1}\gamma\sin s)ds\ . \tag{7.37}$$

Remark 1. The weak limit $\lim^{**}\widehat{W}_T^\varepsilon(\theta,\sigma)$ (without averaging over the angular variables) does not exist (here $\lim^{**}$ stands for the same limit as in (7.35)). However, time averaging can again replace angular averaging (see Remark and Corollary 6.1), i.e. our method easily proves that $\lim^{**}W_T^\varepsilon(x,v)$ exists in a weak sense in all variables (x,v,T), i.e. in the topology of $\mathcal{D}'(\ \mathbb{R}_x\times\mathbb{R}_v\times[0,\infty)_T\ )$, and it satisfies (7.36) weakly in space, velocity and time.

Remark 2. Since the diffusion coefficient $\frac{1}{2}\lambda_0^2 c_\beta$ in (7.36) behaves as β^{-1} for small β (high temperature), we see that Einstein's relation is satisfied at high temperatures. At small temperatures the diffusion does not disappear ($\lim_{\beta\to\infty} c_\beta > 0$), which is due to the ground state quantum fluctuations of the heat bath.

Proof. The proof is similar to the proof of Theorem 6.1, hence we skip certain steps. Let $\phi(x,v)\in C_0^\infty(\mathbb{R}\times\mathbb{R})$. Similarly to (6.14) we obtain from (4.41),

$$\begin{aligned}\langle W_T^\varepsilon,\phi\rangle &= \int \widehat{w}_A^\varepsilon(T\delta^{-1},\theta,\sigma)\overline{\widehat{\phi}(\theta,\sigma)}d\theta\ d\sigma \qquad (7.38)\\ &= \mathbf{E}\int \widehat{w}_0(\xi,\eta)\overline{\widehat{\phi}(\theta,\sigma)}e^{i(x\xi+v\eta)}e^{-i(\theta X(t)+\sigma X'(t))}d\xi\ d\eta\ dx\ dv\ d\theta\ d\sigma\ .\end{aligned}$$

Thanks to (7.6), in the limit $\delta\to 0$ we can replace X by $\widetilde{X}$ and to take the limiting value (7.30) of Q in the formulae (we again have to approximate $\widehat{w}_0$ by bounded functions first). We obtain (cf. (6.15)),

$$\begin{aligned}\lim^{**}\langle W_T^\varepsilon,\phi\rangle &= \lim^{**}\mathbf{E}\int \widehat{w}_0(\xi,\eta)\overline{\widehat{\phi}(\theta,\sigma)}e^{i(x\xi+v\eta)}e^{-i(\theta\widetilde{X}(T\delta^{-1})+\sigma\widetilde{X}'(T\delta^{-1}))}d\xi\ d\eta\ dx\ dv\ d\theta\ d\sigma\\ &= \lim^{**}\int \widehat{w}_0\Big(\xi_{\theta,\sigma}(T\delta^{-1}),\eta_{\theta,\sigma}(T\delta^{-1})\Big)\overline{\widehat{\phi}(\theta,\sigma)}e^{-\frac{1}{2}Q(T\delta^{-1})}d\theta\ d\sigma \qquad (7.39)\end{aligned}$$

where $\lim^{**}$ stands for the limit in (7.35). Recall that the functions $\xi_{\theta,\sigma}$ and $\eta_{\theta,\sigma}$ now depend on the limiting parameters, since φ and ψ do, and they are oscillating, which again prevents the existence of the weak limit in the last line of (7.39) without averaging.

Time averaging is analogous to part b) of Theorem 6.1, and it gives the weak limit in space, velocity and time. We skip the details of the proof of the statement of Remark 1.

Performing a radial avegaring (with respect to the limiting ellipses given by the level curves of $r=r(x,v)$ or $\gamma=\gamma(\theta,\eta)$) is the same as using "radial" testfunctions ϕ which depend only on r; i.e. $\widehat{\phi}(\theta,\sigma)$ depends only on γ hence it can be written as $\widehat{\phi}(\theta,\sigma)=\widetilde{\phi}^*(\gamma)$. In this case

$$\langle\widehat{W}_T^{\#,\varepsilon},\widehat{\phi}\rangle = \langle\widehat{W}_T^\varepsilon,\widehat{\phi}\rangle\ .$$

From the explicit formulas (7.20), (7.22) it is straightforward to check that

$$\lim^{**} \sup_{s\leq T\delta^{-1}} \left| \Big([\xi_{\theta,\sigma}(s)]^2 + \check{a}^2 [\eta_{\theta,\sigma}(s)]^2 \Big) - e^{-I\lambda_0^2 s\delta} \Big([\check{\xi}_{\theta,\sigma}(s)]^2 + \check{a}^2 [\check{\eta}_{\theta,\sigma}(s)]^2 \Big) \right| = 0 \ , \qquad (7.40)$$

where $\check{\xi}$ and $\check{\eta}$ are the solutions to $Y'' + \check{a}^2 Y = 0$, i.e.

$$\check{\xi}_{\theta,\sigma}(s) := \theta \cos(\check{a}s) - \sigma \check{a} \sin(\check{a}s) \ , \qquad \check{\eta}_{\theta,\sigma}(s) := \theta \check{a}^{-1} \sin(\check{a}s) + \sigma \cos(\check{a}s) \ .$$

Since the flow $(\theta,\sigma) \mapsto \Big(\xi_{\theta,\sigma}(s), \eta_{\theta,\sigma}(s) \Big)$ is measure preserving, one can change variables

$$\int_{\mathbb{R}^2} \widehat{w}_0 \Big(\xi_{\theta,\sigma}(t), \eta_{\theta,\sigma}(t) \Big) \overline{\widehat{\phi}(\theta,\sigma)} e^{-\frac{1}{2}Q(t)} \, d\theta \, d\sigma \qquad (7.41)$$
$$= \int_{\mathbb{R}^2} \widehat{w}_0(\theta,\sigma) \overline{\widehat{\phi} \Big(\xi^*_{\theta,\sigma}(t), \eta^*_{\theta,\sigma}(t) \Big)} e^{-\frac{1}{2}Q^*(t)} \, d\theta \, d\sigma \ ,$$

where $\eta^*(t) := \eta(-t)$, $\xi^*(t) := \xi(-t)$ are the backward trajectories. In this way we pushed the trajectories into the argument of $\widehat{\phi}$, where only their $\xi^2 + \check{a}^2\eta^2$ combination matters, and we can apply (7.40) to replace ξ, η by $\check{\xi}, \check{\eta}$, finally we can change variables backwards, now along these new trajectories.

Hence together with (7.30) and with $c'_\beta := c_\beta/2$ for simplicity, we have

$$\begin{aligned} \lim^{**} \langle \widehat{W}_T^{\#,\varepsilon}, \widehat{\phi} \rangle &= \lim^{**} \langle \widehat{W}_T^{\varepsilon}, \widehat{\phi} \rangle \\ &= \lim^{**} \int_{\mathbb{R}^2} \widehat{w}_0 \Big(e^{-I\lambda_0^2 T/2} \check{\xi}_{\theta,\sigma}(T\delta^{-1}) \, , e^{-I\lambda_0^2 T/2} \check{\eta}_{\theta,\sigma}(T\delta^{-1}) \Big) \overline{\widetilde{\phi}^*(\gamma)} e^{-c'_\beta \lambda_0^2 T \gamma^2} d\theta \, d\sigma \ , \end{aligned}$$

if we can show that this latter limit exists. But the right hand side above is in fact independent of the limiting parameters δ, Ω, since we can first integrate on ellipses $\theta^2 + \check{a}^2\sigma^2 = (const)$, similarly to the same calculation in the proof of part c), Theorem 6.1. Hence,

$$\int_{\mathbb{R}^2} \widehat{w}_0 \Big(e^{-I\lambda_0^2 T/2} \check{\xi}_{\theta,\sigma}(T\delta^{-1}) \, , e^{-I\lambda_0^2 T/2} \check{\eta}_{\theta,\sigma}(T\delta^{-1}) \Big) \overline{\widetilde{\phi}^*(\gamma)} e^{-c'_\beta \lambda_0^2 T \gamma^2} d\theta \, d\sigma \qquad (7.42)$$
$$= \int_{\mathbb{R}^2} \widetilde{W}_0^{+,*} \Big(\gamma e^{-I\lambda_0^2 T/2} \Big) \overline{\widetilde{\phi}^*(\gamma)} e^{-c'_\beta \lambda_0^2 T \gamma^2} d\theta \, d\sigma \ ,$$

where we recall the definition of $\widetilde{W}_0^+$ (7.37), which depends only on $\gamma^2 = \theta^2 + \check{a}^2\sigma^2$, and we let $\widetilde{W}_0^{+,*}(\gamma) := \widetilde{W}_0^+(\theta,\sigma)$. Therefore, the relation,

$$\lim^{**} \langle \widehat{W}_T^{\#,\varepsilon}, \widehat{\phi} \rangle = \int_{\mathbb{R}^2} \widetilde{W}_0^{+,*} \Big(\gamma e^{-I\lambda_0^2 T/2} \Big) \overline{\widetilde{\phi}^*(\gamma)} e^{-c'_\beta \lambda_0^2 T \gamma^2} d\theta \, d\sigma$$

defines the weak limit,

$$\widehat{W}_T^+(\theta,\sigma) := \lim^{**} \widehat{W}_T^{\#,\varepsilon}(\theta,\sigma)$$

and it is a function depending only on $\theta^2 + \check{a}^2\sigma^2$, i.e. it can be written as $\widetilde{W}_T^{+,*}(\gamma) := \widehat{W}_T^+(\theta,\sigma)$. Also, we readily obtain the equation satisfied by $\widetilde{W}_T^{+,*}(\gamma)$ by computing,

$$\begin{aligned} \left\langle \partial_T\Big|_{T=0} \widehat{W}_T^+, \widehat{\phi} \right\rangle &= \partial_T\Big|_{T=0} \int_{\mathbb{R}^2} \widetilde{W}_0^{+,*}\Big(\gamma e^{-I\lambda_0^2 T/2}\Big)\overline{\widetilde{\phi}^*(\gamma)} e^{-c'_\beta \lambda_0^2 T\gamma^2} d\theta\, d\sigma \qquad (7.43) \\ &= \int_{\mathbb{R}^2} \Big[-\frac{I\lambda_0^2}{2}\gamma\partial_\gamma - c'_\beta\lambda_0^2\gamma^2 \Big] \widetilde{W}_0^{+,*}(\gamma)\overline{\widetilde{\phi}^*(\gamma)} d\theta\, d\sigma \ , \end{aligned}$$

from which (7.36) follows, recalling that $I = \frac{\pi}{2}$ and the value of $c'_\beta = c_\beta/2$ from (7.31). ■

Acknowledgements. The authors are indebted to H. Spohn for discussions. F.C. and L.E. were partially supported by the Erwin Schrödinger Institute in Vienna (Austria) during their visit, and they thank this institution for its hospitality. This work was supported by the TMR-Network "Asymptotic Methods in Kinetic Theory" number ERB FMBX CT97 0157 (F.C., F.F. and P.A.M.) and by NSF-grant DMS-9970323 (L.E.).

References

[1] A.Arnold, J. L. Lopez, P. A. Markowich and J. Soler, *An Analysis of Quantum Fokker-Planck Models: A Wigner Function Approach*, preprint, 1999

[2] A. Arai, *Long-time behaviour of an electron interacting with a quantized radiation field*, J. Math. Phys., Vol. 32, N. 8, p. 2224-2242 (1991).

[3] C, Boldrighini, L. Bunimovich, Y.Sinai, *On the Boltzmann equation for the Lorentz gas*, J. Stat. Phys. Vol. 32, p. 477-501, (1983).

[4] N. Ben Abdallah, P. Degond, *On a hierarchy of macroscopic models for semi-conductors*, J. Math. Phys., Vol. 37, N. 7, p. 3306-3333 (1996).

[5] A.O. Caldeira, A.J. Leggett, *Path-integral approach to quantum Brownian motion*, Physica A, Vol. 121, 587-616 (1983). Erratum: ibid. Vol. 130, p. 374 (1985).

[6] A.O. Caldeira, A.J. Leggett, *Quantum tunnelling in a dissipative system*, Ann. Phys., Vol. 149, p. 374-456 (1983).

[7] F. Castella, *On the derivation of a Quantum Boltzmann Equation from the periodic Von-Neumann equation*, Mod. Math. An. Num., Vol. 33, N. 2, p. 329-349 (1999).

[8] F. Castella, P. Degond, *From the Von-Neumann equation to the quantum Boltzmann equation in a deterministic framework*, C. R. Acad. Sci., t. 329, sér. I, p. 231-236 (1999) and Preprint (1999).

[9] Y.-C. Chen, J. L. Lebowitz and C. Liverani, *Dissipative quantum dynamics in a boson bath*, Phys. Rev. B, Vol 40, p. 4664-4682 (1989).

[10] C. Cohen-Tannoudji, B. Diu, F. Laloë, **Mécanique Quantique**, I et II, Enseignement des Sciences, Vol. 16, Hermann (1973).

[11] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, **Processus d'interaction entre photons et atomes**, "Savoirs actuels", Intereditions/Editions du CNRS (1988).

[12] H. Dekker, *Multilevel tunneling and coherence: dissipative spin-hopping dynamics at finite temperatures*, Phys. Rev. A, Vol. 44, N. 4, p. 2314-2323 (1991).

[13] P. de Smedt, D. Dürr, J. L. Lebowitz and C. Liverani, *Quantum system in contact with a thermal environment: rigorous treatment of a simple model*, Commun. Math. Phys., Vol. 120, p. 195-231 (1988).

[14] L. Diosi, *On High-Temperature Markovian Equation for Quantum Brownian Motion*, Europhys. Lett., Vol. 22, N. 1, p. 1-3 (1993).

[15] L. Diosi, *Caldeira-Leggett master equation and medium temperatures*, Phys. A, Vol. 199, p. 517-526 (1993).

[16] L. Diosi, N. Gisin, J. Haliwell, I.C. Percival, *Decoherent Histories and Quantum State Diffusion*, Phys. Rev. Lett., Vol. 74, N. 2 p. 203-207 (1995).

[17] D. Dürr, S. Goldstein and J. Lebowitz, *Asymptotic motion of a classical particle in a random potential in two dimensions: Landau model*, Commun. Math. Phys. Vol. 113, p. 209-230 (1987).

[18] L. Erdős, H.T. Yau, *Linear Boltzmann equation as scaling limit of quantum Lorenz gas*, Advances in Differential Equations and Mathematical Physics. Contemporary Mathematics Vol. 217, p. 137-155 (1998).

[19] L. Erdős, H.T. Yau, *Linear Boltzmann equation as the weak coupling limit of a random Schrödinger equation*, to appear in Comm. Pure Appl. Math. (1999).

[20] L. Erdős, H.T. Yau, *Linear Boltzmann equation for electron transport in a weakly coupled phonon field*, In preparation

[21] R. Esposito, M. Pulvirenti, A. Teta, *The Boltzmann Equation for a one-dimensional Quantum Lorentz gas*, Preprint (1998).

[22] R.P. Feynman, A. Hibbs, **Quantum Mechanics and Path Integrals**, McGraw Hill, New York (1965).

[23] R.P. Feynman, F.L. Vernon, *The theory of a general quantum system interacting with a linear dissipative system*, Ann. Phys., Vol. 24, p. 118-173 (1963).

[24] W. Fischer, H. Leschke and P. Müller, *On the averaged quantum dynamics by white-noise Hamiltonian with and without dissipation*, Ann. Physik, Vol. 7, p. 59-100 (1998).

[25] G. Gallavotti, *Rigorous theory of the Boltzmann equation in the Lorentz gas*, Nota inteerna n. 358, Univ. di Roma (1970).

[26] P. Gérard, P.A. Markowich, N. Mauser, F. Poupaud. *Homogenization limits and Wigner transforms*, Comm. Pure Appl. Math., Vol. L, p. 323 - 379 (1997).

[27] Z. Haba, *Classical limit of quantum dissipative systems*, Lett. Math. Phys., Vol 44, p. 121-130 (1998).

[28] T.G. Ho, L.J. Landau, A.J. Wilkins, *On the weak coupling limit for a Fermi gas in a random potential*, Rev. Math. Phys., Vol. 5, N. 2, p. 209-298 (1993).

[29] F. Haake, R. Reibold, *Strong-Damping and low-temperature anomalies for the harmonic oscillator*, Phys. Rev. A, Vol. 32, N. 4, p. 2462-2475 (1985).

[30] V. Jakšić, C.-A. Pillet, *Spectral theory of thermal relaxation*, J. Math. Phys. Vol. 38, p. 1757-1780 (1997)

[31] A. M. Il'in, R. Z. Kas'minskii, *On equations of Brownian motion*, Theory Probab. Appl. (USSR), Vol. 9, p. 421-444 (1964).

[32] J.B. Keller, G. Papanicolaou and L. Ryzhik, *Transport equations for elastic and other waves in random media*, Wave Motion, Vol. 24, N. 4, p. 327-370 (1996).

[33] H. Kesten and G. Papanicolaou, *A limit theorem for stochastic acceleration,* Comm. Math. Phys. Vol. 78, p. 19-63 (1980).

[34] L.J. Landau, *Observation of Quantum Particles on a Large Space-Time Scale*, J. Stat. Phys., Vol. 77, N. 1-2, p. 259-309 (1994).

[35] O. E. Lanford, *On a derivation of the Boltzmann equation*, Astérisque, Vol. 40, p. 117–137 (1976)

[36] E. M. Lifshitz and L. P. Pitaevskii, **Physical Kinetics**, Vol. 10 of Course of Theoretical Physics by L. D. Landau and E. M. Lifshitz, Pergamon Press (1981).

[37] G. Lindblad, *On generators of quantum dynamical semigroups*, Comm. Math. Phys, Vol. 48, p. 119-130 (1999).

[38] P.L. Lions, Th. Paul, *Sur les measures de Wigner*, Revista Mat. Iberoamericana, Vol. 9, p. 533-618 (1993).

[39] P. A. Markowich, C. A. Ringhofer and C. Schmeiser, *Semiconductor Equations*, Springer (1990).

[40] F. Nier, *Asymptotic Analysis of a scaled Wigner equation and Quantum Scattering*, Transp. Theor. Stat. Phys., Vol. 24, N. 4 et 5, p. 591-629 (1995).

[41] F. Nier, *A semi-classical picture of quantum scattering*, Ann. Sci. Ec. Norm. Sup., 4. Sér., t. 29, p. 149-183 (1996).

[42] J.A. Reissland, **The Physics of Phonons**, Wiley Interscience, John Wiley and Sons ltd. (1972).

[43] H. Spohn, *Derivation of the transport equation for electorns moving through random impurities*, J. Stat. Phys., Vol. 17, p. 385-412 (1977).

[44] H. Spohn, *Kinetic equations from Hamiltonian dynamics: Markovian limits*, Rev. Mod. Phys., Vol. 53, N. 3, p.569-615 (1980).

[45] H. Spohn, **Large Scale Dynamics of interacting particles**, Springer, Berlin (1991).

[46] W.G. Unruh, W.H. Zurek, *Reduction of a wave packet in quantum Brownian motion*, Phys. Rev. D, Vol. 40, N. 4, p. 1071-1094 (1989).

[47] E.P. Wigner, *On the Quantum Correction for Thermodynamic Equilibrium*, Phys. Rev. Vol. 40, p. 749-759 (1932).

E-mail addresses: castella@maths.univ-rennes1.fr, lerdos@math.gatech.edu, Florian.Frommlet@esi.ac.at, marko@aurora.tuwien.ac.at